

Asymptotic eigenvalue estimates for a Robin problem with a large parameter

Pavel Exner, Alexander Minakov, and Leonid Parnovski*

Abstract. Robin problem for the Laplacian in a bounded planar domain with a smooth boundary and a large parameter in the boundary condition is considered. We prove a two-sided three-term asymptotic estimate for the negative eigenvalues. Furthermore, improving the upper bound we get a two term asymptotics in terms of the coupling constant and the maximum of the boundary curvature.

Mathematics Subject Classification (2010). 35P15, 35J05.

Keywords. Laplacian, Robin problem, eigenvalue asymptotics.

1. Introduction and the main result

Asymptotic properties of eigenvalues belong among the most often studied problems in the spectral theory. In this paper we are going to discuss an asymptotics of the “attractive” Robin problem for the Laplacian in a bounded domain of \mathbb{R}^2 in the situation when the parameter β in the boundary condition assumes large values. The problem has a physical motivation; it naturally arises in the study of reaction-diffusion equation where a distributed absorption competes with a boundary source – see [5], [6] for details. At the same the question is of mathematical interest. In a recent paper, Levitin and Parnovski [7] investigated the asymptotic behavior of the principal eigenvalue and showed that its leading term is $-c\beta^2$ where $c = 1$ if the domain boundary is smooth and $c > 1$ if it has angles. The same one-term asymptotics is known to hold in the former case also for higher eigenvalues [1].

A related asymptotic problem is encountered in the theory of leaky quantum graphs [2] where the dynamics is not constrained to a bounded region, instead

*The research has been supported by the projects “Support of inter-sectoral mobility and quality enhancement of research teams at Czech Technical University in Prague”, CZ.1.07/2.3.00/30.0034, sponsored by European Social Fund in the Czech Republic, and 14-06818S of the Czech Science Foundation. A.M. expresses his gratitude to Stepan Manko and Satoshi Ohya for useful discussions.

it is governed by a singular Schrödinger operator with an attractive interaction supported by a manifold or complex of a lower dimension. A particularly close analogy occurs in the two-dimensional situation when the interaction support is closed smooth loop; using a combination of bracketing and estimates with separated variables, one is able to derive an asymptotic expansion of negative eigenvalues [4] in which the absolute term with respect to the coupling parameter is given by a one-dimensional Schrödinger operator with a potential determined by the geometry of the problem, specifically the curvature of the loop.

This inspires the question whether the technique used for the singular Schrödinger operators cannot be used also for Robin “billiards” with a smooth boundary. This is the main topic of the present paper. We are going to show first that in distinction to the Schrödinger case the method of [4] does not yield an asymptotic expansion, but two-sided asymptotic estimates only, which squeeze only when the domain is a circular disc. On the other hand, these estimates hold true not only for the principal eigenvalue, and moreover, they have three terms in the powers of β which improves, in particular, the result obtained in [7] for smooth boundaries. On the other hand, the result admits an improvement. Replacing the upper bound by a variational estimate similar to that employed recently by Pankrashkin [8] for the principal eigenvalue, we obtain a bound in which only the maximum of the boundary curvature appears, and as a result, a two-term asymptotic expansion.

Let us now state the problem properly. We suppose that Ω be an open, simply connected set in \mathbb{R}^2 with a closed C^4 Jordan boundary $\partial\Omega = \Gamma : [0, L] \ni s \mapsto (\Gamma_1, \Gamma_2) \in \mathbb{R}^2$ which is parametrized by its arc length; for definiteness we choose the clockwise orientation of the boundary. Let $\gamma : [0, L] \rightarrow \mathbb{R}$ be the signed curvature of Γ , i.e. $\gamma(s) = \Gamma_1''(s)\Gamma_2'(s) - \Gamma_2''(s)\Gamma_1'(s)$. We investigate the spectral boundary-value problem

$$\begin{aligned} -\Delta f &= \lambda f & \text{in } \Omega \\ \frac{\partial f}{\partial n} &= \beta f & \text{on } \Gamma \end{aligned} \tag{1}$$

with a parameter $\beta > 0$, which will be in the following assumed to be large; the symbol $\frac{\partial}{\partial n}$ in (1) denotes the outward normal derivative. It is straightforward to check that the quadratic form

$$q_\beta[f] = \|\nabla f\|_{L^2(\Omega)}^2 - \beta \int_{\Gamma} |f(x)|^2 ds \tag{2}$$

with $\text{Dom}(q_\beta) = H^1(\Omega)$ is closed and below bounded; we denote by H_β the unique self-adjoint operator associated with it. Our main goal is to study the asymptotic behavior of the negative eigenvalues of H_β as parameter β tends to infinity. To state the result, we introduce the one-dimensional Schrödinger operator

$$S = -\frac{d^2}{ds^2} - \frac{1}{4}\gamma^2(s) \quad \text{in } L^2(0, L) \tag{3}$$

with the domain

$$P = \{f \in H^2(0, L) : f(0) = f(L), f'(0) = f'(L)\}. \quad (4)$$

We use the symbol μ_j for the j -th eigenvalue of S counted with the multiplicity, $j \in \mathbb{N}$, and furthermore, we denote $\gamma^* = \max_{[0, L]} \gamma(s)$ and $\gamma_* = \min_{[0, L]} \gamma(s)$.

Our main result reads then as follows.

Theorem 1.1. *Under the stated assumptions, to any fixed integer n there exists a $\beta(n) > 0$ such that the number of negative eigenvalues of H_β is not smaller than n . For $\beta > \beta(n)$ we denote by $\lambda_n(\beta)$ the n -th eigenvalue of H_β counted with the multiplicity. Then $\lambda_n(\beta)$ satisfies for $\beta \rightarrow \infty$ the asymptotic estimates*

$$-\left(\beta + \frac{\gamma^*}{2}\right)^2 + \mu_n + \mathcal{O}\left(\frac{\log \beta}{\beta}\right) \leq \lambda_n(\beta) \leq -\left(\beta + \frac{\gamma_*}{2}\right)^2 + \mu_n + \mathcal{O}\left(\frac{\log \beta}{\beta}\right). \quad (5)$$

Remarks 1.2. (a) It will be clear from the proof that the assumption about simple connectedness of Ω is done mostly for the sake of simplicity. The result extends easily to multiply connected domains, in general with different parameters at different components of the boundary; each of the components then gives rise to a series of negative eigenvalues tending to $-\infty$ in the limit.

(b) In the light of the following result the upper bound in (5) is not of much use. We include it primarily to illustrate the significant difference between the “two-sided” situation discussed in [4] and the “one-sided” one treated here.

As we have indicated, the upper bound can be improved:

Theorem 1.3. *In the asymptotic regime $\beta \rightarrow +\infty$ the inequality*

$$\lambda_n(\beta) \leq -\beta^2 - \gamma^* \beta + \mathcal{O}(\beta^{2/3})$$

is valid for any fixed n . Consequently, the j -th eigenvalue behaves asymptotically as

$$\lambda_n(\beta) = -\beta^2 - \gamma^* \beta + \mathcal{O}(\beta^{2/3}).$$

Thus we obtain a two-term asymptotics which, in contrast to the Schrödinger operator case treated in [4], is not precise enough to distinguish between individual eigenvalues whose mutual distances are expected to be of order of $\mathcal{O}(1)$.

2. Proof of Theorem 1.1

Let us first introduce some quadratic forms and operators which we shall need in the argument. To begin with, we need the following result, which is a straightforward modification of Lemma 2.1 of [4], hence we skip the proof.

Lemma 2.1. *Let Φ be the map*

$$[0, L) \times (0, a) \ni (s, u) \mapsto (\Gamma_1(s) + u\Gamma_2'(s), \Gamma_2(s) - u\Gamma_1'(s)) \in \mathbb{R}^2.$$

Then there exists an $a_1 > 0$ such that the map Φ is injective for any $a \in (0, a_1]$.

Choose an a satisfying $0 < a < a_1$, to be specified later, and let Σ_a be the strip neighborhood of Γ of width a ,

$$\Sigma_a := \Phi([0, L) \times [0, a)).$$

Then $\Omega \setminus \Sigma_a = \Lambda_a$ is a compact simply connected domain with the boundary $\Gamma_a := \Phi([0, L) \times \{a\})$. We define

$$q_{a,\beta}^D[f] := \|\nabla f\|_{\Sigma_a}^2 - \beta \int_{\Gamma} |f(x)|^2 ds \quad \text{for } f \in \{f \in H^1(\Sigma_a) : f|_{\Gamma_a} = 0\},$$

$$q_{a,\beta}^N[f] := \|\nabla f\|_{\Sigma_a}^2 - \beta \int_{\Gamma} |f(x)|^2 ds \quad \text{for } f \in H^1(\Sigma_a),$$

and denote by $L_{a,\beta}^D$ and $L_{a,\beta}^N$ the self-adjoint operators associated with the forms $q_{a,\beta}^D$ and $q_{a,\beta}^N$, respectively. The first key component of the proof is to use the Dirichlet-Neumann bracketing – see [9, Sec. XIII.15, Prop. 4] – imposing additional boundary condition at Γ_a . This yields

$$(-\Delta_{\Lambda_a}^N) \oplus L_{a,\beta}^N \leq H_\beta \leq (-\Delta_{\Lambda_a}^D) \oplus L_{a,\beta}^D \quad (6)$$

in $L^2(\Omega) = L^2(\Lambda_a) \oplus L^2(\Sigma_a)$ where the inequality should be understood, of course, in the variational sense. Since the estimating operators have the direct-sum structure and the first terms in the inequalities (6) referring to the part of Ω separated from the boundary are positive, in order to estimate the negative eigenvalues of H_β it is sufficient to estimate those of $L_{a,\beta}^D$, $L_{a,\beta}^N$.

To this aim we use the second main trick, introducing a “straightening” transformation in the spirit of [3], to produce a pair of operators in $L^2((0, L) \times (0, a))$ that are unitarily equivalent to $L_{a,\beta}^D$ and $L_{a,\beta}^N$, respectively. Specifically, we introduce the following change of variables,

$$f(x_1, x_2) = \frac{1}{(1 - u\gamma(s))^{1/2}} \varphi(s, u);$$

then it is straightforward to check that for any function $f \in H^2(\Sigma_a)$ we have also $\varphi \in H^2((0, L) \times (0, a))$ and

$$|f_{x_1}|^2 + |f_{x_2}|^2 = \left[\frac{1}{(1 - u\gamma(s))^2} \left| \frac{\partial \varphi}{\partial s} \right|^2 + \left| \frac{\partial \varphi}{\partial u} \right|^2 + \tilde{V}(s, u) |\varphi|^2 \right. \\ \left. + \frac{u\gamma'(s)}{2(1 - u\gamma(s))^3} \left(\varphi \frac{\partial \overline{\varphi}}{\partial s} + \overline{\varphi} \frac{\partial \varphi}{\partial s} \right) + \frac{\gamma(s)}{2(1 - u\gamma(s))} \left(\varphi \frac{\partial \overline{\varphi}}{\partial u} + \overline{\varphi} \frac{\partial \varphi}{\partial u} \right) \right]$$

with

$$\tilde{V}(s, u) = \frac{\gamma^2(s)}{4(1 - u\gamma(s))^2} + \frac{u^2(\gamma'(s))^2}{4(1 - u\gamma(s))^4},$$

where we employ the usual shorthands, $f_{x_j} = \frac{\partial f}{\partial x_j}$, and furthermore

$$\begin{aligned} & \iint_{\Sigma_a} (|f_{x_1}|^2 + |f_{x_2}|^2) dx_1 dx_2 - \beta \int_{\Gamma} |f(x)|^2 ds \\ &= \iint_0^a \int_0^L \frac{1}{(1 - u\gamma(s))^2} \left| \frac{\partial \varphi}{\partial s} \right|^2 ds du + \iint_0^a \int_0^L \left| \frac{\partial \varphi}{\partial u} \right|^2 ds du + \iint_0^a \int_0^L V(s, u) |\varphi|^2 ds du \\ & \quad - \int_0^L \left(\frac{\gamma(s)}{2} + \beta \right) |\varphi(s, 0)|^2 ds + \int_0^L \frac{\gamma(s)}{2(1 - a\gamma(s))} |\varphi(s, a)|^2 ds, \end{aligned}$$

where

$$\begin{aligned} V(s, u) &= \tilde{V}(s, u) - \frac{\partial}{\partial s} \left(\frac{u\gamma'(s)}{2(1 - u\gamma(s))^3} \right) - \frac{\partial}{\partial u} \left(\frac{\gamma(s)}{2(1 - u\gamma(s))} \right) \\ &= -\frac{\gamma^2(s)}{4(1 - u\gamma(s))^2} - \frac{u\gamma''(s)}{2(1 - u\gamma(s))^3} - \frac{5}{4} \frac{u^2(\gamma'(s))^2}{(1 - u\gamma(s))^4}. \end{aligned}$$

Armed with these formulæ we can now introduce the two operators in $L^2((0, L) \times (0, a))$ unitarily equivalent to $L_{a,\beta}^D$ and $L_{a,\beta}^N$, respectively. On the domains

$$Q_a^D = \{ \varphi \in H^1((0, L) \times (0, a)) : \varphi(L, \cdot) = \varphi(0, \cdot) \text{ on } (0, a), \varphi(\cdot, a) = 0 \text{ on } (0, L) \}$$

and

$$Q_a^N = \{ \varphi \in H^1((0, L) \times (0, a)) : \varphi(L, \cdot) = \varphi(0, \cdot) \text{ on } (0, a) \},$$

we define the quadratic forms

$$\begin{aligned} b_{a,\beta}^D[\varphi] &= \iint_0^a \int_0^L \frac{1}{(1 - u\gamma(s))^2} \left| \frac{\partial \varphi}{\partial s} \right|^2 ds du + \iint_0^a \int_0^L \left| \frac{\partial \varphi}{\partial u} \right|^2 ds du \\ & \quad + \iint_0^a \int_0^L V(s, u) |\varphi|^2 ds du - \int_0^L \left(\frac{\gamma(s)}{2} + \beta \right) |\varphi(s, 0)|^2 ds \end{aligned} \quad (7)$$

and

$$\begin{aligned}
b_{a,\beta}^N[\varphi] &= \iint_0^a \frac{1}{(1-u\gamma(s))^2} \left| \frac{\partial \varphi}{\partial s} \right|^2 ds du + \iint_0^a \left| \frac{\partial \varphi}{\partial u} \right|^2 ds du \\
&+ \iint_0^a V(s,u) |\varphi|^2 ds du - \int_0^L \left(\frac{\gamma(s)}{2} + \beta \right) |\varphi(s,0)|^2 ds \\
&+ \int_0^L \frac{\gamma(s)}{2(1-a\gamma(s))} |\varphi(s,a)|^2 ds,
\end{aligned}$$

respectively. It is easy to check the following claim analogous to Lemma 2.2 of [4].

Lemma 2.2. *The operators $B_{a,\beta}^D$ and $B_{a,\beta}^N$ associated with above quadratic forms are unitarily equivalent to $L_{a,\beta}^D$ and $L_{a,\beta}^N$, respectively.*

In the next step we estimate $B_{a,\beta}^D$ and $B_{a,\beta}^N$ just introduced by operators with separated variables. We put¹

$$\begin{aligned}
\gamma_+ &= \max_{[0,L]} |\gamma(\cdot)|, \quad \gamma'_+ = \max_{[0,L]} |\gamma'(\cdot)|, \quad \gamma''_+ = \max_{[0,L]} |\gamma''(\cdot)|, \\
V_+(s) &= \frac{-\gamma^2(s)}{4(1+a\gamma_+)^2} + \frac{a\gamma''_+}{2(1-a\gamma_+)^3}, \\
V_-(s) &= \frac{-\gamma^2(s)}{4(1-a\gamma_+)^2} - \frac{a\gamma''_+}{2(1-a\gamma_+)^3} - \frac{5}{4} \frac{a^2(\gamma'_+)^2}{(1-a\gamma_+)^4}.
\end{aligned}$$

For an a satisfying $0 < a < \gamma_+/2$ and φ belonging to Q_a^D and Q_a^N , respectively, we define

$$\begin{aligned}
\tilde{b}_{a,\beta}^D[\varphi] &= (1-a\gamma_+)^{-2} \iint_0^a \left| \frac{\partial \varphi}{\partial s} \right|^2 ds du + \iint_0^a \left| \frac{\partial \varphi}{\partial u} \right|^2 ds du \\
&+ \iint_0^a V_+(s) |\varphi|^2 ds du - \left(\frac{\gamma^*}{2} + \beta \right) \int_0^L |\varphi(s,0)|^2 ds
\end{aligned}$$

and

$$\begin{aligned}
\tilde{b}_{a,\beta}^N[\varphi] &= (1+a\gamma_+)^{-2} \iint_0^a \left| \frac{\partial \varphi}{\partial s} \right|^2 ds du + \iint_0^a \left| \frac{\partial \varphi}{\partial u} \right|^2 ds du \\
&+ \iint_0^a V_-(s) |\varphi|^2 ds du - \left(\frac{\gamma^*}{2} + \beta \right) \int_0^L |\varphi(s,0)|^2 ds - \frac{\gamma_+}{2(1-a\gamma_+)} \int_0^L |\varphi(s,a)|^2 ds.
\end{aligned}$$

¹There is a typo in [4]; the second term in the definition of V_+ there has to be deleted.

Then we have

$$b_{a,\beta}^D[\varphi] \leq \tilde{b}_{a,\beta}^D[\varphi] \quad \text{for } f \in Q_a^D, \quad (8)$$

$$b_{a,\beta}^N[\varphi] \geq \tilde{b}_{a,\beta}^N[\varphi] \quad \text{for } f \in Q_a^N. \quad (9)$$

Let $\tilde{H}_{a,\beta}^D$ and $\tilde{H}_{a,\beta}^N$ be the self-adjoint operators associated with the forms $\tilde{b}_{a,\beta}^D$ and $\tilde{b}_{a,\beta}^N$, respectively. By $T_{a,\beta}^D$ we denote the self-adjoint operator associated with the form

$$t_{a,\beta}^D[\varphi] = \int_0^a |\varphi'(u)|^2 du - \left(\frac{\gamma^*}{2} + \beta \right) |\varphi(0)|^2$$

defined on $\{\varphi \in H^1(0, a) : \varphi(a) = 0\}$. Similarly, $T_{a,\beta}^N$ is the self-adjoint operator associated with the form

$$t_{a,\beta}^N(\varphi, \varphi) = \int_0^a |\varphi'(u)|^2 du - \left(\frac{\gamma^*}{2} + \beta \right) |\varphi(0)|^2, \quad \varphi \in H^1(0, a).$$

Furthermore, we introduce the operators

$$U_a^D = (1 - a\gamma_+)^{-2} \left(-\frac{d^2}{ds^2} \right) + V_+(s), \quad U_a^N = (1 + a\gamma_+)^{-2} \left(-\frac{d^2}{ds^2} \right) + V_-(s)$$

in $L^2(0, L)$, the domain of both of them being P given by (4). Then we have

$$\tilde{H}_{a,\beta}^D = U_a^D \otimes I + I \otimes T_{a,\beta}^D, \quad \tilde{H}_{a,\beta}^N = U_a^N \otimes I + I \otimes T_{a,\beta}^N, \quad (10)$$

and we can estimate contributions from the longitudinal and transverse variables separately. What concerns the former, we denote by $\mu_j^D(a)$, $\mu_j^N(a)$ the j -th eigenvalue of U_a^D , U_a^N , respectively, counted with the multiplicity, and use Proposition 2.3 of [4] which contains the following claim:

Lemma 2.3. *There exists a constant $C > 0$ such that the estimates*

$$|\mu_j^D(a) - \mu_j| \leq C a j^2 \quad (11)$$

and

$$|\mu_j^N(a) - \mu_j| \leq C a j^2 \quad (12)$$

hold for any $j \in \mathbb{N}$ and $0 < a < 1/(2\gamma_+)$. where C is independent on j, a .

We stress that the constant C here is independent of j and a . As for the transverse part, let us estimate first the principal eigenvalue of $T_{a,\beta}^D$.

Lemma 2.4. *Assume that $a(\beta + \frac{\gamma^*}{2}) > \frac{4}{3}$. Then $T_{a,\beta}^D$ has only one negative eigenvalue which we denote by $\zeta_{a,\beta}^D$. It satisfies the inequalities*

$$-\left(\beta + \frac{\gamma^*}{2}\right)^2 \leq \zeta_{a,\beta}^D \leq -\left(\beta + \frac{\gamma^*}{2}\right)^2 + 4\left(\beta + \frac{\gamma^*}{2}\right)^2 e^{-a(\beta + \frac{\gamma^*}{2})}.$$

Proof. Notice that the domain of the operator is

$$D(T_{a,\beta}^D) = \left\{ \varphi \in H^2(0, a) : \varphi'(0) = -\left(\frac{\gamma_*}{2} + \beta\right) \varphi(0), \varphi(a) = 0 \right\}.$$

Assume that $-k^2$ with $k > 0$ is an eigenvalue of $T_{a,\beta}^D$, and let a nonzero φ be the corresponding eigenfunction, then we have

- (1) $-\varphi''(u) = -k^2 \varphi(u)$;
- (2) $\varphi'(0) = -\left(\frac{\gamma_*}{2} + \beta\right) \varphi(0)$;
- (3) $\varphi(a) = 0$.

In view of the first property, the eigenfunction φ is of the form

$$\varphi(u) = Ae^{ku} + Be^{-ku}.$$

Furthermore, the requirements (2) and (3) yield $kA - kB = \left(-\frac{\gamma_*}{2} - \beta\right)(A + B)$ and $Ae^{ka} + Be^{-ka} = 0$, respectively. Thus the coefficients A, B have to satisfy the equation

$$\begin{pmatrix} e^{ka} & e^{-ka} \\ k + \frac{\gamma_*}{2} + \beta & -(k - \frac{\gamma_*}{2} - \beta) \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = 0.$$

Since $(A, B) \neq (0, 0)$, we get

$$\det \begin{pmatrix} e^{ka} & e^{-ka} \\ k + \frac{\gamma_*}{2} + \beta & -(k - \frac{\gamma_*}{2} - \beta) \end{pmatrix} = 0$$

which is equivalent to $g_{a,\beta}(k) := 2ak + \log\left(\beta + \frac{\gamma_*}{2} - k\right) - \log\left(\beta + \frac{\gamma_*}{2} + k\right) = 0$. It is easy to see that also the converse is true: if $g_{a,\beta}(k) = 0$, then $-k^2$ is an eigenvalue of $T_{a,\beta}^D$. Let us now show that $g_{a,\beta}(\cdot)$ has a unique zero in $(0, \beta + \frac{\gamma_*}{2})$. By definition we have $g_{a,\beta}(0) = 0$, and since

$$\frac{dg_{a,\beta}(k)}{dk} = \frac{2a\left(\beta + \frac{\gamma_*}{2}\right)^2 - 2\left(\beta + \frac{\gamma_*}{2}\right) - 2ak^2}{\left(\beta + \frac{\gamma_*}{2}\right)^2 - k^2}$$

we can claim that $g_{a,\beta}$ is monotonically increasing in $\left(0, \beta + \frac{\gamma_*}{2} - \frac{1}{a}\right)$ and it is monotonically decreasing in $\left(\beta + \frac{\gamma_*}{2} - \frac{1}{a}, \beta + \frac{\gamma_*}{2}\right)$. Moreover, we have

$$\lim_{k \rightarrow \beta + \frac{\gamma_*}{2}} g_{a,\beta} = -\infty;$$

this implies that the function $g_{a,\beta}$ has a unique zero in $\left(0, \beta + \frac{\gamma_*}{2}\right)$. Moreover, since $a\left(\beta + \frac{\gamma_*}{2}\right) > \frac{4}{3}$, we have $\sqrt{\left(\beta + \frac{\gamma_*}{2}\right)\left(\beta + \frac{\gamma_*}{2} - \frac{1}{a}\right)} > \frac{1}{2}\left(\beta + \frac{\gamma_*}{2}\right)$. Consequently,

the solution k has the form $k = \beta + \frac{\gamma^*}{2} - s$, $0 < s < \frac{1}{2}(\beta + \frac{\gamma^*}{2})$. Taking into account the relation $g_{a,\beta}(k) = 0$, we get

$$\log s = \log(2\beta + \gamma^* - s) - 2a \left(\beta + \frac{\gamma^*}{2} - s \right) \leq \log(2\beta + \gamma^*) - a \left(\beta + \frac{\gamma^*}{2} \right).$$

Hence we obtain $s \leq (2\beta + \gamma^*)e^{-a(\beta + \gamma^*/2)}$ which concludes the proof. \square

Next we estimate the first eigenvalue of $T_{a,\beta}^N$.

Lemma 2.5. *Assume that $\left(\beta + \frac{\gamma^*}{2}\right) > \max\left\{\frac{\gamma_+}{2(1-a\gamma_+)}, \frac{2\log 5}{3a}\right\}$. Then $T_{a,\beta}^N$ has a unique negative eigenvalue $\zeta_{a,\beta}^N$, and moreover, we have*

$$-\left(\beta + \frac{\gamma^*}{2}\right)^2 - \frac{45}{4}\left(\beta + \frac{\gamma^*}{2}\right)^2 e^{-a(\beta + \frac{\gamma^*}{2})} \leq \zeta_{a,\beta}^N \leq -\left(\beta + \frac{\gamma^*}{2}\right)^2.$$

Proof. The operator domain in this case looks as follows,

$$D(T_{a,\beta}^N) = \left\{ \varphi \in H^2(0, a) : \varphi'(0) = -\left(\frac{\gamma^*}{2} + \beta\right)\varphi(0), \varphi'(a) = \frac{\gamma_+}{2(1-a\gamma_+)}\varphi(a) \right\}.$$

Assume again that $-k^2$ with $k > 0$ is an eigenvalue of $T_{a,\beta}^N$ corresponding to a nonzero eigenfunction φ . As in the proof of Lemma 2.4 we infer that $-k^2$ is an eigenvalue of $T_{a,\beta}^N$ if and only if²

$$e^{2ka} = \frac{k + \frac{\gamma^*}{2} + \beta}{k - \frac{\gamma^*}{2} - \beta} \cdot \frac{k + \frac{\gamma_+}{2(1-a\gamma_+)}}{k - \frac{\gamma_+}{2(1-a\gamma_+)}}. \quad (13)$$

Since the left-hand side of the last equation is strictly increasing, and the right-hand side is strictly decreasing for $k > 0$, then the equation (13) has a unique positive solution which lies in fact in the subinterval $\left(\beta + \frac{\gamma^*}{2}, +\infty\right)$.

Next we will show that (13) has no solutions in the interval $k \geq \frac{3}{2}\left(\beta + \frac{\gamma^*}{2}\right)$. Suppose that the opposite is true. As $\frac{\gamma_+}{2(1-a\gamma_+)} < \beta + \frac{\gamma^*}{2}$, we have

$$\frac{k + \frac{\gamma^*}{2} + \beta}{k - \frac{\gamma^*}{2} - \beta} \cdot \frac{k + \frac{\gamma_+}{2(1-a\gamma_+)}}{k - \frac{\gamma_+}{2(1-a\gamma_+)}} \leq \left(\frac{k + \frac{\gamma^*}{2} + \beta}{k - \frac{\gamma^*}{2} - \beta}\right)^2.$$

However, since we assume $k \geq \frac{3}{2}\left(\beta + \frac{\gamma^*}{2}\right)$, this would imply

$$e^{2ka} \leq \left(\frac{\frac{3}{2}\left(\beta + \frac{\gamma^*}{2}\right) + \frac{\gamma^*}{2} + \beta}{\frac{3}{2}\left(\beta + \frac{\gamma^*}{2}\right) - \frac{\gamma^*}{2} - \beta}\right)^2 = 25.$$

²There is a misplaced exponential in the analogous proof in [4] which does not affect the claim.

On the other hand, we have $e^{2ka} \geq e^{3a\left(\beta + \frac{\gamma^*}{2}\right)} > 25$, so we come to a contradiction. Hence the solution k of (13) is of the form $k = \beta + \frac{\gamma^*}{2} + s$ with $0 < s < \frac{1}{2}\left(\beta + \frac{\gamma^*}{2}\right)$, and using (13) once again we get

$$e^{2ka} \leq \left(\frac{k + \frac{\gamma^*}{2} + \beta}{k - \frac{\gamma^*}{2} - \beta}\right)^2 \leq \left(\frac{2\beta + \gamma^* + s}{s}\right)^2 \leq \left(\frac{\frac{5}{2}\left(\beta + \frac{\gamma^*}{2}\right)}{s}\right)^2,$$

which further implies

$$s \leq \frac{5}{2}\left(\beta + \frac{\gamma^*}{2}\right) e^{-ka} = \frac{5}{2}\left(\beta + \frac{\gamma^*}{2}\right) e^{-a\left(\beta + \frac{\gamma^*}{2}\right) - sa} \leq \frac{5}{2}\left(\beta + \frac{\gamma^*}{2}\right) e^{-a\left(\beta + \frac{\gamma^*}{2}\right)}.$$

This completes the proof of Lemma 2.5. \square

Now we are finally in position to *prove Theorem 1.1*. We use first the bracketing to squeeze the eigenvalues in question between those of the operators (10). Since the latter have separated variables, their eigenvalues are sums of eigenvalues of the longitudinal and transverse component which we have estimated in Lemmata 2.3 and 2.4, 2.5, respectively, and it is sufficient to choose $a = \frac{6}{\beta} \log \beta$ to get (5). \square

Note that while the argument is pretty much the same as in the proof of Theorem 1 of [4], with Propositions 2.3–2.5 there replaced by the above mentioned lemmata, the result is much weaker due to the presence of the last term in the form $b_{a,\beta}^D$ and its counterpart in $b_{a,\beta}^N$. In particular, the estimates of Theorem 1.1 squeeze to produce an exact asymptotic expansion if and only if the curvature is constant, $\gamma^* = \gamma_*$. Let us now look at this case in more detail:

Example 2.6. Let Ω be a disc of radius R centered at the origin. In this case we have

$$\gamma(s) \equiv \gamma = \frac{1}{R} \tag{14}$$

and the eigenvalues μ_j of the comparison operator S given by (3) can be computed explicitly,

$$\mu_j = \left(-\frac{1}{4} + \left[\frac{j}{2}\right]^2\right) R^{-2}, \tag{15}$$

where $[y]$ denotes the maximum integer which less or equal to y . We introduce the usual polar coordinates,

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad 0 \leq r \leq R, \quad 0 \leq \theta < 2\pi,$$

writing with an abuse of notation $f(x, y) \equiv f(r, \theta)$. Equations (1) with $\lambda = -k^2$ now read

$$\begin{cases} \frac{\partial^2 f}{\partial r^2} + \frac{1}{r} \frac{\partial f}{\partial r} + \frac{1}{r^2} \frac{\partial^2 f}{\partial \theta^2} = k^2 f, \\ \frac{\partial f}{\partial r} \Big|_{r=R} = \beta f. \end{cases} \quad (16)$$

Solution to the first equation in (16) is conventionally sought in the form

$$f(r, \theta) = \sum_{m \in \mathbb{Z}} c_m I_m(kr) e^{im\theta}. \quad (17)$$

Furthermore, the Hamiltonian commutes with the angular momentum operator, $-i \frac{\partial}{\partial \theta}$ with periodic boundary conditions, hence the two operators have common eigenspaces, and we can consider sequence $\{c_m\}$ with nonzero c_m corresponding to a single values of $|m|$; it goes without saying that the discrete spectrum has multiplicity two except the eigenvalue corresponding to $m = 0$ which is simple. The boundary condition in (16) can be then rewritten as

$$kI'_m(kR) - \beta I_m(kR) = 0. \quad (18)$$

for a fixed $m \in \mathbb{Z}$. To find its solutions, let us change the variables to $X = kR$, $\alpha = \beta R$, in which case the condition (18) reads

$$\frac{XI'_m(X)}{I_m(X)} = \alpha. \quad (19)$$

The function at the left-hand side of (19) is strictly increasing for $k > 0$, hence (19) has a unique solution for any fixed α and m . As $\alpha \rightarrow +\infty$, so does X in (19), and using the well-known asymptotics of modified Bessel functions, we find

$$\frac{XI'_m(X)}{I_m(X)} = X - \frac{1}{2} + \frac{4m^2 - 1}{8X} + O(X^{-2}), \quad X \rightarrow +\infty.$$

In combination with the spectral condition (19) this yields

$$X = \alpha + \frac{1}{2} - \frac{4m^2 - 1}{8\alpha} + O(\alpha^{-2}), \quad \alpha \rightarrow +\infty.$$

This, in turn, implies the asymptotics for X^2 , and returning to the original variables β, k we find

$$-k^2 = -\left(\beta + \frac{1}{2R}\right)^2 + \left(m^2 - \frac{1}{4}\right)R^{-2} + \mathcal{O}(\beta^{-1}), \quad \beta \rightarrow +\infty.$$

This agrees, of course, with the conclusion of Theorem 1.1 according to (14) and (15). At the same time it shows that there is not much room for improving the error term in the theorem, because it differs from the one in this explicitly solvable example by the logarithmic factor only.

3. Proof of Theorem 1.3

The idea is to replace the crude estimate of $B_{a,\beta}^D$ from Lemma 2.2 by the first operator in (10) by a finer one. Consider first the principal eigenvalue which satisfies $\lambda_1(\beta) \leq b_{a,\beta}^D[\varphi]$ for any $\varphi \in Q_a^D$ and choose the following family of trial functions,

$$\hat{\varphi}(s, u) = \chi_\varepsilon(s) (e^{-\alpha u} - e^{-2a\alpha + u\alpha}),$$

where χ_ε is a smooth function on $[0, L]$ with the support located in an ε -neighborhood of a point s^* in which the curvature reaches its maximum, $\gamma(s^*) = \gamma^*$, and ε is a parameter to be determined later. In view of the boundary compactness and smoothness, at least one such point exists; without loss of generality we may assume that $(s^* - \varepsilon, s^* + \varepsilon) \subset (0, L)$. We shall consider functions of the form

$$\chi_\varepsilon(s) := \chi\left(\frac{s - s^* + \varepsilon}{2\varepsilon}\right),$$

where $\chi(x)$ is a fixed smooth function on \mathbb{R} with the support in the interval $(0, 1)$; then we have

$$\|\chi_\varepsilon\|_{L^2(0,L)}^2 = 2\varepsilon\|\chi\|_{L^2(0,1)}^2, \quad \|\chi'_\varepsilon\|_{L^2(0,L)}^2 = (2\varepsilon)^{-1}\|\chi'\|_{L^2(0,1)}^2. \quad (20)$$

We also note that on the support of χ_ε , i.e. for any $s \in (s^* - \varepsilon, s^* + \varepsilon)$ we have

$$|\gamma(s) - \gamma^*| < \gamma'_+ |s - s^*| < \gamma'_+ \varepsilon.$$

Computing the terms of the form $b_{a,\beta}^D[\varphi]$ we get for the longitudinal kinetic contribution the estimate

$$\begin{aligned} \iint_0^a \int_0^L \frac{1}{(1 - u\gamma(s))^2} \left| \frac{\partial \hat{\varphi}}{\partial s} \right|^2 ds du &\leq \int_0^a \int_{s^* - \varepsilon}^{s^* + \varepsilon} \left(\frac{1}{(1 - u\gamma^*)^2} + C\varepsilon u \right) (e^{-\alpha u} - e^{-2a\alpha + u\alpha})^2 \\ &\quad \times (\chi'(s))^2 ds du \left[\left(\frac{1}{2\alpha} + \mathcal{O}(\alpha^{-2}) \right) + C\varepsilon \left(\frac{1}{4\alpha^2} + \mathcal{O}(\alpha^{-3}) \right) \right] \|\chi\|_{L^2(0,L)}^2, \end{aligned}$$

where $C > 0$ is a generic constant independent of β, a , and ε . Similarly,

$$\iint_0^a \int_0^L \left| \frac{\partial \hat{\varphi}}{\partial u} \right|^2 ds du = \frac{\alpha}{2} (1 + \mathcal{O}(\alpha e^{-2a\alpha})) \|\chi\|_{L^2(0,L)}^2$$

holds for the transverse kinetic term,

$$\begin{aligned} \int_0^a \int_0^L V(s, u) |\hat{\varphi}|^2 ds du &\leq \int_0^a \int_{s^* - \varepsilon}^{s^* + \varepsilon} \left(-\frac{(\gamma^*)^2}{4(1 - u\gamma^*)} - \frac{u\gamma''(s^*)}{2(1 - u\gamma^*)^3} - \frac{5u^2(\gamma'(s^*))^2}{4(1 - u\gamma^*)^4} + C\varepsilon \right) \\ &\quad \times (e^{-\alpha u} - e^{-2a\alpha + u\alpha})^2 |\chi(s)|^2 ds du = \left(\frac{-(\gamma^*)^2}{4} + C\varepsilon \right) \frac{1}{2\alpha} (1 + \mathcal{O}(\alpha^{-1})) \|\chi\|_{L^2(0,L)}^2 \end{aligned}$$

for the potential one, and

$$-\int_0^L \left(\frac{\gamma(s)}{2} + \beta \right) |\hat{\varphi}(s, 0)|^2 ds \geq - \left(\beta + \frac{\gamma^* - \varepsilon}{2} \right) (1 - e^{-2a\alpha})^2 \|\chi\|_{L^2(0,L)}^2,$$

for the boundary one. Finally, the trial function norm satisfies

$$\iint_0^a \int_0^L |\hat{\varphi}(s, u)|^2 ds du = \frac{1}{2\alpha} (1 + \mathcal{O}(\alpha e^{-2a\alpha})) \|\chi\|_{L^2(0,L)}^2.$$

Putting these expressions together and taking (20) into account we get

$$\begin{aligned} \frac{b_{a,\beta}^D[\hat{\varphi}]}{\|\hat{\varphi}\|_{L^2(0,L)}^2} &\leq \frac{1}{4\varepsilon^2} \frac{\|\chi'\|_{L^2(0,1)}^2}{\|\chi\|_{L^2(0,1)}^2} \left(1 + \mathcal{O}(\alpha^{-1}) + C\varepsilon \left(\frac{1}{2\alpha} + \alpha^{-2} \right) \right) \\ &\quad + \alpha^2 (1 + \mathcal{O}(\alpha e^{-2a\alpha})) + \left(\frac{-(\gamma^*)^2}{4} + C\varepsilon \right) (1 + \mathcal{O}(\alpha^{-1})) \\ &\quad - 2\alpha \left(\beta + \frac{\gamma^* - \varepsilon}{2} \right) (1 + \mathcal{O}(\alpha e^{-2a\alpha})). \end{aligned}$$

Now we choose $\alpha = \beta + \frac{\gamma^*}{2}$ in which case the right-hand side of the last inequality becomes

$$\begin{aligned} \frac{1}{4\varepsilon^2} \frac{\|\chi'\|_{L^2(0,1)}^2}{\|\chi\|_{L^2(0,1)}^2} \left(1 + \mathcal{O}(\beta^{-1}) + C\varepsilon \left(\frac{1}{2\beta} + \beta^{-2} \right) \right) - \left(\beta + \frac{\gamma^*}{2} \right)^2 \\ + \varepsilon \left(\beta + \frac{\gamma^*}{2} \right) + \left(\frac{-(\gamma^*)^2}{4} + C\varepsilon \right) (1 + \mathcal{O}(\alpha^{-1})), \end{aligned}$$

and to optimize the last formula with respect to ε we take $\varepsilon = \beta^{-1/3}$, which yields the estimate

$$\frac{b_{a,\beta}^D[\hat{\varphi}]}{\|\hat{\varphi}\|_{L^2(0,L)}^2} \leq - \left(\beta + \frac{\gamma^*}{2} \right)^2 + \mathcal{O}(\beta^{2/3}) \quad (21)$$

proving the result. The argument for the higher eigenfunctions proceeds in the same way. We employ trial functions of the form

$$\hat{\varphi}_j(s, u) = \chi_{\varepsilon,j}(s) (e^{-\alpha u} - e^{-2a\alpha + u\alpha}),$$

where the longitudinal part is constructed from a shifted function χ , for instance

$$\chi_{\varepsilon,j}(s) := \chi \left(\frac{s - s^* + (2j - 1)\varepsilon}{2\varepsilon} \right).$$

The above estimate of the form remains essentially the same, up to the values of the constants involved. By construction, the functions $\chi_{\varepsilon,j}$ with different values of j have disjoint supports, hence $\hat{\varphi}_j$ is orthogonal to $\hat{\varphi}_i$, $i = 1, \dots, j-1$, and by the min-max principle [9, Sec. XIII.1] the eigenvalue $\lambda_j(\beta)$ has again the upper bound given by the right-hand side of (21). \square

References

- [1] D. Daners and J. Kennedy: On the asymptotic behaviour of the eigenvalues of a Robin problem, *Diff. Int. Eq.* **23** (2010), 659–669.
- [2] P. Exner, Leaky quantum graphs: a review, in “Analysis on graphs and its applications”, Proc. Symp. Pure Math., vol. 77; Amer. Math. Soc., Providence, R.I., 2008., pp. 523–564.
- [3] P. Exner and P. Šeba, Bound states in curved quantum waveguides. *J. Math. Phys.* **30**, no. 11 (1989), 2574–2580.
- [4] P. Exner and K. Yoshitomi, Asymptotics of eigenvalues of the Schrödinger operator with a strong δ -interaction on a loop. *J. Geom. Phys.* **41**, no. 4 (2002), 344–358.
- [5] A.A. Lacey, J.R. Ockendon, J. Sabina, and D. Salazar, Perturbation analysis of a semilinear parabolic problem with nonlinear boundary conditions. *Rocky Mountain J. Math.* **26**, no.1 (1996), 195–212.
- [6] A.A. Lacey, J.R. Ockendon, and J. Sabina, Multidimensional reaction diffusion equations with nonlinear boundary conditions. *SIAM J. Appl. Math.* **58**, no.5 (1998), 1622–1647.
- [7] M. Levitin and L. Parnovski, On the principal eigenvalue of a Robin problem with a large parameter. *Math. Nachr.* **281**, no. 2 (2008), 272–281.
- [8] K. Pankrashkin, On the asymptotics of the principal eigenvalue for a Robin problem with a large parameter in planar domains. *Nanosystems: Phys. Chem. Math.* **4**, no. 4 (2013), 474–483.
- [9] M. Reed and B. Simon, *Methods of Modern Mathematical Physics. IV. Analysis of Operators*, Academic Press, San Diego, 1978.

Received

Pavel Exner^{1,3}

E-mail: exner@ujf.cas.cz

Alexander Minakov^{1,2}

E-mail: minakov.ilt@gmail.com

Leonid Parnovski⁴

E-mail: l.parnovski@ucl.ac.uk

⁽¹⁾Doppler Institute for Mathematical Physics and Applied Mathematics, Czech Technical University in Prague, Břehová 7, 11519 Prague, Czech Republic

⁽²⁾Department of Physics, Faculty of Nuclear Science and Physical Engineering,
Czech Technical University in Prague, Pohraniční 1288/1, 40501 Děčín, Czech
Republic

⁽³⁾Department of Theoretical Physics, Nuclear Physics Institute ASCR, 25068 Řež
near Prague, Czech Republic

⁽⁴⁾Department of Mathematics, University College London
Gower Street, London WC1E 6BT, United Kingdom