

# SPECTRAL THEORY FOR THE WEAK DECAY OF MUONS IN A UNIFORM MAGNETIC FIELD

JEAN-CLAUDE GUILLOT

*In memory of Raymond Stora*

ABSTRACT. In this article we consider a mathematical model for the weak decay of muons in a uniform magnetic field according to the Fermi theory of weak interactions with V-A coupling. With this model we associate a Hamiltonian with cutoffs in an appropriate Fock space. No infrared regularization is assumed. The Hamiltonian is self-adjoint and has a unique ground state. We specify the essential spectrum and prove the existence of asymptotic fields from which we determine the absolutely continuous spectrum. The coupling constant is supposed sufficiently small.

## CONTENTS

1. Introduction.	1
2. The Hamiltonian.	2
2.1. The free Hamiltonian.	3
2.2. The Interaction.	5
3. Existence of a unique ground state.	6
4. The absolutely continuous spectrum.	7
4.1. Asymptotic fields.	7
4.2. Existence of a Fock space subrepresentation of the asymptotic CAR	19
Acknowledgements.	22
Appendix A. The Dirac quantized fields for the electrons and the muons.	22
Appendix B. The Dirac quantized fields for $\nu_\mu$ and $\bar{\nu}_e$ .	25
References	27

## 1. INTRODUCTION.

In this paper we consider a mathematical model for the weak decay of muons into electrons, neutrinos and antineutrinos in a uniform magnetic field according to the Fermi theory with V-A (Vector-Axial Vector)coupling,

$$(1.1) \quad \mu_- \rightarrow e_- + \bar{\nu}_e + \nu_\mu$$

$$(1.2) \quad \mu_+ \rightarrow e_+ + \nu_e + \bar{\nu}_\mu$$

(1.2) is the charge conjugation of (1.1).

This is a part of a program devoted to the study of mathematical models for the weak interactions as patterned according to the Fermi theory and the Standard model in Quantum Field Theory. See [21].

---

2010 *Mathematics Subject Classification.* 81V15, 81V10, 81Q10.

*Key words and phrases.* Weak Decay of muons, Fermi's theory, Uniform Magnetic Field, Spectral Theory,

In this paper we restrict ourselves to the study of the decay of the muon  $\mu_-$  whose electric charge is the charge of the electron (1.1). The study of the decay of the antiparticle  $\mu_+$ , whose charge is positive, (1.2) is quite similar and we omit it.

In [23] we have studied the spectral theory of the Hamiltonian associated to the inverse  $\beta$  decay in a uniform magnetic field. We proved the existence and uniqueness of a ground state and we specify the essential spectrum and the spectrum for a small coupling constant and without any low-energy regularization.

In this paper we consider the weak decay of muons into electrons, neutrinos associated with muons and antineutrinos associated with electrons in a uniform magnetic field according to the Fermi theory with V-A coupling. Hence we neglect the small mass of neutrinos and antineutrinos and we define a total Hamiltonian  $H$  acting in an appropriate Fock space involving three fermionic massive particles-the electrons, the muons and the antimuons - and two fermionic massless particles-the neutrinos and the antineutrinos associated to the muons and the electrons respectively. In order to obtain a well-defined operator, we approximate the physical kernels of the interaction Hamiltonian by square integrable functions and we introduce high-energy cutoffs. We do not need to impose any low-energy regularization in this work but the coupling constant is supposed sufficiently small.

We give a precise definition of the Hamiltonian as a self-adjoint operator in the appropriate Fock space and by adapting the methods used in [23] we first state that  $H$  has a unique ground state and we specify the essential spectrum for sufficiently small values of the coupling constant.

In this paper, our main result is the location of the absolutely continuous spectrum of  $H$ . For that we follow the first step of the approach to scattering theory in establishing, for each involved particle, the existence and basic properties of the asymptotic creation and annihilation operators for time  $t$  going to  $\pm\infty$ . We then have a natural definition of unitary wave operators with the right intertwining property from which we deduce the absolutely continuous spectrum of  $H$ . Scattering theory for models in Quantum Field Theory without any external field has been considered by many authors. See, among others, [1, 3, 16, 17, 19, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37] and references therein. A part of the techniques used in this paper are adapted from the ones developed in these references. Note that the asymptotic completeness of the wave operators is an open problem in the case of the weak interactions in the background of a uniform magnetic field. See (1.1) for a study of scattering theory for a mathematical model of the weak interactions without any external field.

In some parts of our presentation we will only give the statement of theorems referring otherwise to some references.

The paper is organized as follows. In the second section we define the regularized self-adjoint Hamiltonian associated to (1.1). In the third section we consider the existence of a unique ground state and we specify the essential spectrum of  $H$ . In the fourth section we carefully prove the existence of asymptotic limits, when time  $t$  goes to  $\pm\infty$ , of the creation and annihilation operators of each involved particle, we define a unitary wave operator and we prove that it satisfies the right intertwining property with the hamiltonian and we deduce the absolutely continuous spectrum of  $H$ . In Appendices A and B we recall the Dirac quantized fields associated to the muon and the electron in a uniform magnetic external field together with the Dirac quantized free fields associated to the neutrino and the antineutrino.

## 2. THE HAMILTONIAN.

In the Fermi theory the decay of the muon  $\mu$  is described by the following four fermions effective Hamiltonian for the interaction in the Schrödinger representation ( see [21], [24] and [41]):

$$(2.1) \quad \begin{aligned} H_{int} = & \\ & \frac{G_F}{\sqrt{2}} \int d^3x (\bar{\Psi}_{\nu_\mu}(x) \gamma^\alpha (1 - \gamma_5) \Psi_\mu(x)) (\bar{\Psi}_e(x) \gamma_\alpha (1 - \gamma_5) \Psi_{\nu_e}(x)) \\ & + \frac{G_F}{\sqrt{2}} \int d^3x (\bar{\Psi}_{\nu_e}(x) \gamma_\alpha (1 - \gamma_5) \Psi_e(x)) (\bar{\Psi}_\mu(x) \gamma^\alpha (1 - \gamma_5) \Psi_{\nu_\mu}(x)) \end{aligned}$$

Here  $\gamma^\alpha$ ,  $\alpha = 0, 1, 2, 3$  and  $\gamma_5$  are the Dirac matrices in the standard representation.  $\Psi_{(\cdot)}(x)$  and  $\bar{\Psi}_{(\cdot)}(x)$  are the quantized Dirac fields for  $e$ ,  $\mu$ ,  $\nu_\mu$  and  $\nu_e$ .  $\bar{\Psi}_{(\cdot)}(x) = \Psi_{(\cdot)}(x)^\dagger \gamma^0$ .  $G_F$  is the Fermi coupling constant with  $G_F \simeq 1.16639(2) \times 10^{-5} GeV^{-2}$ . See [12].

We recall that  $m_e < m_\mu$ .  $\nu_\mu$  and  $\nu_e$  are massless particles.

### 2.1. The free Hamiltonian.

Throughout this work notations are introduced in appendices A and B.

Let

$$(2.2) \quad \mathfrak{F} = \mathfrak{F}_e \otimes \mathfrak{F}_\mu \otimes \mathfrak{F}_\mu \otimes \mathfrak{F}_{\nu_\mu} \otimes \mathfrak{F}_{\bar{\nu}_e}$$

Let

$$(2.3) \quad \begin{aligned} \omega(\xi_1) &= E_n^{(e)}(p^3) \quad \text{for } \xi_1 = (s, n, p^1, p^3) \\ \omega(\xi_2) &= E_n^{(\mu)}(p^3) \quad \text{for } \xi_2 = (s, n, p^1, p^3) \\ \omega(\xi_3) &= |\mathbf{p}| \quad \text{for } \xi_3 = (\mathbf{p}, \frac{1}{2}) \\ \omega(\xi_4) &= |\mathbf{p}| \quad \text{for } \xi_4 = (\mathbf{p}, -\frac{1}{2}). \end{aligned}$$

Let  $H_D^{(e)}$  ( resp.  $H_D^{(\mu-)}, H_D^{(\mu+)}$ , and  $H_D^{(\nu)}$  ) be the Dirac Hamiltonian for the electron ( resp. the muon, the antimuon and the neutrino ).

The quantization of  $H_D^{(e)}$ , denoted by  $H_{0,D}^{(e)}$  and acting on  $\mathfrak{F}_e$ , is given by

$$(2.4) \quad H_{0,D}^{(e)} = \int \omega(\xi_1) b_+^*(\xi_1) b_+(\xi_1) d\xi_1$$

Likewise the quantization of  $H_D^{(\mu-)}, H_D^{(\mu+)}, H_D^{(\bar{\nu}_e)}$  and  $H_D^{(\nu_\mu)}$ , denoted by  $H_{0,D}^{(\mu-)}, H_{0,D}^{(\mu+)}, H_{0,D}^{(\bar{\nu}_e)}$  and  $H_{0,D}^{(\nu_\mu)}$  respectively, acting on  $\mathfrak{F}_\mu, \mathfrak{F}_{\bar{\nu}_e}$  and  $\mathfrak{F}_{\nu_\mu}$  respectively, is given by

$$(2.5) \quad \begin{aligned} H_{0,D}^{(\mu-)} &= \int \omega(\xi_2) b_+^*(\xi_2) b_+(\xi_2) d\xi_2 \\ H_{0,D}^{(\mu+)} &= \int \omega(\xi_2) b_-^*(\xi_2) b_-(\xi_2) d\xi_2 \\ H_{0,D}^{(\bar{\nu}_e)} &= \int \omega(\xi_3) b_-^*(\xi_3) b_-(\xi_3) d\xi_3 \\ H_{0,D}^{(\nu_\mu)} &= \int \omega(\xi_4) b_+^*(\xi_4) b_+(\xi_4) d\xi_4. \end{aligned}$$

We set  $H_{0,D}^{(\mu)} = H_{0,D}^{(\mu-)} \otimes \mathbf{1} + \mathbf{1} \otimes H_{0,D}^{(\mu+)}$ .  $H_{0,D}^{(\mu)}$  is defined on  $\mathfrak{F}_\mu \otimes \mathfrak{F}_\mu$ .

For each Fock space  $\mathfrak{F}$ , let  $\mathfrak{D}^{(\cdot)}$  denote the set of vectors  $\Phi \in \mathfrak{F}^{(\cdot)}$  for which each component  $\Phi^{(r)}$  is smooth and has a compact support and  $\Phi^{(r)} = 0$  for all but finitely many  $r$ . Then  $H_{0,D}^{(\cdot)}$  is well-defined on the dense subset  $\mathfrak{D}^{(\cdot)}$  and it is

essentially self-adjoint on  $\mathfrak{D}^{(\cdot)}$ . The self-adjoint extension will be denoted by the same symbol  $H_{0,D}^{(\cdot)}$  with domain  $D(H_{0,D}^{(\cdot)})$ .

The spectrum of  $H_{0,D}^{(e)}$  in  $\mathfrak{F}^{(e)}$  is given by

$$(2.6) \quad \text{spec}(H_{0,D}^{(e)}) = \{0\} \cup [m_e, \infty)$$

$\{0\}$  is a simple eigenvalue whose the associated eigenvector is the vacuum in  $\mathfrak{F}^{(e)}$  denoted by  $\Omega^{(e)}$ .  $[m_e, \infty)$  is the absolutely continuous spectrum of  $H_{0,D}^{(e)}$ .

Likewise the spectra of  $H_{0,D}^{(\mu)}$ ,  $H_{0,D}^{(\bar{\nu}_e)}$  and  $H_{0,D}^{(\nu_\mu)}$  in  $\mathfrak{F}^{(\mu)} \otimes \mathfrak{F}^{(\mu)}$ ,  $\mathfrak{F}^{(\bar{\nu}_e)}$  and  $\mathfrak{F}^{(\nu_\mu)}$  respectively are given by

$$(2.7) \quad \begin{aligned} \text{spec}(H_{0,D}^{(\mu)}) &= \{0\} \cup [m_\mu, \infty) \\ \text{spec}(H_{0,D}^{(\bar{\nu}_e)}) &= [0, \infty) \\ \text{spec}(H_{0,D}^{(\nu_\mu)}) &= [0, \infty). \end{aligned}$$

$\Omega^{(\mu)}$ ,  $\Omega^{(\bar{\nu}_e)}$  and  $\Omega^{(\nu_\mu)}$  are the associated vacua in  $\mathfrak{F}^{(\mu)} \otimes \mathfrak{F}^{(\mu)}$ ,  $\mathfrak{F}^{(\bar{\nu}_e)}$  and  $\mathfrak{F}^{(\nu_\mu)}$  respectively and are the associated eigenvectors of  $H_{0,D}^{(\mu)}$ ,  $H_{0,D}^{(\bar{\nu}_e)}$  and  $H_{0,D}^{(\nu_\mu)}$  respectively for the eigenvalue  $\{0\}$ .

The vacuum in  $\mathfrak{F}$ , denoted by  $\Omega$ , is then given by

$$(2.8) \quad \Omega = \Omega^{(e)} \otimes \Omega^{(\mu)} \otimes \Omega^{(\bar{\nu}_e)} \otimes \Omega^{(\nu_\mu)}$$

The free Hamiltonian for the model, denoted by  $H_0$  and acting in  $\mathfrak{F}$ , is now given by

$$(2.9) \quad \begin{aligned} H_0 &= H_{0,D}^{(e)} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes H_{0,D}^{(\mu)} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \\ &+ \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes H_{0,D}^{(\bar{\nu}_e)} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes H_{0,D}^{(\nu_\mu)}. \end{aligned}$$

$H_0$  is essentially self-adjoint on  $\mathfrak{D} = \mathfrak{D}^{(e)} \widehat{\otimes} \mathfrak{D}^{(\mu)} \widehat{\otimes} \mathfrak{D}^{(\bar{\nu}_e)} \widehat{\otimes} \mathfrak{D}^{(\nu_\mu)}$ .

Here  $\widehat{\otimes}$  is the algebraic tensor product.

$\text{spec}(H_0) = [0, \infty)$  and  $\Omega$  is the eigenvector associated with the simple eigenvalue  $\{0\}$  of  $H_0$ .

Let  $S^{(e)}$  be the set of the thresholds of  $H_{0,D}^{(e)}$ :

$$S^{(e)} = (s_n^{(e)}; n \in \mathbb{N})$$

with  $s_n^{(e)} = \sqrt{m_e^2 + 2neB}$ .

Likewise let  $S^{(\mu)}$  be the set of the thresholds of  $H_{0,D}^{(\mu)}$ :

$$S^{(\mu)} = (s_n^{(\mu)}; n \in \mathbb{N})$$

with  $s_n^{(\mu)} = \sqrt{m_\mu^2 + 2neB}$ .

Then

$$(2.10) \quad \mathfrak{S} = S^{(e)} \cup S^{(\mu)}$$

is the set of the thresholds of  $H_0$ .

Throughout this work any finite tensor product of annihilation or creation operators associated with the involved particles will be denoted for shortness by the usual product of the operators (see e.g (2.13) and (2.14)).

## 2.2. The Interaction.

Similarly to [8] [5], [9], [10] [11],[22]and [23], in order to get well-defined operators on  $\mathfrak{F}$ , we have to substitute smoother kernels  $F(\xi_2, \xi_4)$  and  $G(\xi_1, \xi_3)$  for the  $\delta$ -distribution associated with (2.1) (conservation of momenta) and for introducing ultraviolet cutoffs.

Let

$$(2.11) \quad \mathbf{r} = \mathbf{p}_3 + \mathbf{p}_4$$

We get a new operator denoted by  $H_I$  and defined as follows

$$(2.12) \quad H_I = H_I^1 + (H_I^1)^* + H_I^2 + (H_I^2)^*$$

Here

$$(2.13) \quad \begin{aligned} H_I^{(1)} &= \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ &\left. (\overline{U}^{(\nu\mu)}(\xi_4) \gamma^\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) U^{(\nu)}(\xi_4)) \right) \\ &F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_4) b_+^*(\xi_1) b_-^*(\xi_3) b_+(\xi_2). \end{aligned}$$

and

$$(2.14) \quad \begin{aligned} H_I^{(2)} &= \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ &\left. (\overline{U}^{(\nu\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) W^{(\overline{\nu e})}(\xi_3)) \right) \\ &F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_4) b_-^*(\xi_2) b_+^*(\xi_1) b_-^*(\xi_3). \end{aligned}$$

$H_I^{(1)}$  describes the decay of the muon and  $H_I^{(2)}$  is responsible for the fact that the bare vacuum will not be an eigenvector of the total Hamiltonian as expected from physics.

We now introduce the following assumptions on the kernels  $F(\xi_2, \xi_4)$  and  $G(\xi_1, \xi_3)$  in order to get well-defined Hamiltonians in  $\mathfrak{F}$ .

### Hypothesis 2.1.

$$(2.15) \quad \begin{aligned} F(\xi_2, \xi_4) &\in L^2(\Gamma_1 \times \mathbb{R}^3) \\ G(\xi_1, \xi_3) &\in L^2(\Gamma_1 \times \mathbb{R}^3) \end{aligned}$$

These assumptions will be needed throughout the paper.

By (2.12)-(2.15)  $H_I$  is well defined as a sesquilinear form on  $\mathfrak{D}$  and one can construct a closed operator associated with this form.

The total Hamiltonian is thus

$$(2.16) \quad H = H_0 + gH_I, \quad g > 0.$$

$g$  is the coupling constant that we suppose non-negative for simplicity. The conclusions below are not affected if  $g \in \mathbb{R}$ .

The self-adjointness of  $H$  is established by the next theorem.

Let

$$(2.17) \quad \begin{aligned} C &= \|\gamma^0 \gamma_\alpha (1 - \gamma_5)\|_{\mathbb{C}^4} \|\gamma^0 \gamma^\alpha (1 - \gamma_5)\|_{\mathbb{C}^4}. \\ \frac{1}{M} &= \frac{1}{m_e} + \frac{1}{m_\mu} \end{aligned}$$

For  $\phi \in D(H_0)$  we have

$$(2.18) \quad \begin{aligned} & \|H_I \phi\| \\ & \leq 2C \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \left( \frac{2}{M} \|H_0 \phi\| + \|\phi\| \right). \end{aligned}$$

(2.18) follows from standard estimates of creation and annihilation operators in Fock space (the  $N_\tau$  estimates, see [20]). Details can be found in [7, proposition 3.7].

**Theorem 2.2.** (*Self-adjointness*). *Let  $g_0 > 0$  be such that*

$$(2.19) \quad 4g_0 \frac{C}{M} \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} < 1.$$

*Then for any  $g$  such that  $g \leq g_0$   $H$  is self-adjoint in  $\mathfrak{F}$  with domain  $\mathcal{D}(H) = \mathcal{D}(H_0)$ . Moreover any core for  $H_0$  is a core for  $H$ .*

By (2.18) and (2.19) the proof of the self-adjointness of  $H$  follows from the Kato-Rellich theorem.

$\sigma(H)$  stands for the spectrum and  $\sigma_{\text{ess}}(H)$  denotes the essential spectrum. We have

**Theorem 2.3.** (*The essential spectrum and the spectrum*) *Setting*

$$E = \inf \sigma(H)$$

*we have for every  $g \leq g_0$*

$$\sigma(H) = \sigma_{\text{ess}}(H) = [E, \infty)$$

*with  $E \leq 0$ .*

In order to prove the theorem 2.3 we easily adapt to our case the proof given in [11] (see also [4], [38] and [23]). The mathematical model considered in [11] involves also one neutrino and one antineutrino. We omit the details.

### 3. EXISTENCE OF A UNIQUE GROUND STATE.

In the sequel we shall make some of the following additional assumptions on the kernels  $F(\xi_2, \xi_4)$  and  $G(\xi_1, \xi_3)$ .

**Hypothesis 3.1.** *There exists a constant  $K(F, G) > 0$  such that for  $\sigma > 0$*

$$(i) \quad \int_{\Gamma_1 \times \mathbb{R}^3} \frac{|F(\xi_2, \xi_4)|^2}{|\mathbf{p}_4|^2} d\xi_1 d\xi_4 < \infty. (ii) \quad \int_{\Gamma_1 \times \mathbb{R}^3} \frac{|G(\xi_1, \xi_3)|^2}{|\mathbf{p}_3|^2} d\xi_1 d\xi_3 < \infty.$$

$$(iii) \quad \left( \int_{\Gamma_1 \times \{|\mathbf{p}_4| \leq \sigma\}} |F(\xi_2, \xi_4)|^2 d\xi_2 d\xi_4 \right)^{\frac{1}{2}} \leq K(F, G)\sigma.$$

$$(iv) \quad \left( \int_{\Gamma_1 \times \{|\mathbf{p}_3| \leq \sigma\}} |G(\xi_1, \xi_3)|^2 d\xi_1 d\xi_3 \right)^{\frac{1}{2}} \leq K(F, G)\sigma.$$

We then have

**Theorem 3.2.** *Assume that the kernels  $F(\cdot, \cdot)$  and  $G(\cdot, \cdot)$  satisfy Hypothesis 2.1 and 3.1. Then there exists  $g_1 \in (0, g_0]$  such that  $H$  has a unique ground state for  $g \leq g_1$ .*

In order to prove theorem 3.1 it suffices to mimick the proofs given in [5], [11] and [23]. We omit the details.

In [2] fermionic hamiltonian models are considered without any external field. Without any restriction on the strength of the interaction a self-adjoint hamiltonian is defined for which the existence of a ground state is proved. Such a result is an open problem in the case of magnetic fermionic models.

## 4. THE ABSOLUTELY CONTINUOUS SPECTRUM.

As stated in the introduction , in order to specify the absolutely continuous spectrum of  $H$ , we follow the first step of the approach to scattering theory in establishing, for each involved particle, the existence and basic properties of the asymptotic creation and annihilation operators for time  $t$  going to  $\pm\infty$ . The existence of a ground state is quite fundamental in order to get a Fock subrepresentation of the asymptotic canonical anticommutation relations from which we localize the absolutely continuous spectrum of  $H$ .

**4.1. Asymptotic fields.** Let

$$(4.1) \quad \begin{aligned} b_{1,+}^\sharp(t)(f_1) &= e^{itH} e^{-itH_0} b_{1,+}^\sharp(f_1) e^{itH_0} e^{-itH} \\ b_{2,\pm}^\sharp(t)(f_2) &= e^{itH} e^{-itH_0} b_{2,\pm}^\sharp(f_2) e^{itH_0} e^{-itH} \\ b_{3,-}^\sharp(t)(f_3) &= e^{itH} e^{-itH_0} b_{3,-}^\sharp(f_3) e^{itH_0} e^{-itH} \\ b_{4,+}^\sharp(t)(f_4) &= e^{itH} e^{-itH_0} b_{4,+}^\sharp(f_4) e^{itH_0} e^{-itH} . \end{aligned}$$

where, for  $i = 1, 2$ ,  $f_i \in L^2(\Gamma_1)$  and, for  $j = 3, 4$ ,  $f_j \in L^2(\mathbb{R}^3)$ .

The strong limits of  $b_{i,t}^\sharp(\cdot)$  when the time  $t$  goes to  $\pm\infty$  for models in Quantum Field Theory have been considered for fermions and bosons by [33]-[35] and [27]-[31] and , more recently, by [17], [3], [26], [36], [37] and [6] and references therein.

In the sequel we shall make some of the following additional assumptions on the kernels  $F(\xi_2, \xi_4)$  and  $G(\xi_1, \xi_3)$ .

**Hypothesis 4.1.**

$$(i) \quad \int \left| \frac{\partial F}{\partial p_\mu^3}(\xi_2, \xi_4) \right|^2 d\xi_2 d\xi_4 < \infty , \int \left| \frac{\partial G}{\partial p_e^3}(\xi_1, \xi_3) \right|^2 d\xi_1 d\xi_3 < \infty .$$

$$(ii) \quad \int \left| \left( \frac{\partial}{\partial p_\mu^3} \right)^2 F(\xi_2, \xi_4) \right|^2 d\xi_2 d\xi_4 < \infty , \int \left| \left( \frac{\partial}{\partial p_e^3} \right)^2 G(\xi_1, \xi_3) \right|^2 d\xi_1 d\xi_3 < \infty .$$

**Hypothesis 4.2.**

$$(i) \quad \int |\nabla_{p_\mu} F(\xi_2, \xi_4)|^2 d\xi_2 d\xi_4 < \infty , \int |\nabla_{p_e} G(\xi_1, \xi_3)|^2 d\xi_1 d\xi_3 < \infty .$$

$$(ii) \quad \int \left| \frac{\partial^2 F}{\partial p_{\nu_\mu}^1 \partial p_{\nu_\mu}^3}(\xi_2, \xi_4) \right|^2 d\xi_2 d\xi_4 < \infty , \int \left| \frac{\partial^2 G}{\partial p_{\nu_e}^1 \partial p_{\nu_e}^3}(\xi_1, \xi_3) \right|^2 d\xi_1 d\xi_3 < \infty .$$

We then have

**Theorem 4.3.** *Suppose Hypothesis 2.1-Hypothesis 4.2 and  $g \leq g_0$ . Let  $f_1, f_2 \in L^2(\Gamma_1)$  and  $f_3, f_4 \in L^2(\mathbb{R}^3)$ . Then the following asymptotic fields*

$$(4.2) \quad \begin{aligned} b_{1,+,\pm\infty}^\sharp(f_1) &:= s\text{-}\lim_{t \rightarrow \pm\infty} b_{1,+}^\sharp(t)(f_1) \\ b_{2,\pm,\pm\infty}^\sharp(f_2) &:= s\text{-}\lim_{t \rightarrow \pm\infty} b_{2,\pm}^\sharp(t)(f_2) \\ b_{3,-,\pm\infty}^\sharp(f_3) &:= s\text{-}\lim_{t \rightarrow \pm\infty} b_{3,-}^\sharp(t)(f_3) \\ b_{4,+,\pm\infty}^\sharp(f_4) &:= s\text{-}\lim_{t \rightarrow \pm\infty} b_{4,+}^\sharp(t)(f_4) . \end{aligned}$$

*exist.*

*Proof.* The norms of the  $b_{i,t}^\sharp(f)$ 's are uniformly bounded with respect to  $t$ . Hence, in order to prove theorem 4.1 it suffices to prove the existence of the strong limits on  $D(H) = D(H_0)$  with smooth  $f$ .

**Strong limits of  $b_{1,+}^\sharp(f_1)$  and  $b_{2,\pm}^\sharp(f_2)$  .**

Let

$$(4.3) \quad \mathfrak{D} = \{f \in l^2(\Gamma_1) | f(s, n, \dots) \in C_0^\infty(\mathbb{R}^2 \setminus \{0\}) \text{ for all } s \text{ and } n, \text{ and} \\ f(\cdot, n, \dots) = 0 \text{ for all but finitely many } n\}.$$

Let  $f_1, f_2 \in \mathfrak{D}$  . According to [27, lemma1] we have

$$(4.4) \quad b_{1,+}^\sharp(f_1)D(H) \subset D(H) \quad \text{and} \quad b_{2,\pm}^\sharp(f_2)D(H) \subset D(H).$$

Moreover we have

$$(4.5) \quad \begin{aligned} e^{itH_0} b_{1,+}^\sharp(f_1) e^{-itH_0} \Psi &= b_{1,+}^\sharp(e^{itE^e} f_1) \Psi, \\ e^{itH_0} b_{2,\pm}^\sharp(f_2) e^{-itH_0} \Psi &= b_{2,\pm}^\sharp(e^{itE^\mu} f_2) \Psi. \end{aligned}$$

where  $\Psi \in D(H)$ .

Let us first prove the existence of  $b_{1,+}^\sharp(f_1)$ .

Let  $\Psi, \Phi \in D(H)$  and  $f_{1,t}(\xi_1) = (e^{-itE^e} f_1)(\xi_1)$ . By (4.4),(4.5) and the strong differentiability of  $e^{itH}$  we get

$$(4.6) \quad \begin{aligned} &(\Phi, b_{1,+}^\sharp(f_1) \Psi) - (\Phi, b_{1,+}^\sharp(f_1) \Psi) \\ &= g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{1,+}^\sharp(f_1) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{1,+}^\sharp(f_{1,t})] e^{-itH} \Psi) dt \end{aligned}$$

By using the usual canpnicol anticommutation relations (CAR) (see(A.4)) we easily get for all  $\Psi \in D(H)$

$$(4.7) \quad \begin{aligned} &[H_I^{(1)}, b_{1,+}^\sharp(f_{1,t})] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ &(\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) W^{(\overline{\nu}_e)}(\xi_3)) \\ &\left. \overline{f_{1,t}(\xi_1)} F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_4) b_-^*(\xi_3) b_+(\xi_2) \Psi. \right) \end{aligned}$$

$$(4.8) \quad \begin{aligned} &[H_I^{(2)}, b_{1,+}^\sharp(f_{1,t})] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ &(\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) W^{(\overline{\nu}_e)}(\xi_3)) \\ &\left. \overline{f_{1,t}(\xi_1)} F(\xi_2, \xi_4) G(\xi_1, \xi_3) \right) b_+^*(\xi_4) b_-^*(\xi_2) b_-^*(\xi_3) \Psi. \end{aligned}$$

$$(4.9) \quad [(H_I^{(1)})^*, b_{1,+}^\sharp(f_{1,t})] \Psi = [(H_I^{(2)})^*, b_{1,+}^\sharp(f_{1,t})] \Psi = 0$$

where  $\overline{U} = U^\dagger \gamma^0$ .

Similarly we get

$$(4.10) \quad \begin{aligned} &(\Phi, b_{1,+}^*(f_1) \Psi) - (\Phi, b_{1,+}^*(f_1) \Psi) \\ &= g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{1,+}^*(f_1) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{1,+}^*(f_{1,t})] e^{-itH} \Psi) dt \end{aligned}$$

with

$$(4.11) \quad [H_I^{(1)}, b_{1,+}^*(f_{1,t})] \Psi = [H_I^{(2)}, b_{1,+}^*(f_{1,t})] \Psi = 0$$

and



$$(4.12) \quad \begin{aligned} & \left[ (H_I^{(1)})^*, b_{1,+}^*(f_{1,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. (\overline{W}^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)) (\overline{U}^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)) \right) \\ & \overline{F}(\xi_2, \xi_4) \overline{G}(\xi_1, \xi_3) f_{1,t}(\xi_1) b_+^*(\xi_2) b_-(\xi_3) b_+(\xi_4) \Psi. \end{aligned}$$

$$(4.13) \quad \begin{aligned} & \left[ (H_I^{(2)})^*, b_{1,+}^*(f_{1,t}) \right] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. (\overline{W}^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)) (W^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)) \right) \\ & \overline{F}(\xi_2, \xi_4) \overline{G}(\xi_1, \xi_3) f_{1,t}(\xi_1) b_-(\xi_2) b_-(\xi_3) b_+(\xi_4) \Psi. \end{aligned}$$

By (4.6) and (4.10), in order to prove the existence of  $b_{1,+,\pm\infty}^\sharp(f_1)$ , we have to estimate

$$e^{itH} [H_I, b_{1,+}(f_{1,t})] e^{-itH} \Psi$$

and

$$e^{itH} [H_I, b_{1,+}^*(f_{1,t})] e^{-itH} \Psi$$

for large  $|t|$ .

By (B.5), the  $N_\tau$  estimates (see [20] and [7, Proposition 3.7]), (A.8), (A.11) and (A.13) we get

$$(4.14) \quad \begin{aligned} & \left\| e^{itH} [H_I^{(1)}, b_{1,+}(f_{1,t})] e^{-itH} \Psi \right\| \leq \\ & C \left( \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U^{(e)}(x^2, \xi_1) f_{1,t}(\xi_1) \overline{G}(\xi_1, \xi_3) \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \| (N_{\mu_-} + 1)^{\frac{1}{2}} e^{-itH} \Psi \|. \end{aligned}$$

and

$$(4.15) \quad \begin{aligned} & \left\| e^{itH} [H_I^{(2)}, b_{1,+}(f_{1,t})] e^{-itH} \Psi \right\| \leq \\ & C \left( \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U^{(e)}(x^2, \xi_1) f_{1,t}(\xi_1) \overline{G}(\xi_1, \xi_3) \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \| (N_{\mu_+} + 1)^{\frac{1}{2}} e^{-itH} \Psi \|. \end{aligned}$$

By (2.18) and (2.19) we have

$$(4.16) \quad \|H_I \Psi\| \leq a \|H_0 \phi\| + b \|\Psi\|$$

with

$$a = 4 \frac{C}{M} \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)}$$

and

$$b = 2C \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)}$$

Hence we obtain

$$(4.17) \quad \|H_0 \Psi\| \leq \tilde{a} \|H \Psi\| + \tilde{b} \|\Psi\|$$

with

$$\tilde{a} = \frac{1}{1 - g_0 a} \quad \text{and} \quad \tilde{b} = \frac{g_0 b}{1 - g_0 a}$$

Therefore we have

$$(4.18) \quad \begin{aligned} \|(N_e + 1)^{\frac{1}{2}} e^{-itH} \Psi\| &\leq \frac{1}{m_e} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_e) \|\Psi\|) \\ \|(N_{\mu\pm} + 1)^{\frac{1}{2}} e^{-itH} \Psi\| &\leq \frac{1}{m_\mu} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_\mu) \|\Psi\|). \end{aligned}$$

where  $m_\mu$  is the mass of the muon.

Hence we get

$$(4.19) \quad \begin{aligned} &\|e^{itH} [H_I, b_{1,+}(f_{1,t})] e^{-itH} \Psi\| \leq \\ &2C \left( \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U^{(e)}(x^2, \xi_1) f_{1,t}(\xi_1) \overline{G(\xi_1, \xi_3)} \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ &\|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_\mu} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_\mu) \|\Psi\|). \end{aligned}$$

Moreover we have

$$(4.20) \quad \begin{aligned} &\int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U^{(e)}(x^2, \xi_1) f_{1,t}(\xi_1) \overline{G(\xi_1, \xi_3)} \right\|_{\mathbb{C}^4}^2 \right) \\ &= \sum_{j=1}^4 \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U_j^{(e)}(x^2, \xi_1) e^{-itE_n^{(e)}(p_e^3)} f_1(\xi_1) \overline{G(\xi_1, \xi_3)} \right\|_{\mathbb{C}^4}^2 \right) \end{aligned}$$

where  $(\bigcup_{j=1}^4 U_j^{(e)}(x^2, \xi_1))$  are the four components of the vectors (A.8) and (A.11)  $\in \mathbb{C}^4$ .

Note that

$$(4.21) \quad e^{-itE_n^{(e)}(p_e^3)} = \frac{1}{it} \frac{E_n^{(e)}(p_e^3)}{p_e^3} \frac{d}{dp_e^3} e^{-itE_n^{(e)}(p_e^3)}.$$

By (4.20) and (4.21), by a two-fold partial integration with respect to  $p_e^3$  and by Hypothesis 4.1 one can show that there exists for every  $j$  a function, denoted by  $H_j^{(e)}(\xi_1, \xi_3)$ , such that

$$(4.22) \quad \begin{aligned} &\sum_{j=1}^4 \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U_j^{(e)}(x^2, \xi_1) e^{-itE_n^{(e)}(p_e^3)} f_1(\xi_1) \overline{G(\xi_1, \xi_3)} \right\|_{\mathbb{C}^4}^2 \right) \\ &= \sum_{j=1}^4 \frac{1}{t^4} \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U_j^{(e)}(x^2, \xi_1) H_j^{(e)}(\xi_1, \xi_3) e^{-itE_n^{(e)}(p_e^3)} \right\|_{\mathbb{C}^4}^2 \right) \\ &\leq C_{f_1} \frac{1}{t^4} \sum_{j=1}^4 \left( \int d\xi_1 d\xi_3 \chi_{f_1}(\xi_1) |H_j^{(e)}(\xi_1, \xi_3)|^2 \right) < \infty. \end{aligned}$$

Here  $\chi_{f_1}(\cdot)$  is the characteristic function of the support of  $f_1(\cdot)$  and (A.13) is used.

By (4.6) and (4.19)-(4.22) the strong limits of  $b_{1,+}(f_1)$  on  $\mathfrak{F}$  when  $t$  goes to  $\pm\infty$  and for all  $f_1 \in L^2(\Gamma_1)$  exist for every  $g \leq g_0$ .

By (4.11)-(4.13) and by mimicking the proof of (4.14) and (4.15) we get

$$(4.23) \quad \begin{aligned} &\text{Sup} \left( \left\| e^{itH} [(H_I^{(1)})^*, b_{1,+}^*(f_{1,t})] e^{-itH} \Psi \right\|, \left\| e^{itH} [(H_I^{(2)})^*, b_{1,+}^*(f_{1,t})] e^{-itH} \Psi \right\| \right) \\ &\leq C \left( \int dx^2 \left( \int d\xi_3 \left\| \int d\xi_1 U^{(e)}(x^2, \xi_1) f_{1,t}(\xi_1) \overline{G(\xi_1, \xi_3)} \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \\ &\times \|F(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_\mu} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_\mu) \|\Psi\|). \end{aligned}$$

It follows from (4.10) and (4.20)-(4.23) that the strong limits of  $b_{1,+t}^*(f_1)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_1 \in L^2(\Gamma_1)$  and for every  $g \leq g_0$ .

We now consider the existence of  $b_{2,\epsilon,\pm\infty}^\sharp(f_2)$

Let  $\Psi, \Phi \in D(H)$  and  $f_{2,t}(\xi_2) = (e^{-itE^\mu} f_2)(\xi_2)$  with  $f_2 \in \mathfrak{D}$ . By (4.4), (4.5) and the strong differentiability of  $e^{itH}$  we get

$$(4.24) \quad \begin{aligned} & (\Phi, b_{2,+T}(f)\Psi) - (\Phi, b_{2,+T_0}(f_2)\Psi) \\ &= g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{2,+t}(f_2)\Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{2,+}(f_{2,t})] e^{-itH} \Psi) dt \end{aligned}$$

with

$$(4.25) \quad [H_I^{(1)}, b_{2,+}(f_{2,t})] \Psi = [H_I^{(2)}, b_{2,+}(f_{2,t})] \Psi = [(H_I^{(2)})^*, b_{2,+}(f_{2,t})] \Psi = 0$$

and

$$(4.26) \quad \begin{aligned} & [(H_I^{(1)})^*, b_{2,+}(f_{2,t})] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. \overline{W^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)} \overline{U^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)} \right) \\ & \overline{F(\xi_2, \xi_4) G(\xi_1, \xi_3) f_{2,t}(\xi_2) b_-(\xi_3) b_+(\xi_1) b_+(\xi_4)} \Psi. \end{aligned}$$

Similarly we obtain

$$(4.27) \quad \begin{aligned} & \|e^{itH} [H_I, b_{2,+}(f_{2,t})] e^{-itH} \Psi\| \leq \\ & C \left( \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 U^{(\mu)}(x^2, \xi_2) F(\xi_2, \xi_4) f_{2,t}(\xi_1) \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \|(N_e + 1)^{\frac{1}{2}} e^{-itH} \Psi\|. \end{aligned}$$

It follows from (4.16)-(4.18) that

$$(4.28) \quad \|(N_e + 1)^{\frac{1}{2}} e^{-itH} \Psi\| \leq \frac{1}{m_e} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_e) \|\Psi\|).$$

Hence

$$(4.29) \quad \begin{aligned} & \|e^{itH} [H_I, b_{2,+}(f_{2,t})] e^{-itH} \Psi\| \leq \\ & C \left( \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 U^{(\mu)}(x^2, \xi_2) F(\xi_2, \xi_4) f_{2,t}(\xi_2) \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_e) \|\Psi\|). \end{aligned}$$

Moreover we have

$$(4.30) \quad \begin{aligned} & \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 U^{(\mu)}(x^2, \xi_2) f_{2,t}(\xi_2) F(\xi_2, \xi_4) \right\|_{\mathbb{C}^4}^2 \right. \\ & \left. = \sum_{j=1}^4 \int dx^2 \left( \int d\xi_4 \left| \int d\xi_2 U_j^{(\mu)}(x^2, \xi_2) e^{-itE_n^{(\mu)}(p_\mu^3)} f_2(\xi_2) F(\xi_2, \xi_4) \right|^2 \right) \right)^2. \end{aligned}$$

where  $(U_{j=1}^4 U_j^{(\mu)}(x^2, \xi_2))$  are the four components of the vectors (A.8) and (A.11)  $\in \mathbb{C}^4$  for  $\alpha = \mu$ .

By (4.30), by a two-fold partial integration with respect to  $p_\mu^3$  and by Hypothesis 4.1 one can show that there exists for every  $j$  a function, denoted by  $H_j^{(\mu)}(\xi_2, \xi_4)$ , such that

$$\begin{aligned}
(4.31) \quad & \sum_{j=1}^4 \int dx^2 \left( \int d\xi_4 \left| \int d\xi_2 U_j^{(\mu)}(x^2, \xi_2) e^{-itE_n^{(\mu)}(p_\mu^3)} f_2(\xi_2) F(\xi_2, \xi_4) \right|^2 \right) \\
& = \sum_{j=1}^4 \frac{1}{t^4} \int dx^2 \left( \int d\xi_4 \left| \int d\xi_1 U_j^{(\mu)}(x^2, \xi_2) H_j^{(\mu)}(\xi_2, \xi_4) e^{-itE_n^{(\mu)}(p_\mu^3)} \right|^2 \right) \\
& \leq C_{f_2} \frac{1}{t^4} \sum_{j=1}^4 \left( \int d\xi_2 d\xi_4 \chi_{f_2}(\xi_2) |H_j^{(\mu)}(\xi_2, \xi_4)|^2 \right) < \infty.
\end{aligned}$$

Here  $\chi_{f_2}(\cdot)$  is the characteristic function of the support of  $f_2(\cdot)$  and (A.13) is used. Similarly we have

$$\begin{aligned}
(4.32) \quad & (\Phi, b_{2,+}^*(f_2) \Psi) - (\Phi, b_{2,+}^*(f_2) \Psi) \\
& = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{2,+}^*(f_2) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{2,+}^*(f_2,t)] e^{-itH} \Psi) dt
\end{aligned}$$

with

$$(4.33) \quad \left[ (H_I^{(1)})^*, b_{2,+}^*(f_2,t) \right] \Psi = \left[ H_I^{(2)}, b_{2,+}^*(f_2,t) \right] \Psi = \left[ (H_I^{(2)})^*, b_{2,+}^*(f_2,t) \right] \Psi = 0$$

and

$$\begin{aligned}
(4.34) \quad & \left[ (H_I^{(1)})^*, b_{2,+}^*(f_2,t) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\
& \left. (\overline{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) W^{(\overline{\nu}_e)}(\xi_3)) G(\xi_1, \xi_3) (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) \right) \\
& F(\xi_2, \xi_4) G(\xi_1, \xi_3) f_{2,t}(\xi_2) b_+^*(\xi_4) b_+^*(\xi_1) b_-^*(\xi_3) \Psi.
\end{aligned}$$

Similarly we obtain

$$\begin{aligned}
(4.35) \quad & \| e^{itH} [H_I, b_{2,+}^*(f_2,t)] e^{-itH} \Psi \| \leq \\
& C \left( \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 U^{(\mu)}(x^2, \xi_2) F(\xi_2, \xi_4) f_{2,t}(\xi_1) \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\
& \| G(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_e) \| \Psi \|).
\end{aligned}$$

It follows from (4.29), (4.31) and (4.35) that the strong limits of  $b^\sharp(2, +, t)(f_2)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_2 \in L^2(\Gamma_1 \times \mathbb{R}^3)$  and for every  $g \leq g_0$ .

Let us now consider the strong limits of  $b^\sharp(2, -, t)(f_2)$ .

We have for all  $f_2 \in \mathfrak{D}$

$$\begin{aligned}
(4.36) \quad & (\Phi, b_{2,-}^*(f) \Psi) - (\Phi, b_{2,-}^*(f_2) \Psi) \\
& = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{2,-}^*(f_2) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{2,-}^*(f_2,t)] e^{-itH} \Psi) dt
\end{aligned}$$

with

$$(4.37) \quad \left[ H_I^{(1)}, b_{2,-}^*(f_2,t) \right] \Psi = \left[ (H_I^{(1)})^*, b_{2,-}^*(f_2,t) \right] \Psi = \left[ (H_I^{(2)})^*, b_{2,-}^*(f_2,t) \right] \Psi = 0$$

(4.38)

$$\begin{aligned} & \left[ (H_I^{(2)}), b_{2,-}(f_{2,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ & \left. (\bar{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) W^{(\bar{\nu}_e)}(\xi_3)) G(\xi_1, \xi_3) (\bar{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2)) \right) \\ & F(\xi_2, \xi_4) G(\xi_1, \xi_3) f_{2,t}(\xi_2) b_+^*(\xi_4) b_+^*(\xi_1) b_-^*(\xi_3) \Psi. \end{aligned}$$

By mimicking the proofs given above we get

$$\begin{aligned} & \| e^{itH} [H_I, b_{2,-}(f_{2,t})] e^{-itH} \Psi \| \leq \\ (4.39) \quad & C \left( \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 W^{(\mu)}(x^2, \xi_2) F(\xi_2, \xi_4) \overline{f_{2,t}(\xi_1)} \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \| G(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_e) \| \Psi \|). \end{aligned}$$

and

$$\begin{aligned} & \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 W^{(\mu)}(x^2, \xi_2) \overline{f_{2,t}(\xi_2)} F(\xi_2, \xi_4) \right\|_{\mathbb{C}^4}^2 \right) \\ (4.40) \quad & = \sum_{j=1}^4 \int dx^2 \left( \int d\xi_4 \left| \int d\xi_2 W_j^{(\mu)}(x^2, \xi_2) e^{-itE_n^{(\mu)}(p_\mu^3)} \overline{f_2(\xi_2)} F(\xi_2, \xi_4) \right|^2 \right). \end{aligned}$$

where  $(\bigcup_{j=1}^4 W_j^{(\mu)}(x^2, \xi_2))$  are the four components of the vectors (A.14) - (A.16)  $\in \mathbb{C}^4$  for  $\alpha = \mu$ .

By (4.40), by a two-fold partial integration with respect to  $p_\mu^3$  and by Hypothesis 4.1 one can show that there exists for every  $j$  a function, denoted by  $\tilde{H}_j^{(\mu)}(\xi_2, \xi_4)$ , such that

$$\begin{aligned} & \sum_{j=1}^4 \int dx^2 \left( \int d\xi_4 \left| \int d\xi_2 W_j^{(\mu)}(x^2, \xi_2) e^{-itE_n^{(\mu)}(p_\mu^3)} \overline{f_2(\xi_2)} F(\xi_2, \xi_4) \right|^2 \right) \\ (4.41) \quad & = \sum_{j=1}^4 \frac{1}{t^4} \int dx^2 \left( \int d\xi_4 \left| \int d\xi_2 W_j^{(\mu)}(x^2, \xi_2) \tilde{H}_j^{(\mu)}(\xi_2, \xi_4) e^{-itE_n^{(\mu)}(p_\mu^3)} \right|^2 \right) \\ & \leq C_{f_2} \frac{1}{t^4} \sum_{j=1}^4 \left( \int d\xi_2 d\xi_4 \chi_{f_2}(\xi_2) |\tilde{H}_j^{(\mu)}(\xi_2, \xi_4)|^2 \right) < \infty. \end{aligned}$$

Here  $\chi_{f_2}(\cdot)$  is the characteristic function of the support of  $f_2(\cdot)$  and (A.17) is used.

It follows from (4.36), (4.39)-(4.41) that the strong limits of  $b(2, -, t)(f_2)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_2 \in L^2(\Gamma_1 \times \mathbb{R}^3)$  and for every  $g \leq g_0$ .

We now have for all  $f_2 \in \mathfrak{D}$

$$\begin{aligned} & (\Phi, b_{2,-,T}^*(f_2) \Psi) - (\Phi, b_{2,-,T_0}^*(f_2) \Psi) \\ (4.42) \quad & = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{2,-,t}^*(f_2) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{2,-}^*(f_{2,t})] e^{-itH} \Psi) dt \end{aligned}$$

with

$$(4.43) \quad \left[ H_I^{(1)}, b_{2,-}^*(f_{2,t}) \right] \Psi = \left[ (H_I^{(1)})^*, b_{2,-}^*(f_{2,t}) \right] \Psi = \left[ H_I^{(2)}, b_{2,-}^*(f_{2,t}) \right] \Psi = 0$$

$$(4.44) \quad \begin{aligned} & \left[ (H_I^{(2)})^*, b_{2,-}^*(f_{2,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. (\overline{W}^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)) (\overline{W}^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)) \right) \\ & \overline{F(\xi_2, \xi_4)} \overline{G(\xi_1, \xi_3)} f_{2,t}(\xi_2) b_-(\xi_3) b_+(\xi_1) b_+(\xi_4) \Psi. \end{aligned}$$

Similarly to (4.39) we get

$$(4.45) \quad \begin{aligned} & \| e^{itH} [H_I, b_{2,-}^*(f_{2,t})] e^{-itH} \Psi \| \leq \\ & C \left( \int dx^2 \left( \int d\xi_4 \left\| \int d\xi_2 W^{(\mu)}(x^2, \xi_2) F(\xi_2, \xi_4) \overline{f_{2,t}(\xi_1)} \right\|_{\mathbb{C}^4}^2 \right) \right)^{\frac{1}{2}} \times \\ & \| G(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_e) \| \Psi \|). \end{aligned}$$

It follows from (4.43), (4.45), (4.40) and (4.41) that the strong limits of  $b(2, -, t)^*(f_2)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_2 \in L^2(\Gamma_1 \times \mathbb{R}^3)$  and for every  $g \leq g_0$ .

**Strong limits of  $b_{3,-,t}^\sharp(f_3)$  and  $b_{4,+,t}^\sharp(f_4)$ .**

Let

$$(4.46) \quad \mathfrak{D}' = \{f(\cdot) \in C_0^\infty(\mathbb{R}^3 \setminus \{(0, 0, p^3)\}); p^3 \in \mathbb{R}\}.$$

Let  $f_1, f_2 \in \mathfrak{D}'$ . According to [27, lemma1] we have

$$(4.47) \quad b_{4,+}^\sharp(f_4) D(H) \subset D(H) \quad \text{and} \quad b_{3,-}^\sharp(f_3) D(H) \subset D(H).$$

Moreover we have

$$(4.48) \quad \begin{aligned} e^{itH_0} b_{3,-}^\sharp(f_3) e^{-itH_0} \Psi &= b_{3,-}^\sharp(e^{it|p_3|} f_3) \Psi, \\ e^{itH_0} b_{4,+}^\sharp(f_4) e^{-itH_0} \Psi &= b_{4,+}^\sharp(e^{it|p_4|} f_4) \Psi. \end{aligned}$$

where  $\Psi \in D(H)$ .

Let  $\Psi, \Phi \in D(H)$  and  $f_{j,t}(\xi_j) = (e^{-it|p_j|} f_j)(\xi_j)$  where  $j = 3, 4$ . By (4.4), (4.5) and the strong differentiability of  $e^{itH}$  we get

$$(4.49) \quad \begin{aligned} & (\Phi, b_{3,-,T}(f_3) \Psi) - (\Phi, b_{3,-,T_0}(f_1) \Psi) \\ &= g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{3,-,t}(f_3) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{3,-}(f_{3,t})] e^{-itH} \Psi) dt \end{aligned}$$

By using the usual anticommutation relations (CAR) (see (A.4) and (B.4)) we easily get for all  $\Psi \in D(H)$

$$(4.50) \quad \begin{aligned} & \left[ H_I^{(1)}, b_{3,-}(f_{3,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ & \left. (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) W^{(\bar{\nu}_e)}(\xi_3)) \right) \\ & \overline{f_{3,t}(\xi_3)} F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_4) b_+^*(\xi_1) b_+(\xi_2) \Psi. \end{aligned}$$

$$(4.51) \quad \begin{aligned} & \left[ H_I^{(2)}, b_{3,-}(f_{3,t}) \right] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ & \left. (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) W^{(\bar{\nu}_e)}(\xi_3)) \right) \\ & \overline{f_{3,t}(\xi_3)} F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_4) b_-^*(\xi_2) b_+^*(\xi_1) \Psi. \end{aligned}$$

and

$$(4.52) \quad \left[ (H_I^{(1)})^*, b_{3,-}(f_{3,t}) \right] \Psi = \left[ (H_I^{(2)})^*, b_{3,-}(f_{3,t}) \right] \Psi = 0$$

By (B.5) we get

$$(4.53) \quad \left\| \left[ H_I^{(1)}, b_{3,-}(f_{3,t}) \right] \Psi \right\| \leq \int dx^2 \left( \int d\xi_1 \left| (\overline{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) \int d\xi_3 e^{-ip_3^2 x^2} W^{(\overline{\nu}_e)}(\xi_3) G(\xi_1, \xi_3) \overline{f_{3,t}(\xi_3)} \right|^2 \right)^{\frac{1}{2}} \left\| \int d\xi_2 d\xi_4 e^{-ip_4^2 x^2} (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) b_+^*(\xi_4) b_+(\xi_2) \Psi \right\|.$$

and

$$(4.54) \quad \left\| \left[ H_I^{(2)}, b_{3,-}(f_{3,t}) \right] \Psi \right\| \leq \int dx^2 \left( \int d\xi_1 \left| (\overline{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) \int d\xi_3 e^{-ip_3^2 x^2} W^{(\overline{\nu}_e)}(\xi_3) G(\xi_1, \xi_3) \overline{f_{3,t}(\xi_3)} \right|^2 \right)^{\frac{1}{2}} \left\| \int d\xi_2 d\xi_4 e^{-ip_4^2 x^2} (\overline{W}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) b_+^*(\xi_4) b_-^*(\xi_2) \Psi \right\|.$$

Moreover we have

$$(4.55) \quad \int d\xi_1 \left\| \int d\xi_3 e^{-ip_3^2 x^2} W^{(\overline{\nu}_e)}(\xi_3) \overline{f_{1,t}(\xi_1)} G(\xi_1, \xi_3) \right\|_{\mathbb{C}^4}^2 = \sum_{j=1}^4 \int d\xi_1 \left| \int d\xi_3 e^{-ip_3^2 x^2} W_j^{(\overline{\nu}_e)}(\xi_3) (e^{it|\mathbf{p}_3|} \overline{f_3(\xi_3)} G(\xi_1, \xi_3)) \right|^2.$$

where  $(\bigcup_{j=1}^4 W_j^{(\overline{\nu}_e)}(\xi_3))$  are the four components of the vector (B.12)  $\in \mathbb{C}^4$ .

By a two-fold partial integration with respect to  $p^3$  and  $p^1$  and by Hypothesis 4.2 one can show that there exists for every  $j$  a function, denoted by  $H_j^{(\overline{\nu}_e)}(\xi_1, \xi_3)$ , such that

$$(4.56) \quad \begin{aligned} & \sum_{j=1}^4 \int d\xi_1 \left| \int d\xi_3 e^{-ip_3^2 x^2} W_j^{(\overline{\nu}_e)}(\xi_3) e^{it|\mathbf{p}_3|} \overline{f_3(\xi_3)} G(\xi_1, \xi_3) \right|^2 \\ &= \sum_{j=1}^4 \frac{1}{t^4} \int d\xi_1 \left| \int d\xi_3 e^{-ip_3^2 x^2} W_j^{(\overline{\nu}_e)}(\xi_3) H_j^{(\overline{\nu}_e)}(\xi_1, \xi_3) e^{it|\mathbf{p}_3|} \right|^2 \\ &\leq C_{f_3}^2 \frac{1}{t^4} \sum_{j=1}^4 \int d\xi_1 d\xi_3 \chi_{f_3}(\xi_3) e^{it|\mathbf{p}_3|} |H_j^{(\overline{\nu}_e)}(\xi_1, \xi_3)|^2 < \infty. \end{aligned}$$

Here  $\chi_{f_3}(\cdot)$  is the characteristic function of the support of  $f_3(\cdot)$ .

By the  $N_\tau$  estimates and by (4.18), (A.13), (A.17) and (B.14) it follows from (4.52)-(4.56) that, for every  $\Psi \in D(H)$ ,

$$(4.57) \quad \begin{aligned} & \| e^{itH} [H_I, b_{3,-}(f_{3,t})] e^{-itH} \Psi \| \leq \\ & CC_{f_3} \frac{1}{t^2} \left( \sum_{j=1}^4 \int d\xi_1 d\xi_3 \chi_{f_3}(\xi_3) |H_j^{(\overline{\nu}_e)}(\xi_1, \xi_3)|^2 \right)^{\frac{1}{2}} \times \\ & \| F(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_\mu} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_\mu) \| \Psi \|) \end{aligned}$$

Furthermore we have

$$(4.58) \quad \begin{aligned} & (\Phi, b_{3,-}^*(f_3)\Psi) - (\Phi, b_{3,-}^*(f_3)\Psi) \\ & = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{3,-}^*(f_3)\Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{3,-}^*(f_{3,t})] e^{-itH} \Psi) dt \end{aligned}$$

with

$$(4.59) \quad \begin{aligned} & \left[ (H_I^{(1)})^*, b_{3,-}^*(f_{3,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. (\overline{W}^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)) (\overline{U}^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)) \right) \\ & \overline{F}(\xi_2, \xi_4) G(\xi_1, \xi_3) f_{3,t}(\xi_2) b_+^*(\xi_2) b_+(\xi_1) b_+(\xi_4) \Psi. \end{aligned}$$

$$(4.60) \quad \begin{aligned} & \left[ (H_I^{(2)})^*, b_{3,-}^*(f_{3,t}) \right] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. (\overline{W}^{(\bar{\nu}_e)}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1)) (W^{(\mu)}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4)) \right) \\ & \overline{F}((\xi_2, \xi_4) G(\xi_1, \xi_3) f_{3,t} b_+(\xi_1) b_-(\xi_2) b_+(\xi_4) \Psi. \end{aligned}$$

and

$$(4.61) \quad \left[ (H_I^{(1)}), b_{3,-}^*(f_{3,t}) \right] \Psi = \left[ (H_I^{(2)}), b_{3,-}^*(f_{3,t}) \right] \Psi = 0$$

By adapting the proof of (4.53)-(4.57) to (4.58)-(4.61) we obtain

$$(4.62) \quad \begin{aligned} & \| e^{itH} [H_I, b_{3,-}^*(f_{3,t})] e^{-itH} \Psi \| \\ & \leq CC_{f_3} \frac{1}{t^2} \left( \sum_{j=1}^4 \int d\xi_1 d\xi_3 \chi_{f_3}(\xi_3) |H_j^{(\bar{\nu}_e)}(\xi_1, \xi_3)|^2 \right)^{\frac{1}{2}} \times \\ & \times \| F(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_\mu} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_\mu) \| \Psi \|). \end{aligned}$$

Here  $\chi_{f_3}(\cdot)$  is the characteristic function of the support of  $f_3(\cdot)$ .

It follows from (4.49), (4.47), (4.58) and (4.62) that the strong limits of  $b_{3,-,t}^\sharp(f_3)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_3 \in L^2(\mathbb{R}^3)$  and for every  $g \leq g_0$ .

$$(4.63) \quad \begin{aligned} & (\Phi, b_{4,+}(f_1)\Psi) - (\Phi, b_{4,+}(f_1)\Psi) \\ & = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{4,+}(f_4)\Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{4,+}(f_{4,t})] e^{-itH} \Psi) dt \end{aligned}$$

By using the usual canonical anticommutation relations (CAR) (see (A.4) and (B.4)) we easily get for all  $\Psi \in D(H)$

$$(4.64) \quad \begin{aligned} & \left[ H_I^{(1)}, b_{4,+}(f_{4,t}) \right] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ & \left. (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma_\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma^\alpha (1 - \gamma_5) W^{(\bar{\nu}_e)}(\xi_3)) \right) \\ & \overline{f_{4,t}}(\xi_4) F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_+^*(\xi_1) b_-^*(\xi_3) b_+(\xi_2) \Psi. \end{aligned}$$



$$(4.65) \quad \begin{aligned} & [H_I^{(2)}, b_{4,+}(f_{4,t})] \Psi = - \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{-ix^2 r^2} \right. \\ & \left. (\overline{U}^{(\nu_\mu)}(\xi_4) \gamma^\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2)) (\overline{U}^{(e)}(x^2, \xi_1) \gamma_\alpha (1 - \gamma_5) W^{(\overline{\nu_e})}(\xi_3)) \right) \\ & \overline{f_{4,t}(\xi_4)} F(\xi_2, \xi_4) G(\xi_1, \xi_3) b_-^*(\xi_2) b_+^*(\xi_1) b_-^*(\xi_3) \Psi. \end{aligned}$$

and

$$(4.66) \quad [(H_I^{(1)})^*, b_{4,+}(f_{4,t})] \Psi = [(H_I^{(2)})^*, b_{4,+}(f_{4,t})] \Psi = 0$$

By (B.5) we get

$$(4.67) \quad \begin{aligned} & \left\| [H_I^{(1)}, b_{4,+}(f_{4,t})] \Psi \right\| \leq \\ & \int dx^2 \left( \int d\xi_2 \left| \left\langle \int d\xi_4 U^{(\nu_\mu)}(\xi_4) f_{4,t}(\xi_4) e^{ip_4^2 x^2} \overline{F(\xi_2, \xi_4)}, \gamma^0 \gamma^\alpha (1 - \gamma_5) U^{(\mu)}(x^2, \xi_2) \right\rangle \right|^2 \right)^{\frac{1}{2}} \\ & \left\| \int d\xi_1 d\xi_3 e^{-ip_3^2 x^2} \left\langle U^{(e)}(x^2, \xi_1), \gamma^0 \gamma_\alpha (1 - \gamma_5) W^{(\overline{\nu_e})}(\xi_3) G(\xi_1, \xi_3) \right\rangle b_+^*(\xi_1) b_-^*(\xi_3) \Psi \right\|. \end{aligned}$$

where  $\langle \cdot, \cdot \rangle$  is the scalar product in  $\mathbb{C}^4$ .

and

$$(4.68) \quad \begin{aligned} & \left\| [H_I^{(2)}, b_{4,+}(f_{4,t})] \Psi \right\| \leq \\ & \int dx^2 \left( \int d\xi_2 \left| \left\langle \int d\xi_4 U^{(\nu_\mu)}(\xi_4) f_{4,t}(\xi_4) e^{ip_4^2 x^2} \overline{F(\xi_2, \xi_4)}, \gamma^0 \gamma^\alpha (1 - \gamma_5) W^{(\mu)}(x^2, \xi_2) \right\rangle \right|^2 \right)^{\frac{1}{2}} \\ & \left\| \int d\xi_1 d\xi_3 e^{-ip_3^2 x^2} \left\langle U^{(e)}(x^2, \xi_1), \gamma^0 \gamma_\alpha (1 - \gamma_5) W^{(\overline{\nu_e})}(\xi_3) G(\xi_1, \xi_3) \right\rangle b_+^*(\xi_1) b_-^*(\xi_3) \Psi \right\|. \end{aligned}$$

By adapting the proof of (4.57) to (4.67) and (4.68) one can show that there exists for every  $j$  a function, denoted by  $H^{\nu_\mu}(\xi_2, \xi_4)$ , such that

$$(4.69) \quad \begin{aligned} & \| e^{itH} [H_I, b_{4,+}(f_{4,t})] e^{-itH} \Psi \| \leq \\ & CC_{f_4} \frac{1}{t^2} \left( \sum_{j=1}^4 \int d\xi_2 d\xi_4 \chi_{f_4}(\xi_4) |H_j^{(\nu_\mu)}(\xi_2, \xi_4)|^2 \right)^{\frac{1}{2}} \times \\ & \| G(\cdot, \cdot) \|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \| H \Psi \| + (\tilde{b} + m_e) \| \Psi \|) \end{aligned}$$

with

$$\sum_{j=1}^4 \int d\xi_2 d\xi_4 \chi_{f_4}(\xi_4) |H_j^{(\nu_\mu)}(\xi_2, \xi_4)|^2 < \infty$$

Here  $\chi_{f_4}(\cdot)$  is the characteristic function of the support of  $f_4(\cdot)$ .

Similarly we have

$$(4.70) \quad \begin{aligned} & (\Phi, b_{4,+}^*(f_{4,T}) \Psi) - (\Phi, b_{4,+}^*(f_{4,T_0}) \Psi) \\ & = g \int_{T_0}^T \frac{d}{dt} (\Phi, b_{4,+}^*(f_{4,t}) \Psi) = ig \int_{T_0}^T (\Phi, e^{itH} [H_I, b_{4,+}^*(f_{4,t})] e^{-itH} \Psi) dt \end{aligned}$$

with

$$(4.71) \quad \begin{aligned} & \left[ (H_I^{(1)})^*, b_{4,+}^*(f_{4,t}) \right] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. \overline{W^{(\bar{\nu}_e)}}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1) \overline{U^{(\mu)}}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4) \right) \\ & \overline{F(\xi_2, \xi_4)} \overline{G(\xi_1, \xi_3)} f_{4,t}(\xi_4) b_+^*(\xi_2) b_-(\xi_3) b_+(\xi_1) \Psi. \end{aligned}$$

$$(4.72) \quad \begin{aligned} & \left[ (H_I^{(2)})^*, b_{4,+}^*(f_{4,t}) \right] \Psi = \int d\xi_1 d\xi_2 d\xi_3 d\xi_4 \left( \int dx^2 e^{ix^2 r^2} \right. \\ & \left. \overline{W^{(\bar{\nu}_e)}}(\xi_3) \gamma^\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1) \overline{W^{(\mu)}}(x^2, \xi_2) \gamma_\alpha (1 - \gamma_5) U^{(\nu_\mu)}(\xi_4) \right) \\ & \overline{F((\xi_2, \xi_4))} \overline{G(\xi_1, \xi_3)} f_{4,t} b_-(\xi_3) b_+(\xi_1) b_-(\xi_2) \Psi. \end{aligned}$$

and

$$(4.73) \quad \left[ H_I^{(1)}, b_{4,+}^*(f_{4,t}) \right] = \left[ H_I^{(2)}, b_{4,+}^*(f_{4,t}) \right] = 0$$

By (B.5) we get

$$(4.74) \quad \begin{aligned} & \left\| \left[ (H_I^{(1)})^*, b_{4,+}^*(f_{4,t}) \right] \Psi \right\| \leq \\ & \int dx^2 \left( \int d\xi_2 \left| \left\langle U^{(\mu)}(x^2, \xi_2), \gamma^0 \gamma^\alpha (1 - \gamma_5) \int d\xi_4 U^{(\nu_\mu)}(\xi_4) f_{4,t}(\xi_4) e^{ip_4^2 x^2} \overline{F(\xi_2, \xi_4)} \right\rangle \right|^2 \right)^{\frac{1}{2}} \\ & \left\| \int d\xi_1 d\xi_3 e^{-ip_3^2 x^2} \left\langle W^{(\bar{\nu}_e)}(\xi_3), \gamma^0 \gamma_\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1) \overline{G(\xi_1, \xi_3)} \right\rangle b_+(\xi_1) b_-(\xi_3) \Psi \right\|. \end{aligned}$$

where  $\langle \cdot, \cdot \rangle$  is the scalar product in  $\mathbb{C}^4$ .

and

$$(4.75) \quad \begin{aligned} & \left\| \left[ (H_I^{(2)})^*, b_{4,+}^*(f_{4,t}) \right] \Psi \right\| \leq \\ & \int dx^2 \left( \int d\xi_2 \left| \left\langle U^{(\mu)}(x^2, \xi_2), \gamma^0 \gamma^\alpha (1 - \gamma_5) \int d\xi_4 U^{(\nu_\mu)}(\xi_4) f_{4,t}(\xi_4) e^{ip_4^2 x^2} \overline{F(\xi_2, \xi_4)} \right\rangle \right|^2 \right)^{\frac{1}{2}} \\ & \left\| \int d\xi_1 d\xi_3 e^{-ip_3^2 x^2} \left\langle W^{(\bar{\nu}_e)}(\xi_3), \gamma^0 \gamma_\alpha (1 - \gamma_5) U^{(e)}(x^2, \xi_1) \overline{G(\xi_1, \xi_3)} \right\rangle b_+(\xi_1) b_-(\xi_3) \Psi \right\|. \end{aligned}$$

By adapting the proof of (4.57) and (4.67) to (4.74) and (4.75) one gets

$$(4.76) \quad \begin{aligned} & \left\| e^{itH} \left[ H_I, b_{4,+}^*(f_{4,t}) \right] e^{-itH} \Psi \right\| \leq \\ & CC_{f_4} \frac{1}{t^2} \left( \sum_{j=1}^4 \int d\xi_2 d\xi_4 \chi_{f_4}(\xi_4) |H_j^{(\nu_\mu)}(\xi_2, \xi_4)|^2 \right)^{\frac{1}{2}} \times \\ & \|G(\cdot, \cdot)\|_{L^2(\Gamma_1 \times \mathbb{R}^3)} \frac{1}{m_e} (\tilde{a} \|H\Psi\| + (\tilde{b} + m_e) \|\Psi\|) \end{aligned}$$

It follows from (4.63), (4.69), (4.70) and (4.76) that the strong limits of  $b_{4,+}^\sharp(f_4)$  exist when  $t$  goes to  $\pm\infty$ , for all  $f_4 \in L^2(\mathbb{R}^3)$  and for every  $g \leq g_0$ .

This concludes the proof of theorem 4.3.  $\square$

#### 4.2. Existence of a Fock space subrepresentation of the asymptotic CAR.

From now on we only consider the case where the time  $t$  goes to  $+\infty$ . The following proposition is an easy consequence of theorem 4.1.

##### Proposition 4.4.

Suppose Hypothesis 2.1-Hypothesis 4.2 and  $g \leq g_1$ . We have

i) Let  $f_1, g_1, f_2, g_2 \in L^2(\Gamma_1)$  and  $f_3, g_3, f_4, g_4 \in L^2(\mathbb{R}^3)$ . The following anticommutation relations hold in the sense of quadratic form.

$$\begin{aligned}
\{b_{1,+,\infty}(f_1), b_{1,+,\infty}^*(g_1)\} &= \langle f_1, g_1 \rangle_{L^2(\Gamma_1)} \mathbf{1} \\
\{b_{2,\epsilon,\infty}(f_2), b_{2,\epsilon',\infty}^*(g_2)\} &= \langle f_2, g_2 \rangle_{L^2(\Gamma_1)} \delta_{\epsilon\epsilon'} \mathbf{1} \\
\{b_{3,-,\infty}(f_3), b_{3,-,\infty}^*(g_3)\} &= \langle f_3, g_3 \rangle_{L^2(\mathbb{R}^3)} \mathbf{1} \\
\{b_{4,+,\infty}(f_4), b_{4,+,\infty}^*(g_4)\} &= \langle f_4, g_4 \rangle_{L^2(\mathbb{R}^3)} \mathbf{1} \\
\{b_{1,+,\infty}(f_1), b_{1,+,\infty}(g_1)\} &= \{b_{1,+,\infty}^*(f_1), b_{1,+,\infty}^*(g_1)\} = 0 \\
\{b_{1,+,\infty}(f_1), b_{2,\epsilon,\infty}^\sharp(f_2)\} &= \{b_{1,+,\infty}(f_1), b_{3,-,\infty}^\sharp(f_3)\} = 0 \\
\{b_{1,+,\infty}(f_1), b_{4,+,\infty}^\sharp(f_4)\} &= 0. \\
\{b_{2,\epsilon,\infty}(f_2), b_{2,\epsilon',\infty}(g_2)\} &= \{b_{2,\epsilon,\infty}^*(f_2), b_{2,\epsilon',\infty}^*(g_2)\} = 0 \\
\{b_{2,\epsilon,\infty}(f_2), b_{3,-,\infty}^\sharp(f_3)\} &= \{b_{2,\epsilon,\infty}(f_2), b_{4,+,\infty}^\sharp(f_4)\} = 0 \\
\{b_{3,-,\infty}(f_3), b_{3,-,\infty}(g_3)\} &= \{b_{3,-,\infty}^*(f_3), b_{3,-,\infty}^*(g_3)\} = 0 \\
\{b_{3,-,\infty}(f_3), b_{4,+,\infty}^\sharp(f_4)\} &= 0 \\
\{b_{4,+,\infty}(f_4), b_{4,+,\infty}(g_4)\} &= \{b_{4,+,\infty}^*(f_4), b_{4,+,\infty}^*(g_4)\} = 0.
\end{aligned}$$

Here  $\epsilon = \pm$ .

ii)

$$\begin{aligned}
e^{itH} b_{1,+,\infty}^\sharp(f_1) &= b_{1,+,\infty}^\sharp(e^{i\omega(\xi_1)t} f_1) e^{itH}. \\
e^{itH} b_{2,\pm,\infty}^\sharp(f_2) &= b_{2,\pm,\infty}^\sharp(e^{i\omega(\xi_2)t} f_2) e^{itH}. \\
e^{itH} b_{3,-,\infty}^\sharp(f_3) &= b_{3,-,\infty}^\sharp(e^{i\omega(\xi_3)t} f_3) e^{itH}. \\
e^{itH} b_{4,+,\infty}^\sharp(f_4) &= b_{4,+,\infty}^\sharp(e^{i\omega(\xi_4)t} f_4) e^{itH}.
\end{aligned}$$

and the following pullthrough formulae are satisfied:

$$\begin{aligned}
[H, b_{1,+,\infty}^*(f_1)] &= b_{1,+,\infty}^*(\omega(\xi_1) f_1), [H, b_{1,+,\infty}(f_1)] = -b_{1,+,\infty}(\omega(\xi_1) f_1) \\
[H, b_{2,\pm,\infty}^*(f_2)] &= b_{2,\pm,\infty}^*(\omega(\xi_2) f_2), [H, b_{2,\pm,\infty}(f_2)] = -b_{2,\pm,\infty}(\omega(\xi_2) f_2) \\
[H, b_{3,-,\infty}^*(f_3)] &= b_{3,-,\infty}^*(\omega(\xi_3) f_3), [H, b_{3,-,\infty}(f_3)] = -b_{3,-,\infty}(\omega(\xi_3) f_3) \\
[H, b_{4,+,\infty}^*(f_4)] &= b_{4,+,\infty}^*(\omega(\xi_4) f_4), [H, b_{4,+,\infty}(f_4)] = -b_{4,+,\infty}(\omega(\xi_4) f_4).
\end{aligned}$$

iii)

$$b_{1,+,\infty}(f_1)\Omega_g = b_{2,\pm,\infty}(f_2)\Omega_g = b_{3,-,\infty}(f_3)\Omega_g = b_{4,+,\infty}(f_4)\Omega_g = 0$$

Here  $\Omega_g$  is the ground state of  $H$ .

Our main result is the following theorem

**Theorem 4.5.** Suppose Hypothesis 2.1-Hypothesis 4.2 and  $g \leq g_1$ . Then we have

$$\sigma_{ac} = [E, \infty).$$

*Proof.* By (2.2) we have, for all sets of integers  $(p, q, \bar{q}, r, s)$  in  $\mathbb{N}^5$ ,

$$(4.77) \quad \mathfrak{F} = \bigoplus_{(p,q,\bar{q},r,s)} \mathfrak{F}^{(p,q,\bar{q},r,s)}.$$

with

$$(4.78) \quad \mathfrak{F}^{(p,q,\bar{q},r,s)} = (\otimes_a^p L^2(\Gamma_1)) \otimes (\otimes_a^q L^2(\Gamma_1)) \otimes (\otimes_a^{\bar{q}} L^2(\Gamma_1)) \otimes (\otimes_a^r L^2(\mathbb{R}^3)) \otimes (\otimes_a^s L^2(\mathbb{R}^3)).$$

Here  $p$  is the number of electrons,  $q$  (resp.  $\bar{q}$ ) is the number of muons (resp. antimuons),  $r$  is the number of antineutrinos  $\bar{\nu}_e$  and  $s$  is the number of neutrinos  $\nu_\mu$ .

Let  $\{e_i^1 | i = 1, 2, \dots\}, \{e_j^2 | j = 1, 2, \dots\}$  and  $\{f_k^2 | k = 1, 2, \dots\}$  be three orthonormal basis of  $L^2(\Gamma_1)$ . Let  $\{e_l^3 | l = 1, 2, \dots\}$  and  $\{e_m^4 | m = 1, 2, \dots\}$  be two orthonormal basis of  $L^2(\mathbb{R}^3)$ .

Consider the following vectors of  $\mathfrak{F}$

$$(4.79) \quad \prod_{1 \leq \alpha \leq p} b_{1,+}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{1,+}^*(e_{l_\alpha}^3) \prod_{1 \leq \alpha \leq s} b_{4,+}^*(e_{m_\alpha}^4) \Omega$$

The indices are assumed ordered,  $i_1 < \dots < i_p$ ,  $j_1 < \dots < j_q$ ,  $k_1 < \dots < k_{\bar{q}}$ ,  $l_1 < \dots < l_r$  and  $m_1 < \dots < m_s$ .

The set, for  $(p, q, \bar{q}, r, s)$  given in  $\mathbb{N}^5$ ,

$$\mathbb{D}^{(p,q,\bar{q},r,s)} = \{\Phi \in \mathfrak{F}^{(p,q,\bar{q},r,s)} \mid \Phi \text{ is a finite linear combination of basis vectors of the form (4.79)}\}$$

is a dense domain in  $\mathfrak{F}^{(p,q,\bar{q},r,s)}$ . The set of vectors of the form (4.79) is an orthonormal basis of  $\mathfrak{F}^{(p,q,\bar{q},r,s)}$  (see [39, Chapter 10]). Hence the vectors obtained in this way for  $p, q, \bar{q}, r, s, = 0, 1, 2, \dots$  form an orthonormal basis of  $\mathfrak{F}$  and the set

$$\mathbb{D} = \{\Psi \in \mathfrak{F} \mid \Psi \text{ is a finite linear combination of basis vectors of the form (4.79) for } p, q, \bar{q}, r, s = 0, 1, 2, \dots\}$$

is a dense domain in  $\mathfrak{F}$ .

On the other hand we now introduce the following vectors of  $\mathfrak{F}$

$$(4.80) \quad \prod_{1 \leq \alpha \leq p} b_{1,+\infty}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+\infty}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-\infty}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{3,-\infty}^*(e_{l_\alpha}^3) \prod_{1 \leq \alpha \leq s} b_{4,+\infty}^*(e_{m_\alpha}^4) \Omega_g$$

Let  $\mathfrak{F}_\infty^{(p,q,\bar{q},r,s)}$  denote the closed linear hull of vectors of the form (4.80). It follows from proposition 4.4 that the set of vectors of the form (4.80) is an orthonormal basis of  $\mathfrak{F}_\infty^{(p,q,\bar{q},r,s)}$ .

The set, for  $(p, q, \bar{q}, r, s)$  given in  $\mathbb{N}^5$ ,

$$\mathbb{D}_\infty^{(p,q,\bar{q},r,s)} = \{\Phi \in \mathfrak{F}_\infty \mid \Phi \text{ is a finite linear combination of basis vectors of the form (4.80)}\}.$$

is a dense domain in  $\mathfrak{F}_\infty^{(p,q,\bar{q},r,s)}$ .

The asymptotic outgoing Fock space denoted by  $\mathfrak{F}_\infty$  is then defined by

$$(4.81) \quad \mathfrak{F}_\infty = \bigoplus_{p,q,\bar{q},r,s} \mathfrak{F}_\infty^{(p,q,\bar{q},r,s)}.$$

The vectors of the form (4.80) obtained for  $p, q, \bar{q}, r, s = 0, 1, 2, \dots$  form an orthonormal basis of  $\mathfrak{F}$  and the set

$$\mathbb{D}_\infty = \{ \Phi \in \mathfrak{F}_\infty \mid \Phi \text{ is a finite linear combination of basis vectors} \\ \text{of the form (4.80) for } p, q, \bar{q}, r, s = 0, 1, 2, \dots \}$$

is a dense domain in  $\mathfrak{F}_\infty$ .

We now introduce the following linear operators, denoted by  $W_\infty^{(p,q,\bar{q},r,s)}$ , and defined on  $\mathbb{D}^{(p,q,\bar{q},r,s)}$  by

$$(4.82) \quad \begin{aligned} & W_\infty^{(p,q,\bar{q},r,s)} \prod_{1 \leq \alpha \leq p} b_{1,+}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{1,+}^*(e_{l_\alpha}^3) \\ & \quad \prod_{1 \leq \alpha \leq s} b_{4,+}^*(e_{m_\alpha}^4) \Omega \\ & = \prod_{1 \leq \alpha \leq p} b_{1,+, \infty}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+, \infty}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-, \infty}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{3,-, \infty}^*(e_{l_\alpha}^3) \\ & \quad \prod_{1 \leq \alpha \leq s} b_{4,+, \infty}^*(e_{m_\alpha}^4) \Omega_g. \end{aligned}$$

$W_\infty^{(p,q,\bar{q},r,s)}$  can be uniquely extended to linear operators from  $\mathbb{D}^{(p,q,\bar{q},r,s)}$  to  $\mathbb{D}^{(p,q,\bar{q},r,s)}$ . It then follows from proposition 4.4. that the operators  $W_\infty^{(p,q,\bar{q},r,s)}$  can be uniquely extended to unitary operators from  $\mathbb{D}^{(p,q,\bar{q},r,s)}$  to  $\mathbb{D}^{(p,q,\bar{q},r,s)}$

Let

$$(4.83) \quad W_\infty = \bigoplus_{p,q,\bar{q},r,s} W_\infty^{(p,q,\bar{q},r,s)}.$$

Hence  $W_\infty$  is a unitary operator from  $\mathfrak{F}$  to  $\mathfrak{F}_\infty$ .

The operators  $b_{1,+, \infty}(f_1), b_{1,+, \infty}(g_1), b_{2,+, \infty}(f_2), b_{2,+, \infty}(g_2), b_{2,-, \infty}(f_2), b_{2,-, \infty}(g_2), b_{3,-, \infty}(f_3), b_{3,-, \infty}(g_3), b_{4,+, \infty}(f_4)$  and  $b_{4,+, \infty}(g_4)$  defined on  $\mathfrak{F}_\infty$  generate a Fock representation of the ACR (see Proposition 4.4 i).

By proposition 4.4 ii) we have

$$(4.84) \quad \begin{aligned} & e^{itH} \prod_{1 \leq \alpha \leq p} b_{1,+, \infty}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+, \infty}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-, \infty}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{3,-, \infty}^*(e_{l_\alpha}^3) \\ & \quad \prod_{1 \leq \alpha \leq s} b_{4,+, \infty}^*(e_{m_\alpha}^4) \Omega_g \\ & = e^{iEt} \prod_{1 \leq \alpha \leq p} b_{1,+, \infty}^*(e^{i\omega(\xi_1)t} e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+, \infty}^*(e^{i\omega(\xi_2)t} e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-, \infty}^*(e^{i\omega(\xi_2)t} f_{k_\alpha}^2) \\ & \quad \prod_{1 \leq \alpha \leq r} b_{3,-, \infty}^*(e^{i\omega(\xi_3)t} e_{l_\alpha}^3) \prod_{1 \leq \alpha \leq s} b_{4,+, \infty}^*(e^{i\omega(\xi_4)t} e_{m_\alpha}^4) \Omega_g. \end{aligned}$$

Hence  $e^{iHt}$  leaves  $\mathfrak{F}_\infty$  invariant and  $H$  is both reduced by  $\mathfrak{F}_\infty$  and  $\mathfrak{F}_\infty^\perp$ . Thus

$$\mathfrak{F} \simeq \mathfrak{F}_\infty \oplus \mathfrak{F}_\infty^\perp$$

In view of (4.5), (4.48) and (4.84) we get

$$\begin{aligned}
& W_\infty e^{itH_0} \prod_{1 \leq \alpha \leq p} b_{1,+}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-}^*(f_{k_\alpha}^2) \prod_{1 \leq \alpha \leq r} b_{3,-}^*(e_{l_\alpha}^3) \\
& \prod_{1 \leq \alpha \leq s} b_{4,+}^*(e_{m_\alpha}^4) \Omega \\
(4.85) \quad & = e^{i(H-E)t} W_\infty \prod_{1 \leq \alpha \leq p} b_{1,+}^*(e_{i_\alpha}^1) \prod_{1 \leq \alpha \leq q} b_{2,+}^*(e_{j_\alpha}^2) \prod_{1 \leq \alpha \leq \bar{q}} b_{2,-}^*(f_{k_\alpha}^2) \\
& \prod_{1 \leq \alpha \leq r} b_{3,-}^*(e_{l_\alpha}^3) \prod_{1 \leq \alpha \leq s} b_{4,+}^*(e_{m_\alpha}^4) \Omega.
\end{aligned}$$

This yields

$$(4.86) \quad W_\infty e^{it(H_0+E)} = e^{iHt} W_\infty$$

Hence the reduction of  $H$  to  $\mathfrak{F}_\infty$  is unitarily equivalent to  $H_0 + E$ . Thus  $\sigma_{ac}(H) = [E, \infty)$ . This concludes the proof of theorem 4.5.  $\square$

#### ACKNOWLEDGEMENTS.

J.-C.G. acknowledges J.-M. Barbaroux, J. Faupin and G. Hachem for helpful discussions.

#### APPENDIX A. THE DIRAC QUANTIZED FIELDS FOR THE ELECTRONS AND THE MUONS.

The appendices are based on the section 2 and section 3 of [23]. See also [13], [14], [15] and [25].

$(s, n, p^1, p^3)$  are quantum variables of the electrons, the positrons, the muons and the antimuons in a uniform magnetic field. Here  $s = \pm 1$ ,  $n \geq 0$ ,  $p^1 \in \mathbb{R}$ ,  $p^3 \in \mathbb{R}$ .

Let  $\xi_1 = (s, n, p_e^1, p_e^3)$  be the quantum variables of a electron and let  $\xi_2 = (s, n, p_\mu^1, p_\mu^3)$  be the quantum variables of a muon and of an antimuon.

We set  $\Gamma_1 = \{-1, 1\} \times \mathbb{N} \times \mathbb{R}^2$  for the configuration space for the electrons, the muons and the antimuons.  $L^2(\Gamma_1)$  is the Hilbert space associated to each species of fermions.

$$(A.1) \quad L^2(\Gamma_1) = l^2(L^2(\mathbb{R}^2)) \oplus l^2(L^2(\mathbb{R}^2))$$

Let  $\mathfrak{F}_e$  and  $\mathfrak{F}_\mu$  denote the Fock spaces for the electrons and the muons respectively. Remark that  $\mathfrak{F}_\mu$  is also the Fock space for the antimuons.

We have

$$(A.2) \quad \mathfrak{F}_e = \mathfrak{F}_\mu = \bigoplus_{n=0}^{\infty} \bigotimes_a^n L^2(\Gamma_1).$$

$\bigotimes_a^n L^2(\Gamma_1)$  is the antisymmetric  $n$ -th tensor power of  $L^2(\Gamma_1)$ .

$\Omega_\alpha = (1, 0, 0, 0, \dots)$  is the vacuum state in  $\mathfrak{F}_\alpha$  for  $\alpha = e, \mu$ .

We shall use the notations

$$\begin{aligned}
(A.3) \quad & \int_{\Gamma_1} d\xi_1 = \sum_{n \geq 0} \int_{\mathbb{R}^2} dp_e^1 dp_e^3 \\
& \int_{\Gamma_1} d\xi_2 = \sum_{s=\pm 1} \sum_{n \geq 0} \int_{\mathbb{R}^2} dp_\mu^1 dp_\mu^3.
\end{aligned}$$

$b_+(\xi_1)$  (resp.  $b_+^*(\xi_1)$ ) is the annihilation (resp. creation) operator for the electron.

Let  $\epsilon = \pm$ .

$b_\epsilon(\xi_2)$  (resp.  $b_\epsilon^*(\xi_2)$ ) are the annihilation (resp. creation) operators for the muon when  $\epsilon = +$  and for the antimuon when  $\epsilon = -$ .

The operators  $b_+(\xi_1)$ ,  $b_+^*(\xi_1)$ ,  $b_\epsilon(\xi_2)$  and  $b_\epsilon^*(\xi_2)$  fulfil the usual anticommutation relations (CAR) (see [40] and [23]). Therefore the following anticommutation relations hold

$$(A.4) \quad \begin{aligned} \{b_+(\xi_1), b_+^*(\xi_1')\} &= \delta(\xi_1 - \xi_1'), \\ \{b_\epsilon(\xi_2), b_{\epsilon'}^*(\xi_2')\} &= \delta_{\epsilon\epsilon'} \delta(\xi_2 - \xi_2'), \\ \{b_+^\sharp(\xi_1), b_{\epsilon'}^\sharp(\xi_2)\} &= 0. \end{aligned}$$

where  $\{b, b'\} = bb' + b'b$  and  $b^\sharp = b$  or  $b^*$ .

In addition, following the convention described in [40, Section 4.1] and [40, Section 4.2], we assume that the fermionic creation and annihilation operators of different species of particles anticommute (see [8] arXiv for explicit definitions). In our case this property will be verified by the creation and annihilation operators for the electrons, the muons, the antimuons, the antineutrinos associated to the electron and the neutrinos associated to the muons.

Recall that for  $\varphi \in L^2(\Gamma_1)$ , the operators

$$(A.5) \quad \begin{aligned} b_{1,+}(\varphi) &= \int_{\Gamma_1} b(\xi_1) \overline{\varphi(\xi_1)} d\xi_1, \\ b_{1,+}^*(\varphi) &= \int_{\Gamma_1} b^*(\xi_1) \varphi(\xi_1) d\xi_1, \\ b_{2,\epsilon}(\varphi) &= \int_{\Gamma_1} b_\epsilon(\xi_2) \overline{\varphi(\xi_2)} d\xi_2, \\ b_{2,\epsilon}^*(\varphi) &= \int_{\Gamma_1} b_\epsilon^*(\xi_2) \varphi(\xi_2) d\xi_2. \end{aligned}$$

are bounded operators on  $\mathfrak{F}_{(e)}$  and  $\mathfrak{F}_{(\mu)}$  respectively satisfying

$$(A.6) \quad \begin{aligned} \|b_{1,+}^\sharp(\varphi)\| &= \|\varphi\|_{L^2} \\ \|b_{2,\epsilon}^\sharp(\varphi)\| &= \|\varphi\|_{L^2} \end{aligned}$$

Set  $\alpha = e, \mu$ . The Dirac quantized fields for the electron and the muon, denoted by  $\Psi_{(\alpha)}(\mathbf{x})$ , are given by

$$(A.7) \quad \begin{aligned} \Psi_{(\alpha)}(\mathbf{x}) &= \frac{1}{2\pi} \int d\xi_1 \left( e^{i(p_\alpha^1 x^1 + p_\alpha^3 x^3)} U^{(\alpha)}(x^2, \xi_\alpha) b_+(\xi_\alpha) \right. \\ &\quad \left. + e^{-i(p_\alpha^1 x^1 + p_\alpha^3 x^3)} W^{(\alpha)}(x^2, \xi_\alpha) b_-^*(\xi_\alpha) \right). \end{aligned}$$

where  $\xi_e = \xi_1$  and  $\xi_\mu = \xi_2$ . See [17].

For  $\xi_\alpha = (s, n, p_\alpha^1, p_\alpha^3)$  we have  $U^{(\alpha)}(x^2, \xi_\alpha) = U_s^{(\alpha)}(x^2, n, p_\alpha^1, p_\alpha^3)$ .

For  $s = 1$  and  $n \geq 1$   $U_1^{(\alpha)}(x^2, n, p^1, p^3)$  is given by

$$(A.8) \quad U_{+1}^{(\alpha)}(x^2, n, p^1, p^3) = \left( \frac{E_n^{(\alpha)}(p^3) + m_\alpha}{2E_n^{(\alpha)}(p^3)} \right)^{\frac{1}{2}} \begin{pmatrix} I_{n-1}(\xi) \\ 0 \\ \frac{p^3}{E_n^{(\alpha)}(p^3) + m_\alpha} I_{n-1}(\xi) \\ -\frac{\sqrt{2neB}}{E_n^{(\alpha)}(p^3) + m_\alpha} I_n(\xi) \end{pmatrix}$$

where

$$(A.9) \quad \begin{aligned} \xi &= \sqrt{eB} \left( x^2 - \frac{p^1}{eB} \right) \\ I_n(\xi) &= \left( \frac{\sqrt{eB}}{n! 2^n \sqrt{\pi}} \right)^{\frac{1}{2}} \exp(-\xi^2/2) H_n(\xi). \end{aligned}$$

Here  $H_n(\xi)$  is the Hermite polynomial of order  $n$  and we define

$$(A.10) \quad I_{-1}(\xi) = 0$$

For  $n = 0$  and  $s = 1$  we set

$$U_{+1}^{(\alpha)}(x^2, 0, p^1, p^3) = 0$$

For  $s = -1$  and  $n \geq 0$   $U_{-1}^{(\alpha)}(x^2, n, p^1, p^3)$  is given by

$$(A.11) \quad U_{-1}^{(\alpha)}(x^2, n, p^1, p^3) = \left( \frac{E_n^{(\alpha)}(p^3) + m_\alpha}{2E_n^{(\alpha)}(p^3)} \right)^{\frac{1}{2}} \begin{pmatrix} 0 \\ I_n(\xi) \\ -\frac{\sqrt{2neB}}{E_n^{(\alpha)}(p^3) + m_\alpha} I_{n-1}(\xi) \\ -\frac{p^3}{E_n^{(\alpha)}(p^3) + m_\alpha} I_n(\xi) \end{pmatrix}$$

Through out this work  $e$  will be the positive unit of charge taken to be equal to the proton charge.

$E_n^{(\alpha)}(p^3)$ ,  $n \geq 0$ , is given by

$$(A.12) \quad E_n^{(\alpha)}(p^3) = \sqrt{m_\alpha^2 + (p^3)^2 + 2neB}$$

Note that

$$(A.13) \quad \int dx^2 U_s^{(\alpha)}(x^2, n, p^1, p^3)^\dagger U_{s'}^{(\alpha)}(x^2, n, p^1, p^3) = \delta_{ss'}$$

where  $\dagger$  is the adjoint in  $\mathbb{C}^4$ .

In order to study the spectral theory of our Hamiltonian it is not necessary to know  $W^{(e)}(x^2, \xi_1)$  in (A.6). We have to know  $W^{(\mu)}(x^2, \xi_2)$  explicitly.

For  $\xi_2 = (s, n, p_\mu^1, p_\mu^3)$  with  $n > 0$  we have

$$(A.14) \quad \begin{aligned} W^{(\mu)}(x^2, \xi_2) &= V_{-1}^{(\mu)}(x^2, n, -p_\mu^1, -p_\mu^3) \quad \text{for } \xi_2 = (1, n, p_\mu^1, p_\mu^3), n \geq 0. \\ W^{(\mu)}(x^2, \xi_2) &= V_{+1}^{(\mu)}(x^2, n, -p_\mu^1, -p_\mu^3) \quad \text{for } \xi_2 = (-1, n, p_\mu^1, p_\mu^3), n \geq 1. \\ W^{(\mu)}(x^2, \xi_2) &= 0 \quad \text{for } \xi_1 = (-1, 0, p_\mu^1, p_\mu^3). \end{aligned}$$

For  $s = 1$  and  $n \geq 1$   $V_{+1}^{(\mu)}(x^2, n, p^1, p^3)$  is given by

$$(A.15) \quad V_{+1}^{(\mu)}(x^2, n, p^1, p^3) = \left( \frac{E_n^{(\mu)}(p^3) + m_\mu}{2E_n^{(\mu)}(p^3)} \right)^{\frac{1}{2}} \begin{pmatrix} -\frac{p^3}{E_n^{(\mu)}(p^3) + m_\mu} I_{n-1}(\xi) \\ \frac{\sqrt{2neB}}{E_n^{(\mu)}(p^3) + m_\mu} I_n(\xi) \\ I_{n-1}(\xi) \\ 0 \end{pmatrix}$$

and for  $n = 0$  we set

$$V_{+1}^{(\mu)}(x^2, 0, p^1, p^3) = 0$$

For  $s = -1$  and  $n \geq 0$   $V_{-1}^{(\mu)}(x^2, n, p^1, p^3)$  is given by



$$(A.16) \quad V_{-1}^{(\mu)}(x^2, n, p^1, p^3) = \left( \frac{E_n^{(\mu)}(p^3) + m_\mu}{2E_n^{(\mu)}(p^3)} \right)^{\frac{1}{2}} \begin{pmatrix} \frac{\sqrt{2neB}}{E_n^{(\mu)}(p^3) + m_\mu} I_{n-1}(\xi) \\ \frac{p^3}{E_n^{(\mu)}(p^3) + m_\mu} I_n(\xi) \\ 0 \\ I_n(\xi) \end{pmatrix}$$

Note that

$$(A.17) \quad \int dx^2 V_s^{(\mu)}(x^2, n, p^1, p^3)^\dagger V_{s'}^{(\mu)}(x^2, n, p^1, p^3) = \delta_{ss'}$$

where  $\dagger$  is the adjoint in  $\mathbb{C}^4$ .

## APPENDIX B. THE DIRAC QUANTIZED FIELDS FOR $\nu_\mu$ AND $\bar{\nu}_e$ .

We suppose that neutrinos and antineutrinos are massless as in the Standard Model.

The quantum variables of the neutrinos and antineutrinos are the momenta and the helicities.

Let  $\mathbf{P} = (P^1, P^2, P^3)$  be the generators of space-translations.  $H^3$  is the helicity operator  $\frac{1}{2} \frac{\mathbf{P} \cdot \boldsymbol{\Sigma}}{|\mathbf{P}|}$  where  $|\mathbf{P}| = \sqrt{\sum_{i=1}^3 (P^i)^2}$  and  $\boldsymbol{\Sigma} = (\Sigma^1, \Sigma^2, \Sigma^3)$  with for  $j = 1, 2, 3$

$$(B.1) \quad \Sigma^j = \begin{pmatrix} \sigma_j & 0 \\ 0 & \sigma_j \end{pmatrix}$$

The helicity of the neutrino associated to the muon is  $-\frac{1}{2}$ .  $\nu_\mu$  is left-handed. The helicity of the antineutrino associated to the electron is  $\frac{1}{2}$ .  $\bar{\nu}_e$  is right-handed.

Let  $\xi_3 = (\mathbf{p}, \frac{1}{2})$  be the quantum variables of the antineutrino  $\bar{\nu}_e$  where  $\mathbf{p} \in \mathbb{R}^3$  is the momentum and  $\frac{1}{2}$  is the helicity. Let  $\xi_4 = (\mathbf{p}, -\frac{1}{2})$  be the quantum variables of the neutrino  $\nu_\mu$  where  $\mathbf{p} \in \mathbb{R}^3$  is the momentum and  $-\frac{1}{2}$  is the helicity.

$L^2(\mathbb{R}^3)$  is the Hilbert space of the states of the neutrinos  $\nu_\mu$  and of the antineutrinos  $\bar{\nu}_e$ . Let  $\mathfrak{F}_{(\nu_\mu)}$  and  $\mathfrak{F}_{(\bar{\nu}_e)}$  denote the Fock spaces for the neutrinos and the antineutrinos respectively.

We have

$$(B.2) \quad \mathfrak{F}_{\nu_\mu} = \mathfrak{F}_{\bar{\nu}_e} = \bigoplus_{n=0}^{\infty} \bigotimes_a^n L^2(\mathbb{R}^3)$$

$\Omega_\beta = (1, 0, 0, 0, \dots)$  is the vacuum state in  $\mathfrak{F}_\beta$  for  $\beta = \bar{\nu}_e, \nu_\mu$ .

In the sequel we shall use the notations

$$(B.3) \quad \int_{\mathbb{R}^3} d\xi_3 = \int_{\mathbb{R}^3} d^3\mathbf{p}$$

$$\int_{\mathbb{R}^3} d\xi_4 = \int_{\mathbb{R}^3} d^3\mathbf{p}$$

$b_-(\xi_3)$  and  $b_*(\xi_3)$  are the annihilation and creation operators for the antineutrino associated to the electron respectively.  $b_+(\xi_4)$  and  $b_*(\xi_4)$  are the annihilation and creation operators for the neutrino associated to the muon respectively. The operators  $b_-^\sharp(\xi_3)$  and  $b_+^\sharp(\xi_4)$ , fulfil the usual anticommutation relations (CAR) and they anticommute with  $b_+^\sharp(\xi_1)$  and  $b_e^\sharp(\xi_2)$  according to the convention described in [40, Section 4.1]. See [8] arXiv for explicit definitions.

Therefore the following anticommutation relations hold

$$\begin{aligned}
\{b_-(\xi_3), b_-^*(\xi'_3)\} &= \delta(\xi_3 - \xi'_3), \\
\{b_+(\xi_4), b_+(\xi'_4)\} &= \delta(\xi_4 - \xi'_4), \\
\{b_-^\sharp(\xi_3), b_+^\sharp(\xi_4)\} &= 0 \\
\{b_-^\sharp(\xi_3), b_\epsilon^\sharp(\xi_2)\} &= \{b_-^\sharp(\xi_3), b_+^\sharp(\xi_1)\} = 0. \\
\{b_+^\sharp(\xi_4), b_\epsilon^\sharp(\xi_2)\} &= \{b_+^\sharp(\xi_4), b_+^\sharp(\xi_1)\} = 0.
\end{aligned}
\tag{B.4}$$

Recall that, for  $\varphi \in L^2(\mathbb{R}^3)$ , the operators

$$\begin{aligned}
b_{4,+}(\varphi) &= \int_{\mathbb{R}^3} b_+(\xi_4) \overline{\varphi(\xi_4)} d\xi_4. \\
b_{3,-}(\varphi) &= \int_{\mathbb{R}^3} b_-(\xi_3) \overline{\varphi(\xi_3)} d\xi_3. \\
b_{4,+}^*(\varphi) &= \int_{\mathbb{R}^3} b_+^*(\xi_4) \varphi(\xi_4) d\xi_4. \\
b_{3,-}^*(\varphi) &= \int_{\mathbb{R}^3} b_-^*(\xi_3) \varphi(\xi_3) d\xi_3.
\end{aligned}
\tag{B.5}$$

are bounded operators on  $\mathfrak{F}_{(\nu_\mu)}$  and  $\mathfrak{F}_{(\bar{\nu}_e)}$  respectively satisfying

$$\|b_4^\sharp(\varphi)\| = \|b_3^\sharp(\varphi)\| = \|\varphi\|_{L^2}.
\tag{B.6}$$

The Dirac quantized fields for the neutrinos and antineutrinos associated to the electron and the muon respectively are denoted by  $\Psi_{(\nu_e)}(\mathbf{x})$  and  $\Psi_{(\nu_\mu)}(\mathbf{x})$ .

We have

$$\begin{aligned}
\Psi_{(\nu_e)}(\mathbf{x}) &= \left(\frac{1}{2\pi}\right)^{\frac{3}{2}} \left( \int d\tilde{\xi}_3 e^{i(\mathbf{p}\cdot\mathbf{x})} U^{(\nu_e)}(\tilde{\xi}_3) b_+(\tilde{\xi}_3) \right. \\
&\quad \left. + \int d\xi_3 e^{-i(\mathbf{p}\cdot\mathbf{x})} W^{(\bar{\nu}_e)}(\xi_3) b_-^*(\xi_3) \right).
\end{aligned}
\tag{B.7}$$

and

$$\begin{aligned}
\Psi_{(\nu_\mu)}(\mathbf{x}) &= \left(\frac{1}{2\pi}\right)^{\frac{3}{2}} \left( \int d\xi_4 e^{i(\mathbf{p}\cdot\mathbf{x})} U^{(\nu_\mu)}(\xi_4) b_+(\xi_4) \right. \\
&\quad \left. + \int d\tilde{\xi}_4 e^{-i(\mathbf{p}\cdot\mathbf{x})} W^{(\bar{\nu}_\mu)}(\tilde{\xi}_4) b_-^*(\tilde{\xi}_4) \right).
\end{aligned}
\tag{B.8}$$

where  $\tilde{\xi}_3 = (\mathbf{p}, -\frac{1}{2})$  and  $\tilde{\xi}_4 = (\mathbf{p}, \frac{1}{2})$  with, for  $\beta = 3, 4$ ,

$$\begin{aligned}
\int_{\mathbb{R}^3} d\xi_\beta &= \int_{\mathbb{R}^3} d^3p \\
\int_{\mathbb{R}^3} d\tilde{\xi}_\beta &= \int_{\mathbb{R}^3} d^3p
\end{aligned}
\tag{B.9}$$

See [17].

For the purpose of this paper one only needs to know  $W^{(\bar{\nu}_e)}(\xi_3)$  and  $U^{(\nu_\mu)}(\xi_4)$  explicitly.  $U^{(\nu_e)}(\tilde{\xi}_3)$  and  $W^{(\bar{\nu}_\mu)}(\tilde{\xi}_4)$  are given in [23].

By [23, (3.6), (3.7), (3.24), (3.32)] and [39] we have

$$U^{(\nu_\mu)}(\xi_4) = U^{(\nu_\mu)}(\mathbf{p}, -\frac{1}{2}) = \frac{1}{\sqrt{2}} \begin{pmatrix} h_-(\mathbf{p}) \\ -h_-(\mathbf{p}) \end{pmatrix}
\tag{B.10}$$

with

$$h_-(\mathbf{p}) = \frac{1}{\sqrt{2|\mathbf{p}|(|\mathbf{p}| - p^3)}} \begin{pmatrix} p^3 - |\mathbf{p}| \\ p^1 + ip^2 \end{pmatrix}
\tag{B.11}$$

and for  $|\mathbf{p}| = p^3$  we set

$$h_-(\mathbf{p}) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Moreover we have

$$(B.12) \quad W^{(\nu_e)}(\xi_3) = V^{(\nu_e)}\left(-\mathbf{p}, \frac{1}{2}\right) = \frac{1}{\sqrt{2}} \begin{pmatrix} -h_+(-\mathbf{p}) \\ h_+(-\mathbf{p}) \end{pmatrix}$$

with

$$(B.13) \quad h_+(-\mathbf{p}) = \frac{1}{\sqrt{2|\mathbf{p}|(|\mathbf{p}| + p^3)}} \begin{pmatrix} -p^1 + ip^2 \\ |\mathbf{p}| + p^3 \end{pmatrix}$$

and for  $|\mathbf{p}| = -p^3$  we set

$$h_+(-\mathbf{p}) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Note that

$$(B.14) \quad \|U^{(\nu_\mu)}(\xi_4)\|_{\mathbb{C}^4} = \|W^{(\bar{\nu}_e)}(\xi_3)\|_{\mathbb{C}^4} = 1$$

#### REFERENCES

- [1] B.L. Alvarez and J. Faupin. Scattering theory for mathematical models of the weak interaction. ArXiv 1809.02456. 2018. [arXiv:1809.02456](#)
- [2] B.L. Alvarez, J. Faupin and J.-C. Guillot Hamiltonians models of interacting fermion fields in Quantum Field Theory.. ArXiv 1810.10924. 2018. [arXiv:1810.10924](#)
- [3] Z. Ammari. Scattering theory for a class of fermionic Pauli-Fierz model. *J. Funct. Anal.*, 208(2) (2004), 302–359.
- [4] A. Arai. Essential spectrum of a self-adjoint operator on a abstract hilbert of Fock type and applications to quantum field Halmitonians. *J. Math. Anal. Appl.*, 246 ,(2000), 189-216.
- [5] W.H. Aschbacher, J.-M. Barbaroux, J. Faupin and J.-C. Guillot. Spectral theory for a mathematical model of the weak interaction: The decay of the intermediate vector bosons  $W^\pm$ . II. *Ann. Henri Poincaré*, 12 ,(2011), 1539–1570. [arXiv:1105.2247](#)
- [6] M. Ballesteros, D.-A. Deckert and F. Hänle. Relation between the resonant and the scattering matrix in the massless spin-boson model. ArXiv 1801.04843.2018. [arXiv:1801.04843](#)
- [7] J.-M. Barbaroux, M. Dimassi and J.-C. Guillot. Quantum electrodynamics of relativistic bound states with cutoffs. *J. Hyperbolic Differ. Equ.*, 1(2), (2004), 271–314.
- [8] J.-M. Barbaroux and J.-C. Guillot. Spectral theory for a mathematical model of the weak interaction: The decay of the intermediate vector bosons  $W^\pm$ . I. *Advances in Mathematical Physics* ID 978903 (2009), [arXiv:0904.3171](#)
- [9] J.-M. Barbaroux, J. Faupin, J.-C. Guillot *Spectral properties for Hamiltonians of weak interactions*, Operator Theory: Advances and Applications, 254, (2016), 11–36.
- [10] J.-M. Barbaroux, J. Faupin, J.-C. Guillot *Spectral Theory near Thresholds for Weak Interactions with Massive Particles*, Journal of Spectral Theory, 6, (2016), 505–555.
- [11] J.-M. Barbaroux, J. Faupin and J.-C. Guillot. Local decay for weak interactions with massless particles. ArXiv 1611.07814. To be published in J. Spectr. Theory 2018. [arXiv:1611.07814](#)
- [12] J. Beranger *et al* (Particle Data Group). *Phys. Rev.D* , 86 , O10001, 2012.
- [13] K. Bhattacharya. Solution of the Dirac equation in presence of an uniform magnetic field. [arXiv:0705.4275](#)
- [14] K. Bhattacharya and P.B. Pal. Inverse beta decay of arbitrarily polarized neutrons in a magnetic field. *Pramana.*, 62, (2004) , 1041. [arXiv:hep-ph/0209053](#)
- [15] K. Bhattacharya and P.B. Pal. Neutrinos and magnetic fields: a short review. *Proc. Ind. Natl. Sci. Acad.*, 70 ,(2004), 145. [arXiv:hep-ph/0212118](#)
- [16] J.-F. Bony, J. Faupin and I.M. Sigal Maximal velocity of photons in non-relativistic QED *Adv. Math.*, 231 ,(2012), 3054–3078.
- [17] J. Dereziński and C. Gérard. Asymptotic completeness in quantum field theory. Massive Pauli-Fierz Hamiltonians. *Rev. Math. Phys.*, 11(4) ,(1999), 383–450.
- [18] J. Dereziński and C. Gérard. *Mathematics of quantization and quantum fields*. Cambridge University Press. 2013.
- [19] J. Fröhlich, M. Griesemer and B. Schlein. Rayleigh scattering at atoms with dynamical nuclei. *Comm. Math. Phys.*, 271 ,(2007), 387–430.

- [20] J. Glimm, A. Jaffe. *Quantum Field Theory and Statistical Mechanics*. Birkhäuser, Boston inc., Boston MA , 1985, Expositions, Reprint of articles published in 1969–1977.
- [21] W. Greiner and B. Müller. *Gauge Theory of weak interactions*. Springer, Berlin, 2000.
- [22] J.C. Guillot. Spectral theory of a mathematical model in quantum field theory for any spin, *Spectral theory and partial differential equations*, 13-37, Contemp. Math., 640, Amer. Math. Soc., Providence, RI, 2015.
- [23] J.C. Guillot Weak Interactions in a Background of a Uniform Magnetic Field. A Mathematical Model for the Inverse Beta Decay.I. *Open Access Library Journal*, 4, e4142.2017. hal-01585239 and mp-arc 18-35.
- [24] C. Guinti and A. Studenikin. Neutrino electromagnetic interactions: a window to new physics. *Rev.Mod.Phys.*, 87 , 2015 , 531 .
- [25] G. Hachem Effet Zeeman pour un électron de Dirac. *Ann. Inst. Henri Poincaré*, 58 , 1993 , 105–123 .
- [26] F. Hiroshima. Ground states and spectrum of quantum electrodynamics of nonrelativistic particles. *Trans. Amer. Math.Soc.*, 353 , (2001) , 4497–4528.
- [27] R. Høegh-Krohn. Asymptotic fields in some models of quantum field theory.I. *J.Math. Phys.*, 9 , (1968) , 2075-2080.
- [28] R. Høegh-Krohn. Asymptotic fields in some models of quantum field theory.II. *J. Math. Phys.*, 10 , (1969) , 639-643.
- [29] R. Høegh-Krohn. Asymptotic fields in some models of quantum field theory.III. *J. Math. Phys.*, 11 , (1969) , 185-188.
- [30] R. Høegh-Krohn. Boson fields under a general class of cut-off interactions. *Comm. Math. Phys.*, 12 , (1969) , 216-225.
- [31] R. Høegh-Krohn. On the scattering operator for quantum fields. *Comm. Math. Phys.*, 18 , (1970) , 109-126.
- [32] M. Hübner and H. Spohn Radiative decay: nonperturbative approaches. *Rev. Math. Phys.*, 7 , (1995) , 363-387.  
Springer-Verlag, New York Inc. (1966).
- [33] Y. Kato and N. Mugibayashi. Regular perturbation and asymptotic limits of operators in quantum field theory. *Progr. Theoret. Phys.* , 30 ,(1963), 103-133.
- [34] N. Mugibayashi and Y. Kato Regular perturbation and asymptotic limits of operators in fixed-source theory. *Progr. Theoret. Phys.* , 31 ,(1964), 300-310.
- [35] Y. Kato and N. Mugibayashi. Asymptotic fields in model field theories.I. *Progr. Theoret. Phys.* , 45 ,(1971), 628-639.
- [36] T. Takaesu. On the spectral analysis of quantum electrodynamics with spatial cutoffs. I. *J. Math. Phys.*, 50 , (2009), 06230.
- [37] T. Takaesu. On generalized spin-boson models with singular perturbations. *Hokkaido Mathematical Journal*, 39 , (2010) , 1-33.
- [38] T. Takaesu. Essential Spectrum of a Fermionic Quantum Field Model. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 17 , No4, (2014), 1450024.
- [39] B. Thaller. *The Dirac Equation*. Texts and Monographs in Physics, Springer Verlag, Berlin, 1992.
- [40] S. Weinberg. *The quantum theory of fields. Vol. I*. Cambridge University Press, Cambridge, 2005. Foundations.
- [41] S. Weinberg. *The quantum theory of fields. Vol. II*. Cambridge University Press, Cambridge, 2005. Modern applications.

(J.-C. Guillot) CENTRE DE MATHÉMATIQUES APPLIQUÉES, UMR 7641, ÉCOLE POLYTECHNIQUE  
- C.N.R.S, 91128 PALAISEAU CEDEX, FRANCE  
Email address: Jean-Claude.Guillot@polytechnique.edu