Ill-defined Block-spin Transformations at arbitrarily high Temperatures

Aernout C. D. van Enter
Institute for Theoretical Physics
Rijksuniversiteit Groningen
P.O. Box 800
9747 AG Groningen
THE NETHERLANDS
AENTER@TH.RUG.NL

April 26, 1997

Abstract

Examples are presented of block-spin transformations which map the Gibbs measures of the Ising model in two or more dimensions at temperature intervals extending to arbitrarily high temperatures onto non-Gibbsian measures. In this way we provide the first example of this kind of pathology for very high temperatures, and as a corollary also the first example of such a pathology happening at a critical point.

1 Introduction

In [27, 28, 26] it was shown how various renormalization-group (RG) maps acting on Gibbs measures produce non-Gibbsian measures. In physicists' language, this means that a "renormalized Hamiltonian" can not be defined. The examples presented there were valid at low temperatures or close to a first-order phase transition. The underlying mechanism — pointed out first by Griffiths, Pearce and Israel [7, 8, 10] — is the fact that for the constraints imposed by particular choices of block-spin configurations, the resulting system exhibits a first-order phase transition. For this to happen, it was expected that the original system should be itself at or in the vicinity of a phase transition. Block-average and majority-rule transformations, however, provided counter-examples to this belief, in that they lead to non-Gibbsianness for arbitrarily large values of the magnetic field (at low temperatures) [28, 26].

On the other hand, there exist complementary results, either proving or providing heuristic but plausible arguments, that renormalized Gibbsian measures will be Gibbsian again at high fields or high temperatures [7, 8, 10, 2, 11], in the uniqueness region after a sufficiently often iterated decimation (possibly combined with another transformation) [18, 19], and in intervals around and including critical points for a rather general class of transformations [12, 1, 29, 4, 13].

For further investigations about the non-Gibbsianness of various measures of interest in statistical mechanics, see [28] and references quoted there, and also for more recent results [16, 6, 24, 23, 17, 30, 15].

Here we present the first examples where such a Griffiths-Pearce-Israel type pathology occurs in the high temperature region within the domain of (strong) Complete Analyticity [5]. Our example also includes a critical temperature pathology. Thus we show that there are no pathology-free domains for general block-spin maps. Note that the Potts decimation example of [26] provides a weaker statement; although a pathology occurs above the transition temperature, strong Complete Analyticity does not apply, even though, at least in the two-dimensional case, one expects that the weak Complete Analyticity property (for sufficiently regular volumes) holds [20, 21, 22, 25].

2 The example

We consider the standard nearest neighbor Ising model with (formal) Hamiltonian

$$H = \Sigma_{<i,j>} \; -\sigma(i)\sigma(j) \tag{2.1}$$

at inverse temperature β .

We consider the following transformation :

Divide the lattice Z^d into blocks B_j^L of linear size L and define block-spins $\sigma'(j)$ by

$$\sigma'(j) = \Sigma_{i \in B_j^L} \; \sigma(i) \tag{2.2}$$

except in the case where all spins in the block are minus,in which case

$$\sigma'(j) = +L^d \tag{2.3}$$

This induces a map T_L on the measures on the space of spin-configurations. For more details we refer to [28].

Now we have the following theorem:

Theorem: Let μ_β be the Gibbs measure of the nearest neighbor Ising model in dimension $d \geq 2$, at inverse temperature $\beta > 0$. Then there exists a constant L_0 such that for all $L > L_0$, $\mu'_L = \mu \circ T_L$ is non-Gibbsian.

Proof: The proof follows the scheme of [28], section 4.2. The special block-spin configuration which will be the point of (essential) discontinuity for a suitably chosen conditional probability will be the choice $\sigma'(j)=L^d$ in each block. This means that within each block all spins are unanimous, either in the plus or in the minus-direction. Thus imposing the special block-spin configuration as a constraint means that the sum of all spins in each block can take only two values (either unanimously positive or unanimously negative). Hence the thus constrained system is again a nearest neighbour Ising model, but now at an inverse temperature $\beta' = L^{d-1} \times \beta$. For L large enough, β' becomes so large that the constrained system is in the phase transition (low temperature) regime (step 1 of the proof). The block-spin boundary conditions selecting the plus and minus measures are $\sigma'(j) = +$ or $-(L^d - 2)$ in each boundary block. This means that in every block there is only one spin opposing all the others. When $L > 2$, a standard Peierls argument proves the phase transition (step 2 of the proof). As the expectation of $\sigma'(0)$ (unfixed according to step 3 of the proof) is different in the plus and minus phases, we see that the scheme of proof of [28], section 4.2, can be straightforwardly applied.

3 Comments and Conclusions

1.As a corollary of the theorem we obtain the first example where a Griffiths-Pearce-Israel pathology occurs *at* a critical point.

2.The block-spin map defined above is admittedly artificial from a physical point of view. However, the heuristic considerations of [1, 4], which were designed for the

block-average transformation formally equally well apply to this map, but lead to an incorrect conclusion. This illustrates the danger of relying on formal as opposed to fully rigorous arguments.

3.If we consider the sequence $\mu \circ T_L$, as a corollary to the (local) central limit theorem [9, 3, 2], it is easy to see that, with a proper rescaling of the block spins, it converges weakly to an independent Gaussian zero-mean measure. The measures in the sequence are almost all (that is, all except for possibly a finite number) non-Gibbsian.

4.Applying a decimation transformation often enough will make the measure Gibbsian again [19]. This seems to mean that there are maps which reinforce, as well as maps which weaken, the Gibbsianness of measures in the uniqueness region.What the behaviour of a "typical" map - whatever that might be- is, remains open. In this sense, how robust the phenomenon of renormalisation-group pathologies is, remains an issue of debate.Our result shows that the pathologies are more widespread than was known before, while the results of [18, 19, 1] go in the opposite direction, indicating how one might make some of these pathologies disappear.

Acknowledgements: I had very useful conversations with R. L. Dobrushin, who first asked me about the possible occurrence of pathologies at very high temperatures during a workshop at the Erwin Schrödinger Institute in Vienna, G. Grimmett, and E. Olivieri, who explained to me how imposing constraints can raise, as well as lower, critical temperatures. The proof given above is based on ideas which I developed in a long and very rewarding collaboration with Roberto Fernández and Alan Sokal. This research has been supported by EU contract CHRX-CT93-0411.

References

[1] G. Benfatto, E. Marinari, and E. Olivieri. Some numerical results on the block spin transformation for the $2d$ Ising model at the critical point. *J. Stat. Phys.*, 78:731-757, 1995.

[2] C. Cammarota. The large block spin interaction. *Nuovo Cimento*, B 96:1-16, 1986.

[3] E. A. Carlen and A. Soffer. Entropy production by block spin summation and central limit theorems . *Comm. Math. Phys.*, 140:339-371, 1992.

[4] M. Cassandro and G. Gallavotti. The Lavoisier law and the critical point. *Nuovo Cimento*, B25:691-705, 1975.

[5] R. L. Dobrushin and S. B. Shlosman. Completely analytical interactions: constructive description. *J. Stat. Phys.*, 46:983–1014, 1987.

[6] R. Fernández and C.-Ed. Pfister. Non-quasilocality of projections of Gibbs measures. EPFL preprint, 1994.

[7] R. B. Griffiths and P. A. Pearce. Position-space renormalization-group transformations: Some proofs and some problems. *Phys. Rev. Lett.*, 41:917–920, 1978.

[8] R. B. Griffiths and P. A. Pearce. Mathematical properties of position-space renormalization-group transformations. *J. Stat. Phys.*, 20:499–545, 1979.

[9] D. Iagolnitzer and B. Souillard. Random fields and limit theorems. In J. Fritz, J. L.Lebowitz, and D. Szász, editors, *Random Fields (Esztergom, 1979), Vol. II*, pages 573-592.North-Holland, Amsterdam, 1981.

[10] R. B. Israel. Banach algebras and Kadanoff transformations. In J. Fritz, J. L. Lebowitz, and D. Szász, editors, *Random Fields (Esztergom, 1979), Vol. II*, pages 593–608. North-Holland, Amsterdam, 1981.

[11] I. A. Kashapov. Justification of the renormalization-group method. *Theor. Math. Phys.*, 42:184-186, 1980.

[12] T. Kennedy. Some rigorous results on majority rule renormalization group transformations near the critical point. *J. Stat. Phys.*, 72:15–37, 1993.

[13] T. Kennedy. Abstract, Budapest meeting on disordered systems and statistical physics, rigorous results , august 1995.

[14] J. Lörinczi. Some results on the projected two-dimensional Ising model. In M. Fannes, C. Maes, and A. Verbeure, editors, *On Three Levels*, pages 373–380, New York and London, 1994. Plenum Press.

[15] J. Lörinczi. Groningen Ph.D. dissertation, 1995,to appear.

[16] J. Lörinczi and K. Vande Velde. A note on the projection of Gibbs measures. *J. Stat. Phys.*, 77:881-887, 1994.

[17] C. Maes and K. Vande Velde. The (non-)Gibbsian nature of states invariant under stochastic transformations. *Physica*A, 206:587-603, 1994.

[18] F. Martinelli and E. Olivieri. Some remarks on pathologies of renormalization-group transformations. *J. Stat. Phys.*, 72:1169–1177, 1993.

[19] F. Martinelli and E. Olivieri. Instability of renormalization-group pathologies under decimation. *J. Stat. Phys.*, 79:25-42, 1995.

[20] F. Martinelli and E. Olivieri. Approach to equilibrium of Glauber dynamics in the one phase region I. *Comm. Math. Phys.*, 161:447-486, 1994.

[21] F. Martinelli and E. Olivieri. Approach to equilibrium of Glauber dynamics in the one phase region II. *Comm. Math. Phys.*, 161:487-515, 1994.

[22] F. Martinelli, E. Olivieri and R. H. Schonmann. For 2-D lattice spin systems weak mixing implies strong mixing. *Comm. Math. Phys.*, 165:33-48, 1994.

[23] C. -Ed. Pfister and K. Vande Velde. Almost sure quasilocality in the random cluster model. *J. Stat. Phys.*, 79:765-774, 1995.

[24] J. Salas. Low temperature series for renormalized operators: the ferromagnetic square Ising model. New York University preprint, 1994.

[25] R. H. Schonmann and S. B. Shlosman. Complete analyticity for 2D Ising completed. *Comm. Math. Phys.*, 170:453-482, 1995.

[26] A. C. D. van Enter, R. Fernández, and R. Kotecký. Pathological behavior of renormalization group maps at high fields and above the transition temperature. *J. Stat. Phys.*, 79:969-992, 1995.

[27] A. C. D. van Enter, R. Fernández, and A. D. Sokal. Renormalization transformations in the vicinity of first-order phase transitions: What can and cannot go wrong. *Phys. Rev. Lett.*, 66:3253–3256, 1991.

[28] A. C. D. van Enter, R. Fernández, and A. D. Sokal. Regularity properties and pathologies of position-space renormalization-group transformations: Scope and limitations of Gibbsian theory. *J. Stat. Phys.*, 72:879–1167, 1993.

[29] K. Vande Velde. Private communication.

[30] K. Vande Velde. On the question of quasilocality in large systems of locally interacting components. K.U.Leuven Ph.D. dissertation, 1995