August 30, 2000

Proof of the Boltzmann-Sinai Ergodic Hypothesis for Typical Hard Disk Systems

Nándor Simányi[1]

University of Alabama at Birmingham
Department of Mathematics
Campbell Hall, Birmingham, AL 35294 U.S.A.
E-mail: simanyi@math.uab.edu

Abstract. We consider the system of N (≥ 2) hard disks of masses $m_1, \dots, m_N$ and radius r in the flat unit torus $\mathbb{T}^2$. We prove the ergodicity (actually, the B-mixing property) of such systems for almost every selection $(m_1, \dots, m_N; r)$ of the outer geometric parameters.

Primary subject classification: 37D50

Secondary subject classification: 34D05

§1. Introduction

Hard ball systems or, a bit more generally, mathematical billiards constitute an important and quite interesting family of dynamical systems being intensively studied by dynamicists and researchers of mathematical physics, as well. These dynamical systems pose many challenging mathematical questions, most of them concerning the ergodic (mixing) properties of such systems. The introduction of hard ball systems and the first major steps in their investigations date back to the 40's and 60's, see Krylov's paper [K(1942)] and Sinai's groundbreaking works [Sin(1963)] and [Sin(1970)], in which the author — among other things — formulated the modern version of Boltzmann's ergodic hypothesis (what we call today the Boltzmann–Sinai ergodic hypothesis) by claiming that every hard ball system in a flat torus is ergodic, of course after fixing the values of the trivial flow-invariant quantities. In the articles [Sin(1970)] and [B-S(1973)] Bunimovich and Sinai proved this hypothesis for two hard disks on the two-dimensional unit torus $\mathbb{T}^2$. The generalization of this result to higher dimensions $\nu > 2$ took fourteen years, and was

[1]Research partially supported by the Hungarian National Foundation for Scientific Research, grants OTKA-26176 and OTKA-29849.

done by Chernov and Sinai in [S-Ch(1987)]. Although the model of two hard balls in $\mathbb{T}^\nu$ is already rather involved technically, it is still a so called strictly dispersive billiard system, i. e. such that the smooth components of the boundary $\partial\mathbf{Q}$ of the configuration space are strictly concave from outside $\mathbf{Q}$. (They are bending away from $\mathbf{Q}$.) The billiard systems of more than two hard balls in $\mathbb{T}^\nu$ are no longer strictly dispersive, but just semi-dispersive (strict concavity of the smooth components of $\partial\mathbf{Q}$ is lost, merely concavity persists), and this circumstance causes a lot of additional technical troubles in their study. In the series of my joint papers with A. Krámli and D. Szász [K-S-Sz(1989)], [K-S-Sz(1990)], [K-S-Sz(1991)], and [K-S-Sz(1992)] we developed several new methods, and proved the ergodicity of more and more complicated semi-dispersive billiards culminating in the proof of ergodicity of four billiard balls in the torus $\mathbb{T}^\nu$ ($\nu \geq 3$), [K-S-Sz(1992)]. Then, in 1992, Bunimovich, Liverani, Pellegrinotti and Sukhov [B-L-P-S(1992)] were able to prove the ergodicity for some systems with an arbirarily large number of hard balls. The shortcoming of their model, however, is that, on one hand, they restrict the types of all feasible ball-to-ball collisions, on the other hand, they introduce some extra scattering effect with the collisions at the strictly concave wall of the container. The only result with an arbitrarily large number of balls in a flat unit torus $\mathbb{T}^\nu$ was achieved in [Sim(1992-A-B)], where the author managed to prove the ergodicity (actually, the K-mixing property) of N hard balls in $\mathbb{T}^\nu$, provided that $N \leq \nu$. The annoying shortcoming of that result is that the larger the number of balls N is, larger and larger dimension ν of the ambient container is required by the method of the proof.

On the other hand, if someone considers a hard ball system in an elongated torus which is long in one direction but narrow in the others, so that the balls must keep their cyclic order in the "long direction" (Sinai's "pencase" model), then the technical difficulties can be handled, thanks to the fact that the collisions of ballss are now restricted to neighbouring pairs. The hyperbolicity of such models in three dimensions and the ergodicity in dimension four have been proved in [S-Sz(1995)].

The positivity of the metric entropy for several systems of hard balls can be proven relatively easily, as was shown in the paper [W(1988)]. The articles [L-W(1995)] and [W(1990)] are nice surveys describing a general setup leading to the technical problems treated in a series of research papers. For a comprehensive survey of the results and open problems in this field, see [Sz(1996)].

Pesin's theory [P(1977)] on the ergodic properties of non-uniformly hyperbolic, smooth dynamical systems has been generalized substantially to dynamical systems with singularities (and with a relatively mild behaviour near the singularities) by A. Katok and J-M. Strelcyn [K-S(1986)]. Since then, the so called Pesin's and Katok-Strelcyn's theories have become part of the folklore in the theory of dynamical systems. They claim that — under some mild regularity conditions, especially near the singularities — every non-uniformly hyperbolic and ergodic flow enjoys the Kolmogorov-mixing property, shortly the K-mixing property.

Later on it was discovered and proven in [C-H(1996)] and [O-W(1998)] that the above metioned fully hyperbolic and ergodic flows with singularities turn out to be automatically having the Bernoulli mixing (B-mixing) property. It is worth noting here that almost every semi-dispersive billiard system, especially every hard ball system, enjoys those mild regularity conditions imposed on the systems (as axioms) by [K-S(1986)], [C-H(1996)], and [O-W(1998)]. In other words, for a hard ball flow $(\mathbf{M}, \{S^t\}, \mu)$ the (global) ergodicity of the systems actually implies its full

hyperbolicity and the B-mixing property, as well.

Finally, in our latest joint venture with D. Szász [S-Sz(1999)], we prevailed over the difficulty caused by the low value of the dimension ν by developing a brand new algebraic approach for the study of hard ball systems. That result, however, only establishes complete hyperbolicity (nonzero Lyapunov exponents almost everywhere) for N balls in $\mathbb{T}^\nu$. The ergodicity appeared to be a much harder task.

We note, however, that the algebraic method developed in [S-Sz(1999)] gets an essential application in the proof of Proposition 3.1 below.

Consider the ν-dimensional ($\nu \geq 2$), standard, flat, unit torus $\mathbb{T}^\nu = \mathbb{R}^\nu/\mathbb{Z}^\nu$ as the vessel containing N (≥ 2) hard balls (spheres) $B_1, \dots, B_N$ with positive masses $m_1, \dots, m_N$ and (just for simplicity) common radius $r > 0$. We always assume that the radius $r > 0$ is not too big, so that even the interior of the arising configuration space $\mathbf{Q}$ is connected. Denote the center of the ball B_i by $q_i \in \mathbb{T}^\nu$, and let $v_i = \dot{q}_i$ be the velocity of the i-th particle. We investigate the uniform motion of the balls $B_1, \dots, B_N$ inside the container $\mathbb{T}^\nu$ with half a unit of total kinetic energy: $E = \frac{1}{2}\sum_{i=1}^N m_i||v_i||^2 = \frac{1}{2}$. We assume that the collisions between balls are perfectly elastic. Since — beside the kinetic energy E — the total momentum $I = \sum_{i=1}^N m_i v_i \in \mathbb{R}^\nu$ is also a trivial first integral of the motion, we make the standard reduction $I = 0$. Due to the apparent translation invariance of the arising dynamical system, we factorize out the configuration space with respect to uniform spatial translations as follows: $(q_1, \dots, q_N) \sim (q_1 + a, \dots, q_N + a)$ for all translation vectors $a \in \mathbb{T}^\nu$. The configuration space $\mathbf{Q}$ of the arising flow is then the factor torus $\Big((\mathbb{T}^\nu)^N / \sim\Big) \cong \mathbb{T}^{\nu(N-1)}$ minus the cylinders

$$C_{i,j} = \Big\{(q_1, \dots, q_N) \in \mathbb{T}^{\nu(N-1)} \colon \operatorname{dist}(q_i, q_j) < 2r\Big\}$$

($1 \leq i < j \leq N$) corresponding to the forbidden overlap between the i-th and j-th cylinders. Then it is easy to see that the compound configuration point

$$q = (q_1, \dots, q_N) \in \mathbf{Q} = \mathbb{T}^{\nu(N-1)} \setminus \bigcup_{1 \leq i < j \leq N} C_{i,j}$$

moves in $\mathbf{Q}$ uniformly with unit speed and bounces back from the boundaries $\partial C_{i,j}$ of the cylinders $C_{i,j}$ according to the classical law of geometric optics: the angle of reflection equals the angle of incidence. More precisely: the post-collision velocity v^+ can be obtained from the pre-collision velocity v^- by the orthogonal reflection across the tangent hyperplane of the boundary $\partial\mathbf{Q}$ at the point of collision. Here we must emphasize that the phrase "ortogonal" should be understood with respect to the natural Riemannian metric (the so called mass metric) $||dq||^2 = \sum_{i=1}^N m_i||dq_i||^2$ in the configuration space $\mathbf{Q}$. For the normalized Liouville measure μ of the arising flow $\{S^t\}$ we obviously have $d\mu = \text{const} \cdot dq \cdot dv$, where dq is the Riemannian volume in $\mathbf{Q}$ induced by the above metric and dv is the surface measure (determined by the restriction of the Riemannian metric above) on the sphere of compound velocities

$$\mathbb{S}^{\nu(N-1)-1} = \left\{(v_1, \dots, v_N) \in (\mathbb{R}^\nu)^N \colon \sum_{i=1}^N m_i v_i = 0 \text{ and } \sum_{i=1}^N m_i||v_i||^2 = 1\right\}.$$

The phase space $\mathbf{M}$ of the flow $\{S^t\}$ is the unit tangent bundle $\mathbf{Q} \times \mathbb{S}^{d-1}$ of the configuration space $\mathbf{Q}$. (We will always use the shorthand notation $d = \nu(N-1)$ for the dimension of the billiard table $\mathbf{Q}$.) We must, however, note here that at the boundary $\partial\mathbf{Q}$ of $\mathbf{Q}$ one has to glue together the pre-collision and post-collision velocities in order to form the phase space $\mathbf{M}$, so $\mathbf{M}$ is equal to the unit tangent bundle $\mathbf{Q} \times \mathbb{S}^{d-1}$ modulo this identification.

A bit more detailed definition of hard ball systems with arbitrary masses, as well as their role in the family of cylindric billiards, can be found in §4 of [S-Sz(2000)] and in §1 of [S-Sz(1999)]. We denote the arising flow by $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$.

The joint work by Ya. G. Sinai and N. I. Chernov [S-Ch(1987)] paved the way for further fundamental results concerning the ergodicity of $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$. They proved there a strong result on local ergodicity: An open neighbourhood $U \subset \mathbf{M}$ of every phase point with a hyperbolic trajectory (and with at most one singularity on its trajectory) belongs to a single ergodic component of the billiard flow $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$, of course, modulo the zero measure sets. An immediate consequence of this result is the (hyperbolic) ergodicity of the hard ball systems with $N = 2$ and $\nu \geq 2$.

In the series of articles [K-S-Sz(1989)], [K-S-Sz(1991)], [K-S-Sz(1992)], and [Sim(1992-A-B)] the authors developed a powerful, three-step strategy for proving the (hyperbolic) ergodicity of hard ball systems. First of all, all these proofs are inductions on the number N of balls involved in the problem. Secondly, the induction step itself consists of the following three major steps:

Step I. To prove that every non-singular (i. e. smooth) trajectory segment $S^{[a,b]}x_0$ with a "combinatorially rich" (in a well defined sense) symbolic collision sequence is automatically sufficient (or, in other words, "geometrically hyperbolic", see below in §2), provided that the phase point x_0 does not belong to a countable union J of smooth submanifolds with codimension at least two. (Containing the exceptional phase points.)

The exceptional set J featuring this result is negligible in our dynamical considerations — it is a so called slim set. For the basic properties of slim sets, see §2.7 below.

Step II. Assume the induction hypothesis, i. e. that all hard ball systems with n balls ($n < N$) are (hyperbolic and) ergodic. Prove that then there exists a slim set $S \subset \mathbf{M}$ (see §2.7) with the following property: For every phase point $x_0 \in \mathbf{M}\setminus S$ the entire trajectory $S^{\mathbb{R}}x_0$ contains at most one singularity and its symbolic collision sequence is combinatorially rich, just as required by the result of Step I.

Step III. By using again the induction hypothesis, prove that almost every singular trajectory is sufficient in the time interval $(t_0, +\infty)$, where t_0 is the time moment of the singular reflection. (Here the phrase "almost every" refers to the volume defined by the induced Riemannian metric on the singularity manifolds.)

We note here that the almost sure sufficiency of the singular trajectories (featuring Step III) is an essential condition for the proof of the celebrated Theorem on Local Ergodicity for semi-dispersive billiards proved by Sinai and Chernov, see Theorem 5 in [S-Ch(1987)]. Under this assumption that theorem asserts that a suitable, open neighbourhood U_0 of any sufficient phase point $x_0 \in \mathbf{M}$ (with at most one singularity on its trajectory) belongs to a single ergodic component of the billiard flow $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$.

In an inductive proof of ergodicity, steps I and II together would ensure that there exists an arcwise connected set $C \subset \mathbf{M}$ with full measure, such that every phase point $x_0 \in C$ is sufficient with at most one singularity on its trajectory. Then the cited Theorem on Local Ergodicity (now taking advantage of the result of Step III) states that for every phase point $x_0 \in C$ an open neighbourhood U_0 of x_0 belongs to one ergodic component of the flow. Finally, the connectedness of the set C and $\mu(\mathbf{M} \setminus C) = 0$ easily imply that the flow $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ (now with N balls) is indeed ergodic, and actually fully hyperbolic, as well.

In the series of articles [K-S-Sz(1991)], [K-S-Sz(1992)] the authors followed the strategy outlined above and obtained the (hyperbolic) ergodicity of three and four hard balls, respectively. Technically speaking, in those papers we always assumed tacitly that the masses of balls are equal.

The twin papers [Sim(1992-A-B)] of mine brought new topological and geometric tools to attack the problem of ergodicity. Namely, in [Sim(1992-A)], a brand new topological method was developed, and that resulted in settling Step II of the induction, once forever.

In the subsequent paper [Sim(1992-B)] a new combinatorial approach for handling Step I was developed in the case when the dimension ν of the toroidal container is not less than the number of balls N. This proves the ergodicity of every hard ball system with $\nu \geq N$.

The main result of this article is our

Theorem. *In the case $\nu = 2$ (i. e. for hard disks in $\mathbb{T}^2$) for almost every selection $(m_1, \ldots, m_N; r)$ of the outer geometric parameters it is true that the billiard flow $(\mathbf{M}_{\vec{m},r}, \{S^t\}, \mu_{\vec{m},r})$ of the N-disk system is ergodic and completely hyperbolic. Then, following from the results of Chernov–Haskell [C-H(1996)] and Ornstein–Weiss [O-W(1998)], such a semi-dispersive billiard system actually enjoys the B-mixing property, as well.*

A few remarks concerning this theorem are now in place. First of all, here we present an inductive proof following the above drafted three-step strategy I–III amended in such a way that the exceptional set J featuring Step I is no longer a countable union of codimension-two (i. e. at least two) sets but, rather, it is only a countable union of proper (i. e. of codimension at least one) submanifolds. This shortcoming of Step I makes it possible (in principle, at least) that countably many open ergodic components $C_1, C_2, \ldots$ coexist in such a way that they are separated from each other by codimension-one, smooth, exceptional submanifolds J of $\mathbf{M}$ featuring Step I. The main contents of the present paper is to exclude this possibility, and this is precisely what is going on in §4–7 below. It is just this proof of the non-existence of separating manifolds J that essentially uses the dimension condition $\nu = 2$.

The other remark concerns the fact that — at least in principle — an unspecified zero measure set of the outer geometric parameters $(m_1, \ldots, m_N; r)$ has to be dropped in the Theorem. But why? The reason is the same as for the dropping of the zero set in the main theorem of [S-Sz(1999)], in which we proved that a hard ball system (in any given dimension $\nu \geq 2$) is almost surely fully hyperbolic, that is, its relevant Lyapunov exponents are nonzero almost everywhere. In fact, in the proof of Proposition 3.1 below (which is required for the proof of the Chernov-Sinai

Ansatz, i. e. Step III) we successfully applied the algebraic method developed in [S-Sz(1999)]. Proposition 3.1 asserts that the intersection of the exceptional set J (featuring Step I) and the singularity set $\mathcal{SR}^+$ (see the two paragraphs preceding Proposition 3.1) has at least two codimensions, that is, J and $\mathcal{SR}^+$ cannot even locally coincide.

The paper is organized as follows. After putting forward the prerequisites in §2, in the subsequent section we carry out the inductive proof of the ergodicity by assuming the non-existence of the separating manifolds J. Then all remaining sections 4–7 are devoted to the proof of the non-existence of J-manifolds. That proof contains a lot of new geometric ideas. Finally, in §8, two remarks conclude the article. One of them regards the role of Proposition 3.1 in the entire proof, while the other one applies the method of [S-W(1989)] to prove the striking fact that a typical (i. e. an ergodic, or B-mixing) hard disk system retains its B-mixing property even if one omits the translation factorization $(q_1, \dots, q_N) \sim (q_1 + a, \dots, q_N + a)$ of the configuration space, despite the fact that the dropping of this factorization introduces 2 zero Lyapunov exponents!

§2. Prerequisites

2.1 Cylindric billiards. Consider the d-dimensional ($d \geq 2$) flat torus $\mathbb{T}^d = \mathbb{R}^d/\mathcal{L}$ supplied with the usual Riemannian inner product $\langle\, .\,,\, .\,\rangle$ inherited from the standard inner product of the universal covering space $\mathbb{R}^d$. Here $\mathcal{L} \subset \mathbb{R}^d$ is assumed to be a lattice, i. e. a discrete subgroup of the additive group $\mathbb{R}^d$ with $\mathrm{rank}(\mathcal{L}) = d$. The reason why we want to allow general lattices, other than just the integer lattice $\mathbb{Z}^d$, is that otherwise the hard ball systems would not be covered. The geometry of the structure lattice $\mathcal{L}$ in the case of a hard ball system is significantly different from the geometry of the standard lattice $\mathbb{Z}^d$ in the standard Euclidean space $\mathbb{R}^d$, see later in this section.

The configuration space of a cylindric billiard is $\mathbf{Q} = \mathbb{T}^d \setminus (C_1 \cup \dots \cup C_k)$, where the cylindric scatterers C_i ($i = 1, \dots, k$) are defined as follows.

Let $A_i \subset \mathbb{R}^d$ be a so called lattice subspace of $\mathbb{R}^d$, which means that $\mathrm{rank}(A_i \cap \mathcal{L}) = \dim A_i$. In this case the factor $A_i/(A_i \cap \mathcal{L})$ is a subtorus in $\mathbb{T}^d = \mathbb{R}^d/\mathcal{L}$ which will be taken as the generator of the cylinder $C_i \subset \mathbb{T}^d$, $i = 1, \dots, k$. Denote by $L_i = A_i^{\perp}$ the orthocomplement of A_i in $\mathbb{R}^d$. Throughout this article we will always assume that $\dim L_i \geq 2$. Let, furthermore, the numbers $r_i > 0$ (the radii of the spherical cylinders C_i) and some translation vectors $t_i \in \mathbb{T}^d = \mathbb{R}^d/\mathcal{L}$ be given. The translation vectors t_i play a crucial role in positioning the cylinders C_i in the ambient torus $\mathbb{T}^d$. Set

$$C_i = \left\{ x \in \mathbb{T}^d \colon \mathrm{dist}\,(x - t_i, A_i/(A_i \cap \mathcal{L})) < r_i \right\}.$$

In order to avoid further unnecessary complications, we always assume that the interior of the configuration space $\mathbf{Q} = \mathbb{T}^d \setminus (C_1 \cup \dots \cup C_k)$ is connected. The phase space $\mathbf{M}$ of our cylindric billiard flow will be the unit tangent bundle of $\mathbf{Q}$ (modulo some natural glueings at its boundary), i. e. $\mathbf{M} = \mathbf{Q} \times \mathbb{S}^{d-1}$. (Here $\mathbb{S}^{d-1}$ denotes the unit sphere of $\mathbb{R}^d$.)

The dynamical system $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$, where S^t $(t \in \mathbb{R})$ is the dynamics defined by the uniform motion inside the domain $\mathbf{Q}$ and specular reflections at its boundary (at the scatterers), and μ is the Liouville measure, is called a cylindric billiard flow we want to investigate.

We note that the cylindric billiards — defined above — belong to the wider class of so called semi-dispersive billiards, which means that the smooth components $\partial\mathbf{Q}_i$ of the boundary $\partial\mathbf{Q}$ of the configuration space $\mathbf{Q}$ are (not necessarily strictly) concave from outside of $\mathbf{Q}$, i. e. they are bending away from the interior of $\mathbf{Q}$. As to the notions and notations in connection with semi-dispersive billiards, the reader is kindly referred to the article [K-S-Sz(1990)].

2.2 Hard ball systems. Hard ball systems in the standard unit torus $\mathbb{T}^\nu = \mathbb{R}^\nu/\mathbb{Z}^\nu$ $(\nu \geq 2)$ with positive masses $m_1, \ldots, m_N$ are described (for example) in §1 of [S-Sz(1999)]. These are the dynamical systems describing the motion of N (≥ 2) hard balls with radii $r_1, r_2, \ldots, r_N$ and positive masses $m_1, \ldots, m_N$ in the standard unit torus $\mathbb{T}^\nu = \mathbb{R}^\nu/\mathbb{Z}^\nu$. (For simplicity we assume that these radii have the common value r.) The center of the i-th ball is denoted by q_i $(\in \mathbb{T}^\nu)$, its time derivative is $v_i = \dot{q}_i$, $i = 1, \ldots, N$. One uses the standard reduction of kinetic energy $E = \frac{1}{2}\sum_{i=1}^N m_i||v_i||^2 = \frac{1}{2}$. The arising configuration space (still without the removal of the scattering cylinders $C_{i,j}$) is the torus

$$\mathbb{T}^{\nu N} = (\mathbb{T}^\nu)^N = \{(q_1, \ldots, q_N)\colon\ q_i \in \mathbb{T}^\nu,\ i = 1, \ldots, N\}$$

supplied with the Riemannian inner product (the so called mass metric)

$$\langle v, v'\rangle = \sum_{i=1}^N m_i \langle v_i, v_i'\rangle \tag{2.2.1}$$

in its common tangent space $\mathbb{R}^{\nu N} = (\mathbb{R}^\nu)^N$. Now the Euclidean space $\mathbb{R}^{\nu N}$ with the inner product (2.2.1) plays the role of $\mathbb{R}^d$ in the original definition of cylindric billiards, see §2.1 above.

The generator subspace $A_{i,j} \subset \mathbb{R}^{\nu N}$ $(1 \leq i < j \leq N)$ of the cylinder $C_{i,j}$ (describing the collisions between the i-th and j-th balls) is given by the equation

$$A_{i,j} = \left\{(q_1, \ldots, q_N) \in (\mathbb{R}^\nu)^N\colon\ q_i = q_j\right\}, \tag{2.2.2}$$

see (4.3) in [S-Sz(2000)]. Its orthocomplement $L_{i,j} \subset \mathbb{R}^{\nu N}$ is then defined by the equation

(2.2.3)

$$L_{i,j} = \left\{(q_1, \ldots, q_N) \in (\mathbb{R}^\nu)^N\colon\ q_k = 0 \text{ for } k \neq i, j, \text{ and } m_i q_i + m_j q_j = 0\right\},$$

see (4.4) in [S-Sz(2000)]. Easy calculation shows that the cylinder $C_{i,j}$ (describing the overlap of the i-th and j-th balls) is indeed spherical and the radius of its base sphere is equal to $r_{i,j} = 2r\sqrt{\frac{m_i m_j}{m_i+m_j}}$, see §4, especially formula (4.6) in [S-Sz(2000)].

The structure lattice $\mathcal{L} \subset \mathbb{R}^{\nu N}$ is clearly the integer lattice $\mathcal{L} = \mathbb{Z}^{\nu N}$.

Due to the presence of an extra invariant quantity $I = \sum_{i=1}^N m_i v_i$, one usually makes the reduction $\sum_{i=1}^N m_i v_i = 0$ and, correspondingly, factorizes the configuration space with respect to uniform spatial translations:

$$(q_1, \dots, q_N) \sim (q_1 + a, \dots, q_N + a), \quad a \in \mathbb{T}^\nu.$$

The natural, common tangent space of this reduced configuration space is then

$$\mathcal{Z} = \left\{ (v_1, \dots, v_N) \in (\mathbb{R}^\nu)^N : \sum_{i=1}^N m_i v_i = 0 \right\} = \left(\bigcap_{i<j} A_{i,j} \right)^\perp = (\mathcal{A})^\perp \tag{2.2.4}$$

supplied again with the inner product (2.2.1), see also (4.1) and (4.2) in [S-Sz(2000)]. The base spaces $L_{i,j}$ of (2.2.3) are obviously subspaces of $\mathcal{Z}$, and we take $\tilde{A}_{i,j} = A_{i,j} \cap \mathcal{Z} = P_{\mathcal{Z}}(A_{i,j})$ as the orthocomplement of $L_{i,j}$ in $\mathcal{Z}$. (Here $P_{\mathcal{Z}}$ denotes the orthogonal projection onto the space $\mathcal{Z}$.)

Note that the configuration space of the reduced system (with $\sum_{i=1}^N m_i v_i = 0$) is naturally the torus $\mathbb{R}^{\nu N}/(\mathcal{A} + \mathbb{Z}^{\nu N}) = \mathcal{Z}/P_{\mathcal{Z}}(\mathbb{Z}^{\nu N})$.

2.3 Trajectory Branches. We are going to briefly describe the discontinuity of the flow $\{S^t\}$ caused by a multiple collision at time t_0. Assume first that the pre–collision velocities of the particles are given. What can we say about the possible post–collision velocities? Let us perturb the pre–collision phase point (at time $t_0 - 0$) infinitesimally, so that the collisions at $\sim t_0$ occur at infinitesimally different moments. By applying the collision laws to the arising finite sequence of collisions, we see that the post-collision velocities are fully determined by the time– ordering of the considered collisions. Therefore, the collection of all possible time-orderings of these collisions gives rise to a finite family of continuations of the trajectory beyond t_0. They are called the **trajectory branches**. It is quite clear that similar statements can be said regarding the evolution of a trajectory through a multiple collision **in reverse time**. Furthermore, it is also obvious that for any given phase point $x_0 \in \mathbf{M}$ there are two, ω-high trees $\mathcal{T}_+$ and $\mathcal{T}_-$ such that $\mathcal{T}_+$ ($\mathcal{T}_-$) describes all the possible continuations of the positive (negative) trajectory $S^{[0,\infty)}x_0$ ($S^{(-\infty,0]}x_0$). (For the definitions of trees and for some of their applications to billiards, cf. the beginning of §5 in [K-S-Sz(1992)].) It is also clear that all possible continuations (branches) of the whole trajectory $S^{(-\infty,\infty)}x_0$ can be uniquely described by all possible pairs (B_-, B_+) of ω-high branches of the trees $\mathcal{T}_-$ and $\mathcal{T}_+$ ($B_- \subset \mathcal{T}_-, B_+ \subset \mathcal{T}_+$).

Finally, we note that the trajectory of the phase point x_0 has exactly two branches, provided that $S^t x_0$ hits a singularity for a single value $t = t_0$, and the phase point $S^{t_0}x_0$ does not lie on the intersection of more than one singularity manifolds. (In this case we say that the trajectory of x_0 has a "single singularity".)

2.4 Neutral Subspaces, Advance, and Sufficiency. Consider a **nonsingular** trajectory segment $S^{[a,b]}x$. Suppose that a and b are **not moments of collision**.

Definition 2.4.1. *The neutral space $\mathcal{N}_0(S^{[a,b]}x)$ of the trajectory segment $S^{[a,b]}x$ at time zero ($a < 0 < b$) is defined by the following formula:*

$$\mathcal{N}_0(S^{[a,b]}x) = \big\{ W \in \mathcal{Z} \colon \exists (\delta > 0) \text{ s. t. } \forall \alpha \in (-\delta, \delta)$$
$$p\left(S^a\left(Q(x) + \alpha W, V(x)\right)\right) = p(S^a x) \text{ and } p\left(S^b\left(Q(x) + \alpha W, V(x)\right)\right) = p(S^b x) \big\}.$$

(The formula for the tangent space $\mathcal{Z}$ can be found in (2.2.4).)

It is known (see (3) in §3 of [S-Ch (1987)]) that $\mathcal{N}_0(S^{[a,b]}x)$ is a linear subspace of $\mathcal{Z}$ indeed, and $V(x) \in \mathcal{N}_0(S^{[a,b]}x)$. The neutral space $\mathcal{N}_t(S^{[a,b]}x)$ of the segment $S^{[a,b]}x$ at time $t \in [a,b]$ is defined as follows:

$$\mathcal{N}_t(S^{[a,b]}x) = \mathcal{N}_0\left(S^{[a-t,b-t]}(S^t x)\right). \tag{2.4.2}$$

It is clear that the neutral space $\mathcal{N}_t(S^{[a,b]}x)$ can be canonically identified with $\mathcal{N}_0(S^{[a,b]}x)$ by the usual identification of the tangent spaces of $\mathbf{Q}$ along the trajectory $S^{(-\infty,\infty)}x$ (see, for instance, §2 of [K-S-Sz(1990)]).

Our next definition is that of the **advance**. Consider a non-singular orbit segment $S^{[a,b]}x$ with the symbolic collision sequence $\Sigma = (\sigma_1, \ldots, \sigma_n)$ $(n \geq 1)$ as at the beginning of the present section. For $x = (Q,V) \in \mathbf{M}$ and $W \in \mathcal{Z}$, $\|W\|$ sufficiently small, denote $T_W(Q,V) := (Q+W, V)$.

Definition 2.4.3. *For any $1 \leq k \leq n$ and $t \in [a,b]$, the advance*

$$\alpha(\sigma_k)\colon \mathcal{N}_t(S^{[a,b]}x) \to \mathbb{R}$$

of the collision σ_k is the unique linear extension of the linear functional $\alpha(\sigma_k)$ defined in a sufficiently small neighbourhood of the origin of $\mathcal{N}_t(S^{[a,b]}x)$ in the following way:

$$\alpha(\sigma_k)(W) := t_k(x) - t_k(S^{-t}T_W S^t x).$$

It is now time to bring up the basic notion of **sufficiency** (or, sometimes it is also called hyperbolicity) of a trajectory (segment). This is the utmost important necessary condition for the proof of the fundamental theorem for semi-dispersive billiards, see Condition (ii) of Theorem 3.6 and Definition 2.12 in [K-S-Sz(1990)].

Definition 2.4.4.

(1) *The nonsingular trajectory segment $S^{[a,b]}x$ (a and b are supposed not to be moments of collision) is said to be* **sufficient** *if and only if the dimension of $\mathcal{N}_t(S^{[a,b]}x)$ ($t \in [a,b]$) is minimal, i.e. dim $\mathcal{N}_t(S^{[a,b]}x) = 1$.*

(2) *The trajectory segment $S^{[a,b]}x$ containing exactly one singularity (a so called "single singularity", see above) is said to be* **sufficient** *if and only if both branches of this trajectory segment are sufficient.*

Definition 2.4.5. *The phase point $x \in \mathbf{M}$ with at most one singularity is said to be sufficient if and only if its whole trajectory $S^{(-\infty,\infty)}x$ is sufficient, which means, by definition, that some of its bounded segments $S^{[a,b]}x$ are sufficient.*

In the case of an orbit $S^{(-\infty,\infty)}x$ with a single singularity, sufficiency means that both branches of $S^{(-\infty,\infty)}x$ are sufficient.

2.5 No accumulation (of collisions) in finite time. By the results of Vaserstein [V(1979)], Galperin [G(1981)] and Burago-Ferleger-Kononenko [B-F-K(1998)], in a semi-dispersive billiard flow there can only be finitely many collisions in finite time intervals, see Theorem 1.1 in [B-F-K(1998)]. Thus, the dynamics is well defined as long as the trajectory does not hit more than one boundary components at the same time.

2.6 Collision graphs. Let $S^{[a,b]}x$ be a nonsingular, finite trajectory segment with the collisions $\sigma_1, \ldots, \sigma_n$ listed in time order. (Each σ_k is an unordered pair (i,j) of different labels $i, j \in \{1, 2, \ldots, N\}$.) The graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ with vertex set $\mathcal{V} = \{1, 2, \ldots, N\}$ and set of edges $\mathcal{E} = \{\sigma_1, \ldots, \sigma_n\}$ is called the **collision graph** of the orbit segment $S^{[a,b]}x$.

2.7 Slim sets. We are going to summarize the basic properties of codimension-two subsets A of a smooth manifold M. Since these subsets A are just those negligible in our dynamical discussions, we shall call them **slim**. As to a broader exposition of the issues, see [E(1978)] or §2 of [K-S-Sz(1991)].

Note that the dimension $\dim A$ of a separable metric space A is one of the three classical notions of topological dimension: the covering (Čech-Lebesgue), the small inductive (Menger-Urysohn), or the large inductive (Brouwer-Čech) dimension. As it is known from general general topology, all of them are the same for separable metric spaces.

Definition 2.7.1. *A subset A of M is called slim if and only if A can be covered by a countable family of codimension-two (i. e. at least two) closed sets of μ–measure zero, where μ is a smooth measure on M. (Cf. Definition 2.12 of [K-S-Sz(1991)].)*

Property 2.7.2. *The collection of all slim subsets of M is a σ-ideal, that is, countable unions of slim sets and arbitrary subsets of slim sets are also slim.*

Lemma 2.7.3. *A subset $A \subset M$ is slim if and only if for every $x \in A$ there exists an open neighbourhood U of x in M such that $U \cap A$ is slim. (Locality, cf. Lemma 2.14 of [K-S-Sz(1991)].)*

Property 2.7.4. *A closed subset $A \subset M$ is slim if and only if $\mu(A) = 0$ and $\dim A \leq \dim M - 2$.*

Property 2.7.5 (Integrability). *If $A \subset M_1 \times M_2$ is a closed subset of the product of two manifolds, and for every $x \in M_1$ the set*

$$A_x = \{y \in M_2 \colon (x, y) \in A\}$$

is slim in M_2, then A is slim in $M_1 \times M_2$.

The following lemmas characterize the codimension-one and codimension-two sets.

Lemma 2.7.6. *For any closed subset $S \subset M$ the following three conditions are equivalent:*

(i) $\dim S \leq \dim M - 2$;

(ii) *$\mathrm{int} S = \emptyset$ and for every open connected set $G \subset M$ the difference set $G \setminus S$ is also connected;*

(iii) *$\mathrm{int} S = \emptyset$ and for every point $x \in M$ and for any open neighbourhood V of x in M there exists a smaller open neighbourhood $W \subset V$ of the point x such that for every pair of points $y, z \in W \setminus S$ there is a continuous curve γ in the set $V \setminus S$ connecting the points y and z.*

(See Theorem 1.8.13 and Problem 1.8.E of [E(1978)].)

Lemma 2.7.7. *For any subset $S \subset M$ the condition* $\dim S \leq \dim M - 1$ *is equivalent to* $\mathrm{int} S = \emptyset$. *(See Theorem 1.8.10 of [E(1978)].)*

We recall an elementary, but important lemma (Lemma 4.15 of [K-S-Sz(1991)]). Let R_2 be the set of phase points $x \in \mathbf{M} \setminus \partial\mathbf{M}$ such that the trajectory $S^{(-\infty,\infty)}x$ has more than one singularities.

Lemma 2.7.8. *The set R_2 is a countable union of codimension-two smooth submanifolds of M and, being such, it is slim.*

The next lemma establishes the most important property of slim sets which gives us the fundamental geometric tool to connect the open ergodic components of billiard flows.

Lemma 2.7.9. *If M is connected, then the complement $M \setminus A$ of a slim set $A \subset M$ necessarily contains an arcwise connected, G_δ set of full measure. (See Property 3 of §4.1 in [K-S-Sz(1989)]. The G_δ sets are, by definition, the countable intersections of open sets.)*

§3. The Inductive Proof of the Theorem (Using the results of §4–7)

In this section we prove our theorem by using an induction on the number of disks N (≥ 2). Consider therefore an N-disk billiard flow

$$\left(\mathbf{M}_{\vec{m},r}, \{S^t\}_{t\in\mathbb{R}}, \mu_{\vec{m},r}\right) = \left(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu\right)$$

in the standard unit 2-torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ with the $N+1$-tuple of outer geometric parameters $(m_1, \ldots, m_N; r)$, for which even the interior of the phase space is connected, see the previous section.

As Corollary 3.24 and Lemma 4.2 of [Sim(2000)] state, there exists a positive integer $C(N)$ with the following property: If the non-singular trajectory segment $S^{[0,T]}x_0$ of $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ has at least $C(N)$ consecutive, connected collision graphs, then there exists an open neighbourhood U_0 of x_0 in $\mathbf{M}$ and a proper (i. e. of codimension at least one) algebraic set $N_0 \subset U_0$ such that $S^{[0,T]}y$ is sufficient (or, geometrically hyperbolic, see §2) for all $y \in U_0 \setminus N_0$.

Consider the $(2d-3)$-dimensional, compact cell complex $\mathcal{SR}^+$ of singular reflections $x = (q, v^+) \in \partial\mathbf{M}$ supplied with the outgoing (post collision) velocity v^+, so that the positive orbit is well defined, as long as there is no other singularity on $S^{(0,\infty)}x$. Recall that, as it follows from Lemma 4.1 of [K-S-Sz(1990)], the set of phase points with more than one singularities is a countable union of smooth submanifolds with codimension at least two (see (2.7.8) above), thus the positive semi-trajectory $S^{(0,\infty)}x$ is non-singular for ν-almost every $x \in \mathcal{SR}^+$, where ν denotes the hypersurface measure on $\mathcal{SR}^+$. Also recall that $2d - 1 = 4N - 5$ is the dimension of the phase space $\mathbf{M}$.

Let $\Sigma = (\sigma_1, \ldots, \sigma_n)$ be any fixed symbolic collision sequence with at least $C(N)$ consecutive, connected collision graphs, i. e. a so called $C(N)$-rich symbolic

sequence in the sense of Key Lemma 4.1 and Remark 4.1/b of [S-Sz(1999)]. Let, moreover, $x_0 \in \mathcal{SR}^+$ be an element of a $(2d-3)$-dimensional, open cell C of $\mathcal{SR}^+$ (that is, x_0 does not belong to the $(2d-4)$-skeleton of the $(2d-3)$-dimensional cell complex $\mathcal{SR}^+$) with a non-singular trajectory segment $S^{[0,T]}x_0$, for which the symbolic collision sequence is the given Σ. We will need the following generalization of Key Lemma 4.1 of [S-Sz(1999)], which claims that the exceptional algebraic set N_0 from above (containing all phase points $y \in U_0$ around x_0 for which $S^{[0,T]}y$ is not sufficient) cannot even locally coincide with the invariant hull of $\mathcal{SR}^+$.

Proposition 3.1. *For almost every $(N+1)$-tuple $(m_1, \ldots, m_N; r)$ of the outer geometric parameters the set*

$$\Big\{ y \in U_0 \cap C\colon\ S^{[0,T]}y \text{ is not sufficient} \Big\} \tag{3.2}$$

has an empty interior in C.

Proof (A brief outline). Since this generalization of Key Lemma 4.1 of [S-Sz(1999)] is a direct application of the proof of that lemma (in which application all steps of the mentioned proof need to be repeated with only minor changes), hereby we will only briefly sketch the proof by mainly sheding light on the important steps, during which we point out the differences between the original proof of Key Lemma 4.1 and its modification that proves Proposition 3.1 above. This sketch of the proof will be subdivided into 12 points, as follows.

1^o In order to facilitate the use of arithmetics for the kinetic variables, we lift the entire system to the universal covering space $\mathbb{R}^2$ of $\mathbb{T}^2$ by introducing the notion of adjustment vectors, see Propositions 3.1 and 3.3 of [S-Sz(1999)].

2^o We need to complexify the system, to introduce the algebraically independent initial variables, the polynomial equations defining the algebraic dynamics, the algebraic functions in terms of the initial kinetic variables, and the tower of fields made up by all kinetic variables of orbit segments with a symbolic collision sequence $(\sigma_1, \ldots, \sigma_n;\ a_1, \ldots, a_n)$, along the lines of pp. 49–54 of [S-Sz(1999)].

3^o We should introduce the complex neutral space $\mathcal{N}(\omega)$, just as in (3.21) of [S-Sz(1999)]. By dropping the factorization with respect to uniform spatial translations, the condition of sufficiency now becomes $\dim\mathcal{N}(\omega) = \nu + 1 = 3$.

4^o Just for technical reasons, the reciprocal $1/L$ of the size L of the container $\mathbb{T}^2_L = \mathbb{R}^2/L\cdot\mathbb{Z}^2$ (containing disks with unit radius) is replaced by the radius of disks r moving in the unit torus $\mathbb{R}^2/\mathbb{Z}^2$.

5^o By using the defining equation of the actual singularity containing the cell C, we eliminate one variable out of the initial ones to gain again algebraic independence, despite considering singular phase points on C. We express the sufficiency of $\omega = S^{[0,T]}x$ $(x \in C)$ in terms of the remaining, algebraically independent initial variables. Non-sufficiency again proves to be equivalent to the simultaneous vanishing of finitely many polynomials, in the spirit of Lemma 4.2 from [S-Sz(1999)].

6^o One reformulates the claim of Proposition 3.1 in terms of the initial kinetic variables. The negation of that assertion proves to be the identical vanishing of certain algebraic functions, see also Lemma 4.2 and its proof in [S-Sz(1999)].

7^o Use Property (A) and the concept of combinatorial richness of the symbolic sequence of $S^{[0,T]}x$ (of containing at least $C(N)$ consecutive, connected collision graphs), just like in Key Lemma 4.1 and Remark 4.1/b in [S-Sz(1999].

8^o Carry out an inductive proof for Proposition 3.1 above. The induction goes on with respect to the number of disks N (≥ 2), and this induction is independent of the outer induction to be carried out to prove the Theorem. The statement is obviously true for $N = 2$. We assume $N > 2$ and the induction hypothesis, and perform an indirect proof for the induction step by assuming the negation of Proposition 3.1 for the complexified N-disk system. By using the combinatorial richness formulated in Key Lemma 4.1 of [S-Sz(1999)], one selects a label $i \in \{1, 2, \ldots, N\}$ for the substitution $m_i = 0$, along the lines of Lemma 4.43 of [S-Sz(1999]. The substitution $m_i = 0$ results in a derived scheme $(\Sigma', \mathcal{A}')$ by also preserving Property (A), see Definition 4.11, Main Lemma 4.21, Remark 4.22, and Corollary 4.35 of [S-Sz(1999].

9^o Describe non-sufficiency in the case $m_i = 0$ along the lines of Lemma 4.9 from [S-Sz(1999].

10^o From the indirect assumption one obtains that the induction hypothesis is false for the $(N-1)$-disk system, just like in Lemma 4.40 of [S-Sz(1999)].

11^o From the complex version of the analogue of Key Lemma 4.1 one switches to the real case, just as in the fourth paragraph on page 88 of [S-Sz(1999].

12^o From the real version of the analogue of Key Lemma 4.1 one obtains Proposition 3.1 of this article by dropping a null set of $(N+1)$-tuples $(m_1, \ldots, m_N; r)$ of outer geometric parameters, precisely as in the first paragraph of page 93 of [S-Sz(1999)]. $\square$

By the results of Vaserstein [V(1979)], Galperin [G(1981)], and Burago-Ferleger-Kononenko [B-F-K(1998)], in a semi-dispersive billiard flow there can only be finitely many collisions in finite time, see Theorem 1.1 of [B-F-K(1998)], see also 2.5. Thus the dynamics is well defined, as long as the trajectory does not hit more than one boundary components at the same time.

Lemma 4.1 of [K-S-Sz(1990)] claims that the set

$$\Delta_2 = \left\{x \in \mathbf{M}\text{: } \exists \text{ at least 2 singularities on } S^{\mathbb{R}}x\right\}$$

is a countable union of smooth submanifolds of $\mathbf{M}$ with codimension at least two. Especially, the set Δ_2 is slim, i. e. negligible in our considerations, see also Lemma 2.7.8.

By using Proposition 3.1 and the results of §4–7, we are now going to prove the theorem by an induction on the number of disks N (≥ 2). For $N = 2$ the result is proven by Sinai in [Sin(1970)].

Suppose now that $N > 2$, and the theorem has been proven for every number of disks $N' < N$. Assume, further, that the hard disk system $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ with the outer geometric parameters $(m_1, \ldots, m_N; r)$ has the property that neither this system, nor any of its sub-billiard systems belongs to the exceptional null set featuring Proposition 3.1. Theorem 6.1 of [Sim(1992-A)] states that for almost every phase point $x \in \mathcal{SR}^+$ (with respect to the surface measure on the singularity set $\mathcal{SR}^+$) the positive semi-trajectory $S^{(0,\infty)}x$ contains infinitely many consecutive,

connected collision graphs. Then Corollary 3.24 and Lemma 4.2 of [Sim(2000)] together claim that such non-sufficient phase points $x \in \mathcal{SR}^+$ necessarily belong to a countable family of proper (i. e. of codimension at least one) submanifolds J of the phase space $\partial\mathbf{M}$ of the Poincare section mapping. Finally, Proposition 3.1 above implies that these exceptional submanifolds J have codimension-two (in $\partial\mathbf{M}$) intersection with $\mathcal{SR}^+$ and, therefore, typical phase points $x \in \mathcal{SR}^+$ are sufficient. This precisely means that the so called Chernov-Sinai Ansatz (a necessary condition for the Theorem on Local Ergodicity, see Condition 3.1 in [K-S-Sz(1990)]) holds true.

Quite similarly, Theorem 5.1 of [Sim(1992-A)] together with the slimness of the set Δ_2 of doubly singular phase points assert that there exists a slim subset $S_1 \subset \mathbf{M}$ of the phase space such that for every $x \in \mathbf{M} \setminus S_1$ the phase point x has at most one singularity on its trajectory $S^{\mathbb{R}}x$, and each branch of $S^{\mathbb{R}}x$ contains infinitely many consecutive, connected collision graphs. By Corollary 3.24 and Lemma 4.2 of [Sim(2000)], there exists a locally finite (and, therefore, countable) family of codimension-one, smooth, exceptional submanifolds $J_i \subset \mathbf{M}$ such that for every phase point $x \notin \bigcup_i J_i \cup S_1$ the trajectory $S^{\mathbb{R}}x$ has at most one singularity and it is sufficient. According to the celebrated Theorem on Local Ergodicity by Chernov and Sinai (Theorem 5 in [S-Ch(1987)], see also Corollary 3.12 in [K-S-Sz(1990)]) an open neighbourhood $U_x \ni x$ of such a phase point $x \notin \bigcup_i J_i \cup S_1$ belongs to one ergodic component of the billiard flow $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$, therefore this billiard flow has (at most countably many) open ergodic components $C_1, C_2, \ldots$. We note that theorems 5.1 and 6.1 of [Sim(1992-A)] use the induction hypothesis.

We carry out an indirect proof for the induction step. Assume that, contrary to the assertion of the Theorem, the number of the (open) ergodic components $C_1, C_2, \ldots$ is at least two. The main question is how different ergodic components C_i and C_j can be separated in $\mathbf{M}$?

The above argument showed that in the case of more than one ergodic components C_i there must exist a codimension-one, smooth (actually, analytic) exceptional submanifold $J \subset \mathbf{M} \setminus \partial\mathbf{M}$, a non-singular phase point $x_0 \in J$, and an open ball neighbourhood $B_0 = B_0(x_0, \epsilon_0) \subset \mathbf{M} \setminus \partial\mathbf{M}$ of x_0 with the following properties:

(0) The pair of sets $(B_0, B_0 \cap J)$ is diffeomorphic to the standard pair $(\mathbb{R}^{2d-1}, \mathbb{R}^{2d-2})$, where $2d-1 = \dim\mathbf{M}$, and the two connected components B^1 and B^2 of $B_0 \setminus J$ belong to distinct ergodic components C_i and C_j;

(1) For every $x \in B_0$, the semi-trajectory $S^{[0,\infty)}x$ is sufficient if and only if $x \notin J$;

(2) The dimension $\dim\mathcal{N}_0\left(S^{[0,\infty)}x_0\right)$ (≥ 2) of the neutral space of $S^{[0,\infty)}x_0$ (see §2) has the minimum possible value for all such separating manifolds J and phase points $x \in J$;

(3) There exists a countable union E of proper (i. e. of codimension at least one) submanifolds of J such that for every phase point $x \in J \setminus E$ the positive trajectory $S^{[0,\infty)}x$ is non-singular, and the base point x_0 is selected from the set $J \setminus E$.

We say that J is a "separating manifold". The results of §4–7 assert that such a separating manifold J does not exist. This contradiction finishes the proof of the Theorem. $\square$

§4. Non-Existence of Separating J-Manifolds. Part A: The Neutral Sector Opens Up

As we have seen in §3, the only obstacle on the road of successfully proving (by induction) the ergodicity of almost every hard disk system $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ is the following situation: There exists a codimension-one, smooth (actually, analytic) submanifold $J \subset \mathbf{M} \setminus \partial\mathbf{M}$, a phase point $x_0 \in J$, and an open ball neighborhood $B_0 = B_0(x_0, \epsilon_0) \subset \mathbf{M} \setminus \partial\mathbf{M}$ of x_0 in $\mathbf{M}$ with the following properties:

(0) The pair of sets $(B_0, B_0 \cap J)$ is diffeomorphic to the standard pair $(\mathbb{R}^{2d-1}, \mathbb{R}^{2d-2})$, where $2d-1 = \dim\mathbf{M}$, and the two connected components B^1 and B^2 of $B_0 \setminus J$ belong to distinct ergodic components C_i and C_j;

(1) For every $x \in B_0$ the semi-orbit $S^{(0,\infty)}x$ is sufficient (hyperbolic) if and only if $x \notin J$;

(2) The dimension $\dim\mathcal{N}_0(S^{(0,\infty)}x_0)$ of the neutral space of the semi-orbit $S^{(0,\infty)}x_0$ (see §2) is the minimum possible value for all separating manifolds J and phase points $x \in J$. Then, by the upper semi-continuity of this dimension, we can assume that the neighbourhood B_0 is already small enough to ensure that

$$\dim\mathcal{N}_0(S^{(0,\infty)}x) = \dim\mathcal{N}_0(S^{(0,\infty)}x_0) \text{ for all } x \in J \cap B_0; \tag{4.1}$$

(3) There exists a countable union E of proper (i. e. of codimension at least one) submanifolds of J, such that for every $x \in J \setminus E$ the positive orbit $S^{(0,\infty)}x$ is non-singular, and $x_0 \in J \setminus E$.

We say that J is a separating manifold, or briefly a J-manifold.

The assumed minimality of $\dim\mathcal{N}_0(S^{(0,\infty)}x_0)$ will have profound geometric consequences in the upcoming sections.

First of all, we need to introduce a few notions and notations. Let $w_0 \in \mathcal{N}_0(S^{(0,\infty)}x_0)$ be a unit neutral vector of x_0 with the additional property $\langle w_0, v_0\rangle = 0$, where v_0 is the velocity component of the phase point $x_0 = (q_0, v_0)$. For any pair of real numbers (τ_1, τ_2) ($|\tau_i| < \epsilon_1$, $\epsilon_1 > 0$ is fixed, chosen sufficiently small) we define $T_{\tau_1,\tau_2}x_0$ as the phase point

$$T_{\tau_1,\tau_2}x_0 = \Big(q_0 + \tau_1 w_0,\ (1+\tau_2^2)^{-1/2}(v_0 + \tau_2 w_0)\Big). \tag{4.2}$$

It follows immediately from the properties of the exceptional manifold J that

$$T_{\tau_1,\tau_2}x_0 \in J \cap B_0 \text{ for } |\tau_i| < \epsilon_1, \tag{4.3}$$

as long as the upper bound ϵ_1 is selected small enough.

Basic Properties of $S^{(0,\infty)}x$ for $x = T_{\tau_1,\tau_2}x_0$

We want to investigate the positive semi-trajectories $S^{(0,\infty)}x$ of the phase points $x = T_{\tau_1,\tau_2}x_0 \in J \cap B_0$, $|\tau_i| < \epsilon_1$. The key point in their investigation is that both translations $\tau_1 w_0$ and $\tau_2 w_0$ (one for the configuration, the other for the velocity) are neutral for $S^{\mathbb{R}}x_0$, i. e. they do not cause any velocity change in the velocity history of the semi-trajectory.

Lemma 4.4 (Lemma on the "Neutral Trapezoid"). *Assume that* $|\tau_i| < \epsilon_1$, $\tau_1 \cdot \tau_2 \geq 0$, $t > 0$. *Introduce the notations* $t^* = (1+\tau_2^2)^{-1/2}t$, $x^* = S^{t^*}x_0 = (q_0^*, v_0^*)$, $w_0^* = \left(DS^{t^*}\right)(w_0)$, $\tau_1^* = \tau_1 + t^*\tau_2$. *Then the phase point* $S^t x$ ($x = T_{\tau_1,\tau_2}x_0$) *is equal to*

$$T_{\tau_1^*,\tau_2}x^* = \left(q_0^* + \tau_1^* w_0^*,\, (1+\tau_2^2)^{-1/2}(v_0^* + \tau_2 w_0^*)\right),$$

provided that the so called "neutral trapezoid"

$$\begin{aligned} NT &= NT(x_0, w_0, \tau_1, \tau_2, t) \\ &= \left\{ S^{t'}\left(T_{\tau_1',\tau_2'}x_0\right) :\, |\tau_i'| \leq |\tau_i|,\ \tau_i' \cdot \tau_i \geq 0,\ 0 \leq t' \leq t \right\} \end{aligned} \tag{4.5}$$

is free of singularities, i. e. of multiple or tangential collisions.

Remark 4.6. It should be noted, however, that — when defining the translates

$$T_{s,\tau_2}x^* = \left(q_0^* + s \cdot w_0^*,\, (1+\tau_2^2)^{-1/2}(v_0^* + \tau_2 w_0^*)\right)$$

($0 \leq s \leq \tau_1^*$) — we may hit the boundary of the phase space, which means that the time moment of a collision reaches the value zero. In that case, in order to continue these translations beyond s, by definition, we reflect both the direction vector w_0^* of the spatial variation and the velocity $(1+\tau_2^2)^{-1/2}(v_0^* + \tau_2 w_0^*)$ with respect to the tangent hyperplane of the boundary $\partial\mathbf{Q}$ of the configuration space at the considered point of reflection. Although we use this reflection in our definition of $T_{s,\tau_2}x^*$, in order to keep our notations simpler, we do not indicate the arising change in w_0^* and $v_0^* + \tau_2 w_0^*$ in the formulas. This convention will not cause any confusion in the future.

Proof of Lemma 4.4. The whole point is that — as long as we do not hit any singularity — there is a linear $\mathbb{R}^2$-action $A_{\alpha,\beta}$ ($(\alpha,\beta) \in \mathbb{R}^2$) lurking in the background:

$$\begin{cases} A_{\alpha,\beta}x_0 = S^{\alpha}\left(T_{\beta,0}x_0\right), \\ A_{\alpha,\beta}\left(A_{\alpha',\beta'}x_0\right) = S^{\alpha+\alpha'}\left(T_{\beta+\beta',0}x_0\right) \end{cases} \tag{4.7}$$

acting on the sheet $\left\{A_{\alpha,\beta}x_0 :\, (\alpha,\beta) \in \mathbb{R}^2\right\}$. There is no problem with the neutrality (linearity, i. e. no velocity change) and smoothness of this action $A_{\alpha,\beta}$ as long as we know that the rectangle

$$\{A_{\alpha',\beta'}x_0 :\, |\alpha'| \leq |\alpha|,\ |\beta'| \leq |\beta|, \alpha \cdot \alpha' \geq 0,\ \beta \cdot \beta' \geq 0\}$$

does not hit any singularity. The neutral trapezoid $NT = NT(x_0, w_0, \tau_1, \tau_2, t)$ of (4.5) can be expressed in terms of the $A_{\alpha,\beta}$-action as follows:

(4.8)

$$\begin{aligned} &NT(x_0, w_0, \tau_1, \tau_2, t) \\ &= \{T_{0,\lambda}A_{\alpha,\beta}x_0 :\, 0 \leq \alpha \leq t^*,\ |\beta| \leq |\tau_1| + \alpha|\tau_2|,\ |\lambda| \leq |\tau_2|,\ \lambda \cdot \tau_i \geq 0,\ \beta \cdot \tau_i \geq 0\}. \end{aligned}$$

The statement of Lemma 4.4 is then easily provided by the commutativity and neutrality of the $A_{\alpha,\beta}$-action. □

Let $b > 0$ be a suitably big number so that

$$\dim\mathcal{N}_0\left(S^{[0,b]}x_0\right) = \dim\mathcal{N}_0\left(S^{[0,\infty)}x_0\right). \tag{4.9}$$

(The number b is assumed to be not a moment of collision.) We can further assume that the upper bound $\epsilon_1 > 0$ is already chosen so small that for $|\tau_i| < \epsilon_1$, $\tau_1 \cdot \tau_2 \geq 0$, and for the orbit segment $S^{[0,b]}x$ of $x = T_{\tau_1,\tau_2}x_0$ we have

$$\begin{aligned}\dim\mathcal{N}_0\left(S^{[0,b]}x\right) &= \dim\mathcal{N}_0\left(S^{[0,\infty)}x\right)\\ &\left(= \dim\mathcal{N}_0(S^{[0,\infty)}x_0)\right).\end{aligned} \tag{4.10}$$

Key Lemma 4.11 (Infinite sector opens up in the neutral sheet). *For a typically selected phase point $x_0 \in J$ (more precisely, apart from a first category subset of J) the following holds true:*

For every pair of real numbers (τ_1, τ_2) with $|\tau_i| < \epsilon_1$, $\tau_1 \cdot \tau_2 \geq 0$, the positive trajectory $S^{[0,\infty)}x$ of $x = T_{\tau_1,\tau_2}x_0$ ($\in J \cap B_0$) does not hit any singularity.

Proof. We will argue by the absurd. Suppose that $S^t\left(T_{\tau_1,\tau_2}x_0\right)$ hits a singularity at time moment $t = t_0$ (> 0). Due to the smoothness of the orbit segments $S^{[0,b]}\left(T_{\tau_1,\tau_2}x_0\right)$, the number t_0 is necessarily greater than b. The considered singularity can be one of the following two types:

Type I. Tangential collision between the disks i and j at time $t = t_0$. To simplify the notations, we assume that $\tau_i \geq 0$, $i = 1, 2$. Let us understand the relationship between the curve $\gamma(s) = T_{\tau_1+s,\tau_2}x_0$ ($|s| << 1$) and the invariant hull $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$ of the considered tangential singularity $\mathcal{S}$. Due to the doubly neutral (linear) nature of the perturbations $T_{\tau_1,\tau_2}x_0$, for the parameter values $s < 0$ the disks i and j must avoid each other (pass by each other) for $t \approx t_0$. Otherwise, if these disks collided on the orbit of $\gamma(s)$ near $t = t_0$ for $s < 0$, then the further neutral perturbations $\gamma(s)$ with $s \nearrow 0$ would not set these disks apart at $t = t_0$, thanks to the doubly neutral nature of the perturbations $T_{\tau_1+s,\tau_2}x_0$. Since, by our indirect assumption, for the value $s = 0$ the curve $\gamma(s)$ hits $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$, we have that $\gamma(s)$ is transversal to the manifold $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$ at the point $\gamma(0)$. However, this transversality "kills" the neutral vector $w_0 \in \mathcal{N}_0(S^{[0,\infty)}x) = \mathcal{N}_0(S^{[0,b]}x)$ ($x = T_{\tau_1,\tau_2}x_0 = \gamma(0)$) by lowering the dimension of $\mathcal{N}_0\left(S^{[0,\infty)}\gamma(s)\right)$ for $s > 0$. (We note that new neutral vectors cannot appear because of the stable nature of $\mathcal{N}_0\left(S^{[0,b]}\gamma(s)\right)$.) The latest statement, however, contradicts to the assumed minimality of $\dim\mathcal{N}_0\left(S^{[0,\infty)}x\right) = \dim\mathcal{N}_0\left(S^{[0,\infty)}x_0\right)$, see also (4.1) and the text surrounding it.

Type II. A multiple collision singularity (of type (i,j)–(j,k)) at $t = t_0$.

We begin with an important remark. The multiple collision singularity of type (i,j)–(j,k) means that on each side of the singularity a finite sequence of alternating collisions (i,j) and (j,k) takes place in such a way that on one side of the singularity this finite sequence starts with (i,j), while on the other side it starts with (j,k). Purely to simplify the notations, hereby we are presenting a study of this type of singularity in the case when both collision sequences are made up by

two collisions. This is not a restriction of generality, but merely a simplification of the notations.

Just as above, we again consider the curve $\gamma(s) = T_{\tau_1+s,\tau_2} x_0$ ($|s| << 1$) and its relationship with the invariant hull $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$ of the considered double collision singularity $\mathcal{S}$. Suppose that on the side $s < 0$ of $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$ the collision $(i,j) = \sigma_l$ precedes the collision $(j,k) = \sigma_{l+1}$. Then the derivative of the time difference $t(\sigma_{l+1}) - t(\sigma_l)$ with respect to s at $s = 0$ must be negative, and, therefore, the curve $\gamma(s)$ transversally intersects the invariant hull $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$ of the studied double singularity at the point $\gamma(0)$. More precisely, denote by v_i^-, v_j^-, and v_k^- the velocities of the disks i, j, k right before the collision σ_l on the trajectory of $\gamma(s) = T_{\tau_1+s,\tau_2} x_0$ for $s < 0$, $|s| << 1$. Similarly, let v_i^+, v_j^+, and $v_k^+ = v_k^-$ the corresponding velocities between σ_l and σ_{l+1} on the orbit of the phase point $\gamma(s)$ for $s < 0$, $|s| << 1$, and let

$$\begin{cases} w_0^- = (\delta q_1^-, \dots, \delta q_N^-) = \left(DS^{t(\sigma_l)-\epsilon}(\gamma(s))\right)(w_0), \\ w_0^+ = (\delta q_1^+, \dots, \delta q_N^+) = \left(DS^{t(\sigma_l)+\epsilon}(\gamma(s))\right)(w_0), \end{cases} \tag{4.12}$$

for $s < 0$, $|s| << 1$. By the neutrality of w_0^- and by the conservation of the momentum we immediately obtain

$$\begin{cases} \delta q_i^- - \delta q_j^- = \alpha(v_i^- - v_j^-), \\ \delta q_i^+ - \delta q_j^+ = \alpha(v_i^+ - v_j^+), \\ \delta q_i^+ - \delta q_i^- = \alpha(v_i^+ - v_i^-), \\ \delta q_k^+ = \delta q_k^-, \end{cases} \tag{4.13}$$

where α is the advance of the collision $\sigma_l = (i,j)$ with respect to the neutral vector w_0, see also §2. From the equations (4.13) and from the conservation of the momentum (which is obviously also true for the components δq_a of neutral vectors) we obtain

$$\begin{cases} \delta q_i^+ = \delta q_i^- + \alpha(v_i^+ - v_i^-), \\ \delta q_j^+ = \delta q_j^- + \alpha(v_j^+ - v_j^-), \\ \delta q_j^+ - \delta q_k^+ = \beta(v_j^+ - v_k^+) = \delta q_j^- - \delta q_k^- + \alpha(v_j^+ - v_j^-). \end{cases} \tag{4.14}$$

Here β denotes the advance of the collision $\sigma_{l+1} = (j,k)$ with respect to w_0.

Let us study now the quite similar phenomenon on the other side of the singularity $\bigcup_{t\in\mathbb{R}} S^t(\mathcal{S})$, i. e. for $s > 0$. Since

$$\mathcal{N}_0\left(S^{[0,b]}\gamma(s)\right) = \mathcal{N}_0\left(S^{[0,b]}x_0\right) = \mathcal{N}_0\left(S^{[0,\infty)}x_0\right)$$

and $\mathcal{N}_0\left(S^{[0,\infty)}x_0\right)$ has the minimum value of all such dimensions, we obtain that $w_0 \in \mathcal{N}_0\left(S^{[0,\infty)}\gamma(s)\right)$. Thus, similar thing can be stated about the velocities and neutral vectors as above. Namely, denote by $\tilde{v}_j^+$, $\tilde{v}_k^+$, and $\tilde{v}_i^+ = v_i^-$ the velocities of the disks j, k, i between the collisions $\sigma_l = (j,k)$ and $\sigma_{l+1} = (i,j)$ (Observe that the order of the two collisions is now inverted!) on the orbit $S^{[0,\infty)}\gamma(s)$, $s > 0$, $s << 1$. Let, moreover,

$$\tilde{w}_0^+ = (\delta\tilde{q}_1^+, \dots, \delta\tilde{q}_N^+) = \left(DS^{t(\sigma_l)+\epsilon}(\gamma(s))\right)(w_0) \tag{4.15}$$

for $s > 0$, $s << 1$, and $\tilde{\beta}$, $\tilde{\alpha}$ be the advances of the collisions $\sigma_l = (j,k)$, and $\sigma_{l+1} = (i,j)$, respectively. Then $\delta q_j^- - \delta q_k^- = \tilde{\beta}(v_j^- - v_k^-)$ in the last equation of (4.14), so we get that

$$\beta(v_j^+ - v_k^+) = \tilde{\beta}(v_j^- - v_k^-) + \alpha(v_j^+ - v_j^-). \tag{4.16}$$

By neutrality, for all orbits $S^{[0,\infty)}\gamma(s)$ ($|s| << 1$) the (i,j) collision near $t = t_0$ (which is either σ_l or σ_{l+1}, depending on which side of the singularity we are) has the same normal vector $\vec{n}_1$ and, similarly, for all orbits $S^{[0,\infty)}\gamma(s)$ the (j,k) collision near $t = t_0$ has the same normal vector $\vec{n}_2$. How can we take now advantage of (4.16)? First of all, we can assume that the relative velocities $v_j^- - v_k^-$ and $v_j^+ - v_k^+$ are nonzero, for each of the equations $v_j^- - v_k^- = 0$ and $v_j^+ - v_k^+ = 0$ defines a codimension-two set, which is atypical in J, so we can assume that these vectors are nonzero on the orbit of $\gamma(0)$ (or, equivalently, on the orbit of x_0) by typically choosing the starting phase point x_0. Secondly, by adding an appropriate scalar multiple of v_0 to the neutral vector w_0, we can achieve that $\alpha = 0$ in (4.16), see also §2. We infer, therefore, that the relative velocities $v_j^- - v_k^-$ and $v_j^+ - v_k^+$ are parallel, as long as at least one of the advances β and $\tilde{\beta}$ is nonzero. However, $\alpha = \beta = \tilde{\beta} = 0$ would mean that $\delta q_i^- = \delta q_j^- = \delta q_k^-$, which is impossible, for in that case the time difference $t(\sigma_{l+1}) - t(\sigma_l)$ would not change (and, therefore, it could not tend to zero) as $s \nearrow 0$. Thus, we conclude that $v_j^- - v_k^- \parallel v_j^+ - v_k^+$. However, the difference of these vectors is obviously parallel to the collision normal $\vec{n}_1$, so we get

$$v_j^- - v_k^- \parallel \vec{n}_1. \tag{4.17}$$

A similar argument yields

$$v_i^- - v_j^- \parallel \vec{n}_2. \tag{4.18}$$

However, the events described in (4.17–18) together define a codimension-two subset of the phase space, so we can assume that the typically selected starting phase point $x_0 \in J$ is outside of all such codimension-two submanifolds. This finishes the proof of Main Lemma 4.11. □

§5. Non-Existence of J-Manifolds. Part B: The Weird Behaviour of the ω-limit Set

Let us study now the non-empty, compact ω-limit set

$$\Omega(x_0) = \left\{ x_\infty \in \mathbf{M}\colon \exists \text{ a sequence } t_n \nearrow \infty \text{such that } x_\infty = \lim_{n\to\infty} S^{t_n} x_0 \right\} \tag{5.1}$$

of the trajectory $S^{\mathbb{R}}x_0 = \{S^t x_0 = x_t\colon t \in \mathbb{R}\}$. Consider an arbitrary phase point $x_\infty \in \Omega(x_0)$, $x_\infty = \lim_{n\to\infty} x_{t_n}$, $t_n \nearrow \infty$. Although the trajectory $S^{\mathbb{R}}x_\infty$ of x_∞ may be singular, we can assume that whenever x_∞ belongs to a singularity $S^{-t}\mathcal{S}$, the sequence of points x_{t_n} converges to x_∞ from one side of the codimension-one submanifold $S^{-t}\mathcal{S}$. This can be achieved by using Cantor's diagonal method and

switching to a subsequence of the sequence $t_n \nearrow \infty$. Then for $t \in \mathbb{R}$ the phase points x_{t_n+t} will converge as $n \to \infty$, and we will define the limit $\lim_{n\to\infty} x_{t_n+t}$ as $S^t x_\infty$. In this way we correctly define a trajectory branch $S^{\mathbb{R}} x_\infty$ of the phase point x_∞. As for the concept of trajectory branches, see §2.3.

By switching again – if necessary – to a suitable subsequence of $t_n \nearrow \infty$, we can assume that the unit neutral vectors $w_{t_n} = (DS^{t_n}(x_0))(w_0)$ converge to a (unit) neutral vector $w_\infty \in \mathcal{N}_0(S^{\mathbb{R}} x_\infty)$, which is then necessarily perpendicular to the velocity $v_\infty = v(x_\infty) = (v_1^\infty, \dots, v_N^\infty)$ of x_∞. We write

$$w_\infty = (\delta q_1^\infty, \dots, \delta q_N^\infty), \quad x_\infty = (q_1^\infty, \dots, q_N^\infty; v_1^\infty, \dots, v_N^\infty).$$

We would like to point out again that the well defined orbit $S^{\mathbb{R}} x_\infty$ may be singular. In the case of a multiple collision, according to what was said above, the infinitesimal time-ordering of the collisions (taking place at the same time) is determined, just as the resulting product of reflections connecting the incoming velocity v^- with the outgoing velocity v^+. As far as the other type of singularity — the tangential collisions — is concerned, here there are two possibilities. The first one, in which case the tangentially colliding disks i and j have proper collisions on the nearby approximating trajectories $S^{\mathbb{R}} x_{t_n}$, $n \to \infty$. The second case is when the tangentially colliding disks i and j pass by each other without collision on the approximating orbit $S^{\mathbb{R}} x_{t_n}$, $n \to \infty$. In both cases, we do not include a tangential collision in the symbolic collision sequence of $S^{\mathbb{R}} x_\infty$. In the sequel we will exclusively deal with non-tangential collisions, i. e. collisions with nonzero momentum exchange. They are called proper collisions. This note has particular implications when defining the connected components of the collision graph of the entire trajectory $S^{\mathbb{R}} x_\infty$.

Definition 5.2. Let $\{1, 2, \dots, N\} = H_1 \cup H_2 \cup \cdots \cup H_k$ be the partition of the vertex set into the connected components of the collision graph $\mathcal{G}\left(S^{\mathbb{R}} x_\infty\right)$ of the orbit $S^{\mathbb{R}} x_\infty$. For any i, $1 \le i \le k$, we denote by $\{S_i^t\}_{t\in\mathbb{R}} = \{S_i^t\}$ the internal dynamics of the subsystem H_i, i. e. the dynamics in which we

(a) reduce the total momentum of the subsystem H_i to zero by observing it from a suitably moving reference system;

(b) do not make any distinction between two configurations of H_i differing only by a uniform spatial translation; (Factorizing with respect to uniform spatial translations, see also §1.)

(c) carry out a time-rescaling, so that the total kinetic energy of the internal system $\{S_i^t\}_{t\in\mathbb{R}}$ is equal to 1.

Let, moreover, $M_i = \sum_{j\in H_i} m_j$ the total mass, $I_i = \sum_{j\in H_i} m_j v_j^\infty$ the total momentum, and $V_i = I_i/M_i$ the average velocity of the subsystem H_i. Similarly, we write $W_i = (M_i)^{-1} \sum_{j\in H_i} m_j \delta q_j^\infty = (M_i)^{-1} \sum_{j\in H_i} m_j w_j^\infty$ for the total (average) displacement of the system H_i under the action of the neutral vector $w_\infty = (\delta q_1^\infty, \dots, \delta q_N^\infty) = (w_1^\infty, \dots, w_N^\infty)$. Finally, let $|H_i| \ge 2$ for $i \in \{1, 2, \dots, s\}$, $|H_i| = 1$ for $i \in \{s+1, \dots, k\}$.

First of all, we prove

Lemma 5.3. *Let λ, $\mu \in \mathbb{R}$ be given numbers, and form the neutral vector*

$$n(\mu, \lambda) = \mu v_\infty + \lambda w_\infty \in \mathcal{N}_0(x_\infty) = \mathcal{N}_0(S^{\mathbb{R}} x_\infty).$$

Define $T_{n(\mu,\lambda),0}x_\infty = (q_\infty + n(\mu,\lambda), v_\infty)$ as the neutral translation of $x_\infty = (q_\infty, v_\infty)$ by the vector $n(\mu,\lambda)$, where we use the natural convention of Remark 4.6. Let, finally, i and j be labels of disks belonging to different components H_l, say, to H_1 and H_2. We claim that the orbit $S^{\mathbb{R}}T_{n(\mu,\lambda),0}x_\infty$ of $T_{n(\mu,\lambda),0}x_\infty$ cannot have a proper (i. e. non-tangential) collision between the disks i and j.

Proof. Assume the contrary. Then, by a simple continuity argument, one finds some real numbers μ_0, λ_0 for which the orbit of $T_{n(\mu_0,\lambda_0),0}x_\infty$ hits a tangential singularity between the disks i and j. By using a suitably accurate approximation $(x_{t_n}, w_{t_n}) \approx (x_\infty, w_\infty)$, one finds a neutral, spatial translation of x_{t_n} by a vector $\mu_1 v_{t_n} + \lambda_1 w_{t_n}$ ($\mu_1 \approx \mu_0$, $\lambda_1 \approx \lambda_0$) such that the orbit of $(q_{t_n} + \mu_1 v_{t_n} + \lambda_1 w_{t_n},\, v_{t_n})$ hits a tangential singularity between the disks i and j, which is impossible by Lemma 4.11. This finishes the indirect proof of 5.3. $\square$

The main step in the indirect proof of the Theorem is

Key Lemma 5.4. *There exists a finite collection of nonzero lattice vectors*

$$l_0, l_1, \ldots, l_p \in \mathbb{Z}^2$$

(depending only on N and the common radius r of the N disks moving in the standard unit torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$) with the following properties: For every separating manifold J, for every phase point $x_0 \in J$ fulfilling conditions (0)—(3) listed at the beginning of the previous section, and for every ω-limit point $x_\infty = \lim_{n\to\infty} x_{t_n}$ of the orbit $S^{\mathbb{R}}x_0$ it is true that $k \geq 2$ (i. e. the collision graph $\mathcal{G}$ of $S^{\mathbb{R}}x_\infty$ is not connected, see Definition 5.2 above), and there is an index $j \in \{0, 1, \ldots, p\}$ such that all velocities v_i^∞ $(i = 1, \ldots, N)$ are parallel to the lattice vector l_j.

Remark 5.4/a. It is easy to see that the scenario described in the key lemma (i. e. that all velocities are parallel to l_j for all time $t \in \mathbb{R}$) can only take place if the dynamically connected components of the motion — the connected components of the collision graph of $S^{\mathbb{R}}x_\infty$ — move on closed geodesics of $\mathbb{T}^2$ being parallel to l_j.

Proof. First of all, we prove the geometric

Sublemma 5.5. *Consider the standard x–y coordinate plane with the usual unit vectors $e_1 = (1,0)$, $e_2 = (0,1)$. Suppose that infinitely many disks of radius r and centers at $q_i + je_2 \in \mathbb{R}^2$ $(i = 1, \ldots, N;\ j \in \mathbb{Z})$ are moving uniformly in $\mathbb{R}^2$ and colliding elastically. We assume that the disk centered at $q_i + je_2$ has mass m_i, and its velocity $v_i = \dot{q}_i$ is also independent of j, $i = 1, \ldots, N$, $j \in \mathbb{Z}$. We claim that if the trajectory of such an e_2-periodic system remains in the vertical strip $0 \leq x \leq L$ (for all time $t \in \mathbb{R}$, the number $T > 0$ is given), then all velocities $\dot{q}_i(t) = v_i(t)$ are parallel to e_2.*

Proof of 5.5. We carry out an induction on the number of disks N of the e_2-factorized system. For $N = 1$ the statement is obviously true. Let $N > 1$, and assume that the sublemma has been proven for all numbers $N' < N$.

Let $i_1, i_2, \ldots, i_a$ be the labels i of disks with the largest value of the inner product $\langle q_i, e_1\rangle$ at time $t = 0$. To simplify the notations, we assume that the indices $i_1, i_2, \ldots, i_a$ are just $1, 2, \ldots, a$.

Suppose first that the x-coordinate $\langle v_i, e_1\rangle$ of the velocity $v_i = \dot{q}_i$ is nonzero for some $i \leq a$. By reversing time, if necessary, we can assume that $\langle v_i, e_1\rangle > 0$ for some $i \leq a$. This means that among the disks with the rightmost position at least

one moves to the right. Denote by $i = 1, 2, \dots, b$ $(1 \le b \le a)$ the labels of disks i $(i \le a)$ for which the inner product $\langle v_i, e_1 \rangle$ is maximal. Now it is easy to see that the first velocity component $\langle v_1, e_1 \rangle = \cdots = \langle v_b, e_1 \rangle$ (> 0) cannot decrease in time. As a matter of fact, two things can only happen to the disk(s) i with the rightmost position and maximum value of $\langle v_i, e_1 \rangle$: Either the disk i collides with another disk coming from the left, or another disk j with a larger velocity component $\langle v_j, e_1 \rangle$ passes by q_i, thus by snapping the "title" of having the rightmost position. In either case, the maximum value of the first velocity component $\langle v_i, e_1 \rangle$ of the rightmost disk(s) can only increase. This argument shows that at least one disk i will escape to the right ($\langle q_i, e_1 \rangle \to +\infty$), which is impossible by our assumption on the boundedness of the x-coordinates.

Therefore, only the second possibility can occur, i. e. that $\langle v_i, e_1 \rangle = 0$ for all $i \le a$. This should then remain valid for all time $t \in \mathbb{R}$ by the above argument. However, this also means that the disks $q_i + je_2$, $i \le a$, $j \in \mathbb{Z}$, collide among themselves, while all of them have vertical velocities. In the case $a = N$ we are done, while in the case $a < N$ we can use the induction hypothesis, which says that all velocities v_i, $i > a$, are also vertical. This finishes the proof of 5.5. $\square$

Let us return now to the proof of Key Lemma 5.4. Its proof will be divided into several parts.

1^o **First we prove that $k \ge 2$, i. e. the collision graph $\mathcal{G}$ of $S^{\mathbb{R}}x_\infty$ is not connected.** Assume, on the contrary, the connectedness of $\mathcal{G}$. Let us focus on the limiting neutral vector

$$w_\infty = (\delta q_1^\infty, \dots, \delta q_N^\infty) = (w_1^\infty, \dots, w_N^\infty) = \lim_{n \to \infty} w_{t_n},$$

for which $\sum_{i=1}^N m_i w_i^\infty = 0$, $\sum_{i=1}^N m_i ||w_i^\infty||^2 = 1$, and $\sum_{i=1}^N m_i \langle w_i^\infty, v_i^\infty \rangle = 0$, where $x_\infty = (q_1^\infty, \dots, q_N^\infty; v_1^\infty, \dots, v_N^\infty)$. Let $\Sigma = (\dots, \sigma_{-1}, \sigma_0, \sigma_1, \dots)$ be the symbolic collision sequence of $S^{\mathbb{R}}x_\infty$, and denote by $\alpha_j = \alpha(\sigma_j)$ the advance of the collision σ_j with respect to the neutral vector w_∞, see §2. Since w_∞ is not parallel to $v_\infty = (v_1^\infty, \dots, v_N^\infty)$, by using the assumed connectedness of $\mathcal{G}$ we get that not all advances α_j $(j \in \mathbb{Z})$ are equal, see also Remark 2.2 and Proposition 3.4 of [Sim(1992-B)]. By switching from w_∞ to $-w_\infty$, if necessary, we can assume that there is an index $j \in \mathbb{Z}$ for which $\alpha_j < \alpha_{j+1}$. This means, however, that the translated copy $(q_\infty + \lambda w_\infty, v_\infty)$ of $x_\infty = (q_\infty, v_\infty)$ will hit a double collision singularity $t(\sigma_l) = t(\sigma_{l+1})$ for some value

$$0 < \lambda \le \lambda^* = (\alpha_{j+1} - \alpha_j)^{-1} \cdot (t(\sigma_{j+1}) - t(\sigma_j)) .$$

By considering some well approximating pair $(x_{t_n}, w_{t_n}) \approx (x_\infty, w_\infty)$, we get that some translated copy $(q_{t_n} + \lambda' w_{t_n}, v_{t_n})$ of $x_{t_n} = (q_{t_n}, v_{t_n})$ will hit a double collision singularity for some $\lambda' \approx \lambda$. However, this statement contradicts to Lemma 4.11. Therefore, the collision graph $\mathcal{G}$ of $S^{\mathbb{R}}x_\infty$ is not connected, the number k of the connected components of $\mathcal{G}$ is at least two. $\square$

2^o **Next we prove Key Lemma 5.4 in the case $k = N$, i. e. when no proper collision at all takes place on the trajectory $S^{\mathbb{R}}x_\infty$.** (By the way, this phenomenon can only occur if the maximum lengths $\tau(x_{t_n})$ and $\tau(-x_{t_n})$ of the collision-free path of $x_{t_n} = (q_{t_n}, v_{t_n})$ and $-x_{t_n} = (q_{t_n}, -v_{t_n})$ tend to infinity, as $n \to \infty$.)

First we prove

Sublemma 5.6. *For every pair of labels (i,j) $(1 \le i < j \le N)$ it is true that*

$$\dim span\{v_i^\infty - v_j^\infty,\, w_i^\infty - w_j^\infty\} \le 1.$$

Proof. Assume the opposite, i. e. $\dim \mathrm{span}\{v_i^\infty - v_j^\infty,\, w_i^\infty - w_j^\infty\} = 2$. Then there exists a linear combination $z = \mu v_\infty + \lambda w_\infty$ $(\lambda, \mu \in \mathbb{R})$ of v_∞ and w_∞ with the following properties:

(i) $\mathrm{dist}\{q_i^\infty + \mu v_i^\infty + \lambda w_i^\infty,\, q_j^\infty + \mu v_j^\infty + \lambda w_j^\infty\} = 2r$;
(ii) $v_i^\infty - v_j^\infty \perp q_i^\infty - q_j^\infty + \mu(v_i^\infty - v_j^\infty) + \lambda(w_i^\infty - w_j^\infty)$.

(Observe that here we significantly use the condition $\nu = 2$, i. e. that the dimension of the container torus is two.)
This means that the neutral spatial translate $(q_\infty + z,\, v_\infty)$ of $x_\infty = (q_\infty,\, v_\infty)$ hits a tangential collision singularity between the disks i and j. By using again a suitably good approximation $(x_{t_n}, w_{t_n}) \approx (x_\infty, w_\infty)$, one finds a neutral spatial translate $(q_{t_n} + \mu' v_{t_n} + \lambda' w_{t_n},\, v_{t_n})$ of $x_{t_n} = (q_{t_n},\, v_{t_n})$ $(\lambda' \approx \lambda,\ \mu' \approx \mu)$ which is on the tangential collision singularity between the disks i and j. However, this statement contradicts to Lemma 4.11, by finishing the indirect proof of Sublemma 5.6. □

Without restricting generality, we can assume that all mutually distinct velocities out of $v_1^\infty, \ldots, v_N^\infty$ are $v_1^\infty, \ldots, v_\rho^\infty$. In virtue of $\sum_{i=1}^N m_i v_i^\infty = 0$ and $\sum_{i=1}^N m_i ||v_i^\infty||^2 = 1$, not all velocities are the same, i. e. $\rho \ge 2$.

Sublemma 5.7. *All velocities $v_1^\infty, \ldots, v_N^\infty$ are parallel to each other, that is,*

$$\dim span\{v_1^\infty, \ldots, v_N^\infty\} = 1.$$

Proof. Assume the opposite, i. e. $\dim \mathrm{span}\{v_1^\infty, \ldots, v_N^\infty\} = 2$. (Again an essential use of the condition $\nu = 2$.) Due to the relation $\sum_{i=1}^N m_i v_i^\infty = 0$, the points $v_1^\infty, \ldots, v_N^\infty$ of the plane $\mathbb{R}^2$ do not lie on the same line, not even on a line not passing through the origin. Thus, the mutually distinct points $v_1^\infty, \ldots, v_\rho^\infty$ of $\mathbb{R}^2$ are not collinear. According to the Sylvester-Gallai theorem from combinatorial geometry (see p. 167 of [A-P(1995)] or p. 28 of [K(1948)]) there is a line l_0 in $\mathbb{R}^2$ such that l_0 contains exactly two points, say v_1^∞ and v_2^∞, out of the points $v_1^\infty, \ldots, v_\rho^\infty$. Now consider an arbitrary v_i^∞ which is different from both v_1^∞ and v_2^∞, i. e. for which $v_i^\infty \notin l_0$. By Sublemma 5.6, we have that $w_2^\infty - w_1^\infty = \alpha(v_2^\infty - v_1^\infty)$, $w_i^\infty - w_2^\infty = \beta(v_i^\infty - v_2^\infty)$, and $w_i^\infty - w_1^\infty = \gamma(v_i^\infty - v_1^\infty)$. Therefore,

$$\gamma(v_i^\infty - v_1^\infty) = \gamma(v_i^\infty - v_2^\infty) + \gamma(v_2^\infty - v_1^\infty) = \alpha(v_2^\infty - v_1^\infty) + \beta(v_i^\infty - v_2^\infty),$$

that is,

$$(\alpha - \gamma)(v_2^\infty - v_1^\infty) + (\beta - \gamma)(v_i^\infty - v_2^\infty) = 0.$$

Due to the linear independence of $v_2^\infty - v_1^\infty$ and $v_i^\infty - v_2^\infty$, we get that $\alpha = \beta = \gamma$. Let us replace now the indices 1 and 2 of v_1^∞ and v_2^∞ by some other indices j and l with $v_j^\infty = v_1^\infty$, $v_l^\infty = v_2^\infty$, and repeat the above argument with the triplet of indices (j, l, i). We obtain that

$$w_l^\infty - w_j^\infty = \alpha(v_l^\infty - v_j^\infty) = \alpha(v_2^\infty - v_1^\infty)$$

with the same α as above. Therefore, we get

$$w_i^\infty - w_1^\infty = \alpha(v_i^\infty - v_1^\infty), \quad i = 1, 2, \dots, N. \tag{5.8}$$

Thanks to the conventions $\sum_{i=1}^N m_i w_i^\infty = \sum_{i=1}^N m_i v_i^\infty = 0$, the equations (5.8) can only be fulfilled by the solution $w_i^\infty = \alpha v_i^\infty$ $(i = 1, \dots, N)$, which is impossible, for the vector w_∞ is not parallel to v_∞. This contradiction finishes the indirect proof of Sublemma 5.7. $\square$

Now continue the proof of Key Lemma 5.4 in the case $k = N$. We got that all velocities v_i^∞ in $S^{\mathbb{R}}x_\infty$ are parallel to the same direction vector $0 \neq l \in \mathbb{R}^2$. Since the uniformly moving disks of the orbit $S^{\mathbb{R}}x_\infty$ have no proper collision, we get that

$$\operatorname{dist}\{q_2^\infty - q_1^\infty + t \cdot l,\, 0\} \geq 2r$$

for all $t \in \mathbb{R}$. This means, however, that the direction vector l is parallel to an irreducible (non-divisible) lattice vector $0 \neq l_0 \in \mathbb{Z}^2$, such that $||l_0|| \leq \dfrac{1}{4r}$. There are only finitely many choices for such a lattice vector $l_0 \in \mathbb{Z}^2$. This completes the proof of Key Lemma 5.4 in the case $k = N$. $\square$

3^o The case $s = k$ (≥ 2), i. e. when $|H_i| \geq 2$ for all i, $i = 1, \dots, k$.

Let us study, first of all, the relationship between the subsystems H_1 and H_2 (and their internal dynamics $\{S_1^t\}$, $\{S_2^t\}$) with particular emphasis on their relation to the limiting neutral vector $w_\infty \in \mathcal{N}_0(S^{\mathbb{R}}x_\infty)$. The proof presented in (1^o) yields that the advances of all collisions of $\{S_1^t\}_{t\in\mathbb{R}}$ with respect to w_∞ are equal to the same number α and, similarly, all collisions of the internal flow $\{S_2^t\}_{t\in\mathbb{R}}$ share the same advance β with respect to the neutral vector w_∞. Select and fix an arbitrary real number t_0, and consider the linear combination

$$n(\lambda) = (t_0 - \alpha\lambda)v_\infty + \lambda w_\infty \in \mathcal{N}_0(S^{\mathbb{R}}x_\infty)$$

with variable $\lambda \in \mathbb{R}$. Also consider the corresponding neutral spatial translation

$$x_\infty = (q_\infty,\, v_\infty) \longmapsto T_{n(\lambda),0}x_\infty = (q_\infty + n(\lambda),\, v_\infty)$$

of x_∞ with the natural convention of Remark 4.6. Observe that the neutral translation $T_{n(\lambda),0}$ has the following effect on the internal dynamics $\{S_1^t\}_{t\in\mathbb{R}}$ and $\{S_2^t\}_{t\in\mathbb{R}}$: The advance of the subsystem H_1 is t_0, i. e. the internal time of evolution of H_1 will be the fixed number t_0. On the other hand, the advance of H_2 is obviously $t_0 + \lambda(\beta - \alpha)$. We distinguish between two, quite differently behaving situations:

Case (A): $\alpha \neq \beta$. The internal time of the subsystem H_2 (under the translation $T_{n(\lambda),0}x_\infty$, now $\lambda \in \mathbb{R}$ plays the role of time) changes linearly with λ, it is equal to $t_0 + \lambda(\beta - \alpha)$, while the internal time of H_1 is constantly t_0. How about the relative motion of the non-interacting groups H_1 and H_2? Recall that $V_i = (M_i)^{-1}\sum_{j\in H_i} m_j v_j^\infty$ is the average velocity of the subsystem H_i, while $W_i = (M_i)^{-1}\sum_{j\in H_i} m_j w_j^\infty$ is the average displacement of the subsystem H_i under the translation by the neutral vector $w_\infty \in \mathcal{N}_0(S^{\mathbb{R}}x_\infty)$. (Note that $M_i = \sum_{j\in H_i} m_j$.) The relative position of the subsystem H_1 with respect to H_2 can be measured, for example, by the relative position $q_{j_1}^\infty - q_{j_2}^\infty$ of the disks $j_1 \in H_1$, $j_2 \in H_2$, j_1, j_2

fixed. To simplify the notations, we assume that $j_1 = 1$, $j_2 = 2$. Thus the relative position of the subsystem H_1 with respect to H_2 varies with λ as follows:

$$\begin{aligned} q_1^\infty(\lambda) - q_2^\infty(\lambda) &= q_1^\infty - q_2^\infty + (t_0 - \alpha\lambda)(V_1 - V_2) + \lambda(W_1 - W_2) \\ &= q_1^\infty - q_2^\infty + t_0(V_1 - V_2) + \lambda\left[W_1 - W_2 - \alpha(V_1 - V_2)\right]. \end{aligned} \tag{5.9}$$

Now we would like to paint a global picture (global, that is, in the universal covering space $\mathbb{R}^2$) of the orbit of H_2 under the neutral spatial translations $T_{n(\lambda),0}$, $\lambda \in \mathbb{R}$. Due to the factorization with respect to uniform spatial translations when defining our model (see §1), in order to lift the dynamics from $\mathbb{T}^2$ to its universal covering space $\mathbb{R}^2$ (in a $\mathbb{Z}^2$-periodic manner), it is necessary and sufficient to specify the position of the lifted copy $\bar{q}_1^\infty(\lambda) \in \mathbb{R}^2$ of $q_1^\infty(\lambda) = q_1\left(T_{n(\lambda),0}x_\infty\right)$. We take

$$\bar{q}_1(\lambda) = \bar{q}_1 = \int_0^{t_0} \left(v_1(S^t x_\infty) - V_1\right) dt$$

(independently of λ, so that the "baricenter" of H_1 is unchanged while t_0 is changing later on), since the internal time of the subsystem H_1 is constantly t_0, and we want to describe the motion of H_2 relative to H_1. For $i = 1, 2, \ldots, N$ let the resulting $\mathbb{Z}^2$-periodic lifting to $\mathbb{R}^2$ of $q_i^\infty(\lambda) = q_i\left(T_{n(\lambda),0}x_\infty\right)$ be

$$\bar{q}_i(\lambda) + l, \qquad l \in \mathbb{Z}^2, \tag{5.10}$$

where the lifting $\bar{q}_i(\lambda) \in \mathbb{R}^2$ is selected in such a way that it depends on λ continuously. We point out here that currently the translation parameter λ plays the role of time. Also note that for any $j \in H_1$ we have $\bar{q}_j(\lambda) = \text{const}$ (independent of λ), for $\bar{q}_1(\lambda) = \int_0^{t_0} \left(v_1(S^t x_\infty) - V_1\right) dt$, and the internal time of the subsystem H_1 is not changing by the translations $T_{n(\lambda),0}$. We want to pay special attention to the orbit of points $\bar{q}_j(\lambda) + l \in \mathbb{R}^2$, $j \in H_2$, $l \in \mathbb{Z}^2$. We define the open $2r$-neighborhood $U = U(x_\infty, w_\infty, H_1, H_2, t_0)$ of the set

$$\left\{\bar{q}_j(\lambda) + l\colon\ j \in H_2,\ l \in \mathbb{Z}^2,\ \lambda \in \mathbb{R}\right\}$$

as follows:

$$U = \left\{x \in \mathbb{R}^2\colon\ \exists\, j \in H_2,\ l \in \mathbb{Z}^2,\ \lambda \in \mathbb{R} \text{ s. t. } \operatorname{dist}\left(x,\, \bar{q}_j(\lambda) + l\right) < 2r\right\}. \tag{5.11}$$

According to Lemma 5.3, the points $\bar{q}_j(\lambda) = \bar{q}_j$ $(j \in H_1)$ do not belong to the $\mathbb{Z}^2$-periodic open set U. Let us understand the connected components of the set U. Since the open set U is $\mathbb{Z}^2$-periodic, the $\mathbb{Z}^2$-translations will just permute the connected components of U among themselves. The following lemma essentially uses the $2 - D$ topology of $\mathbb{R}^2$:

Sublemma 5.12. *Let $U_0 \subset U$ be a connected component of the $\mathbb{Z}^2$-periodic open set U of (5.11). Then exactly one of the following possibilities occurs:*

(1) U_0 is bounded;

(2) U_0 is unbounded, l_0-periodic with some lattice vector $0 \neq l_0 \in \mathbb{Z}^2$, and U_0 is bounded in the dirction perpendicular to l_0;

(3) U_0 is $\mathbb{Z}^2$-periodic.

Remark 5.13. In the case (2) all periodicity vectors $l \in \mathbb{Z}^2$ of U_0 are integer multiples of an irreducible lattice vector l_0, which is uniquely determined up to a sign.

Proof. Assume that U_0 is unbounded. Let $y_0, y_1, y_2, \cdots \in U_0$, $\mathrm{dist}(y_n, \partial U_0) \geq r$, $|y_n - y_0| \to \infty$, as $n \to \infty$. By switching to a subsequence, if necessary, we can assume that $p(y_n) \to p(y_\infty)$ $(n \to \infty)$ in $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$, where p: $\mathbb{R}^2 \to \mathbb{T}^2$ is the natural projection. The relation $p(y_n) \to p(y_\infty)$ means that $\mathrm{dist}(y_m - y_n, \mathbb{Z}^2) \to 0$, as $m, n \to \infty$. Let $0 < n < m$ be such integers that there exists a nonzero vector $l_0 \in \mathbb{Z}^2$ for which $\mathrm{dist}(y_m - y_n, l_0) < r/10$. Since the translate $U_0 + l_0 \subset U$ intersects U_0 at the point $y_n + l_0$, it follows from the $\mathbb{Z}^2$-translation invariance of U that $U_0 + l_0 = U_0$, i. e. U_0 is l_0-periodic.

Suppose now that U_0 is unbounded in the direction of $(l_0)^\perp$. Then we can repeat the above argument with another sequence $y_0', y_1', y_2', \cdots \in U_0$ for which, in addition to the above properties of the sequence (y_n), it is also true that the sequence $(y_n' - y_0')/||y_n' - y_0'||$ converges to a (unit) vector perpendicular to l_0. Then the above argument provides another vector of periodicity $l_1 \in \mathbb{Z}^2$ for U_0, so that l_0 and l_1 are not parallel. The periodicity of U_0 with respect to the lattice $\mathcal{L}(l_0, l_1)$ spanned by l_0 and l_1 implies, however, that every $\mathbb{Z}^2$-translate of U_0 intersects U_0 and, therefore, U_0 proves to be indeed $\mathbb{Z}^2$-periodic. This concludes the proof of Sublemma 5.12. $\square$

The next sublemma takes into account that the open set U_0 was defined by a special dynamical system.

Sublemma 5.14. *Out of the three cases listed in 5.12, in fact only one of them, namely (2) can occur.*

Proof.

1. The impossibility of (1): Observe that for every $\bar{q}_j(\lambda)$ $(j \in H_2,\ \lambda \in \mathbb{R})$ there exists a lattice vector $l \in \mathbb{Z}^2$ such that $\bar{q}_j(\lambda) + l \in U_0$. This follows simply from the connectedness of the collision graph of the H_2 subsystem $\{S_2^t\}$. If U_0 were bounded, then there would be a bounded cluster (enclosure) of a billiard dynamics with positive kinetic energy inside U_0, which is impossible for many reasons, for example, by Sublemma 5.5. Thus, U_0 is necessarily unbounded.

2. The impossibility of (3): Assume that U_0 is $\mathbb{Z}^2$-periodic. Then U_0 contains an e_1-periodic, continuous curve γ_1, and an e_2-periodic, continuous curve γ_2, as well. The $\mathbb{Z}^2$-periodic system of curves

$$\bigcup_{m \in \mathbb{Z}} (\gamma_1 + me_2) \cup \bigcup_{m \in \mathbb{Z}} (\gamma_2 + me_1) \subset U_0$$

shows that the connected components of $\mathbb{R}^2 \setminus U_0$ are bounded. (Here we essentially use the $2-D$ topology of $\mathbb{R}^2$.) Therefore, the points $\bar{q}_j(\lambda) = \bar{q}_j(0)$ $(j \in H_1,\ \lambda \in \mathbb{R})$ are enclosed in bounded clusters, for they do not belong to U, see Lemma 5.3. Recall that, as the number t_0 varies, the whole set U and all of its connected components U_0 are moving in $\mathbb{R}^2$ at the velocity $V_2 - V_1$ (as the derivative with respect to t_0 shows), see the term containing t_0 in (5.9). However, for the representatives $\bar{q}_j + l$ $(l \in \mathbb{Z}^2,\ j \in H_1)$ of the H_1-dynamics $\{S_1^t\}$ (now the time parameter is t_0) it is impossible to remain in a uniformly moving, bounded enclosure by Sublemma 5.5, for in that case all velocities v_j^∞ $(j \in H_1)$ would be the same, contradicting to

the fact $|H_1| \geq 2$ and the connectedness of the collision graph of H_1. This proves Sublemma 5.14. $\square$

The joint conclusion of sublemmas 5.14 and 5.5 is that all velocities of the internal flow $\{S_2^t\}$ of H_2 are parallel to the vector of periodicity l_0 of the connected component U_0. Moreover, we constructed the lifting $\bar{q}_i(\lambda) \in \mathbb{R}^2$ in such a way that the l_0-periodic strip U_0 — forbidden zone for the points $\bar{q}_i(\lambda) + \mathbb{Z}^2$ $(i \in H_1)$ — moves at the velocity $V_2 - V_1$, see the term containing t_0 in (5.9). Since the lifting of the H_1-subsystem has no drift (the "baricenter" is not moving when t_0 is changing, see the definition of $\bar{q}_1(\lambda) = \bar{q}_1$ above), we get that the relative velocity $V_2 - V_1$ must also be parallel to l_0. This also means that the $\mathbb{R}^2$-lifting of the internal flow $\{S_1^t\}$ is confined to an l_0-periodic, infinite strip bounded by two translated copies of U_0. By using Sublemma 5.5 again, we obtain that all velocities of the internal flow $\{S_1^t\}$ are also parallel to the vector of periodicity l_0. $\square$

Remark 5.14/a. Since $n(\lambda) = (t_0 - \alpha\lambda)v_\infty + \lambda w_\infty$, the drift (i. e. the average derivative of the positions with respect to the variable λ)

$$(M_2)^{-1} \cdot \sum_{j \in H_2} m_j \frac{d}{d\lambda} \bar{q}_j(\lambda)$$

of the subsystem H_2 is equal to $(W_2 - W_1) - \alpha(V_2 - V_1)$ where, as we recall,

$$V_i = (M_i)^{-1} \cdot \sum_{j \in H_i} m_j v_j^\infty, \quad W_i = (M_i)^{-1} \cdot \sum_{j \in H_i} m_j w_j^\infty.$$

Obviously, this drift must be parallel to the vector of periodicity l_0. Since $V_2 - V_1$ is parallel to l_0, we conclude that $W_2 - W_1$ is also parallel to l_0. This remark will be used later in this section.

The second major case in (3^o) is

Case (B): $\alpha = \beta$**.** Let us consider now the modified neutral vector $w_\infty - \alpha v_\infty \in \mathcal{N}_0(S^{\mathbb{R}} x_\infty)$. The advance of both subsystems H_1 and H_2 is zero with respect to the neutral vector $w_\infty - \alpha v_\infty$, thus

$$\begin{aligned} w_j^\infty - \alpha v_j^\infty &= h_1, \quad \forall\, j \in H_1, \\ w_j^\infty - \alpha v_j^\infty &= h_2, \quad \forall\, j \in H_2, \end{aligned} \tag{5.15}$$

for some vectors h_1, $h_2 \in \mathbb{R}^2$. In other words, the effect of the neutral translation by the vector $w_\infty - \alpha v_\infty$ on the non-interacting groups H_1 and H_2 is that H_i gets displaced (translated) by the vector h_i, $i = 1, 2$. Now there are again two subcases:

Subcase B/1: $h_1 \neq h_2$**.** In this case the result of the neutral translation by the vector $n(\lambda) = \lambda(w_\infty - \alpha v_\infty)$ ($\lambda \in \mathbb{R}$ is now the varying parameter) is that the relative translation of the H_2 subsystem with respect to H_1 is $\lambda(h_2 - h_1)$ with the velocity $0 \neq h_2 - h_1 \in \mathbb{R}^2$. The point is that Lemma 5.3 is again readily applicable (so that $n(\mu, \lambda)$ is replaced by $n(\lambda)$), meaning that on the orbit $S^{\mathbb{R}} T_{n(\lambda),0} x_\infty$ of $T_{n(\lambda),0} x_\infty$ no proper collision takes place between the groups H_1 and H_2. This fact has the following consequence on the $\mathbb{Z}^2$-periodic, $\mathbb{R}^2$-lifting

$$\left\{\bar{q}_i(t) + l \in \mathbb{R}^2 \colon\ i \in H_1 \cup H_2,\ l \in \mathbb{Z}^2,\ t \in \mathbb{R}\right\} \tag{5.16}$$

of the subsystem $H_1 \cup H_2$ with the bericenter normalization $\sum_{i\in H_1} m_i \frac{d}{dt}\bar{q}_i(t) = 0$:

$$\text{dist}\,(\bar{q}_i(t),\, \bar{q}_j(t) + \lambda(h_2 - h_1) + l) \geq 2r, \tag{5.17}$$

for $i \in H_1$, $j \in H_2$, $t, \lambda \in \mathbb{R}$, $l \in \mathbb{Z}^2$. In other words, the $2r$-wide, infinite strips with the direction of $h_2 - h_1$ containing $\bar{q}_i(t)$ on their medium line ($i \in H_1$) are disjoint from the similarly constructed infinite strips containing $\bar{q}_j(t)$ ($j \in H_2$) on their medium line. Similarly to the closing part of the discussion of Case (A), we conclude, first of all, that the relative motion (drift) $V_2 - V_1$ between H_2 and H_1 must be parallel to $h_2 - h_1$ and then, according to Sublemma 5.5, all velocities of the internal dynamics $\{S_1^t\}$ and $\{S_2^t\}$ must also be parallel to $h_2 - h_1$. Since the $(h_2 - h_1)$-parallel strips of width $2r$ are disjoint modulo $\mathbb{Z}^2$, we immediately get that $h_2 - h_1$ has a lattice direction, and the shortest nonzero lattice vector l_0 parallel to $h_2 - h_1$ has length at most $1/(4r)$. $\square$

Remark 5.17/a. Let us observe that everything that has been said about the pair $(H_1,\, H_2)$ in Case B/1 can be repeated almost word-by-word if one of the groups H_i, say H_2, has only one element. This remark will have a particular relevance later in this section.

Subcase B/2: $h_1 = h_2$. In this situation the united subsystem $H_1 \cup H_2$ gets uniformly translated by the vector $h_1 = h_2$ under the action of the neutral vector $w_\infty - \alpha v_\infty$. This is an open possibility, indeed, and nothing else can be said about it.

Now we are in the position of quickly finishing the discussion of (3^o). Recall that $s = k$ (≥ 2), i. e. $|H_i| \geq 2$ for $i = 1, \ldots, k$. Consider the advances $\alpha_i = \alpha(H_i)$ of the subsystems $H_1, \ldots, H_k$ with respect to the limiting neutral vector $w_\infty = \lim_{n\to\infty} w_{t_n} \in \mathcal{N}_0(S^{\mathbb{R}} x_\infty)$. Unfortunately, we again have to distinguish between two cases.

Case I. Not all α_i's are the same, e. g. $\alpha_1 \neq \alpha_2$. In this situation the result of Case (A) above says that $V_1 - V_2$ and all velocities of the internal flows $\{S_1^t\}$ and $\{S_2^t\}$ are parallel to the same nonzero lattice vector $l_0 \in \mathbb{Z}^2$. For any other subsystem H_i $(i > 2)$ we have that $\alpha_i \neq \alpha_1$ or $\alpha_i \neq \alpha_2$. Assume that $\alpha_i \neq \alpha_1$. The result of Case (A) above says that $V_i - V_1$ and all velocities of the internal flows $\{S_1^t\}$ and $\{S_i^t\}$ are parallel to the same nonzero lattice vector $l_1 \in \mathbb{Z}^2$. The common presence of the flow $\{S_1^t\}$ in these statements shows that $l_0 = l_1$ (or, at least they are parallel to each other). Summarizing these results, we finish the discussion of Case I by concluding that all average velocities V_i and all velocities of the internal flows $\{S_i^t\}$ $(i = 1, \ldots, k)$ are parallel to the same (nonzero) lattice vector l_0 whose magnitude is at most $1/(4r)$. $\square$

Case II. $\alpha_1 = \alpha_2 = \cdots = \alpha_k =: \alpha$. Consider, as in Case (B) above, the neutral vector $w_\infty - \alpha v_\infty \in \mathcal{N}_0(S^{\mathbb{R}} x_\infty)$. The advance of each H_i with respect to $w_\infty - \alpha v_\infty$ is zero, so (5.15) applies:

$$w_j^\infty - \alpha v_j^\infty = h_i \text{ for } j \in H_i,\ i = 1, 2, \ldots, k. \tag{5.18}$$

Since $w_\infty - \alpha v_\infty \neq 0$ and $\sum_{i=1}^k M_i h_i = 0$ $(M_i = \sum_{j\in H_i} m_j)$, we conclude that not all vectors $h_1, \ldots, h_k \in \mathbb{R}^2$ are the same, e. g. $h_1 \neq h_2$. Then for every $i > 2$ we

have either $h_i \neq h_1$, or $h_i \neq h_2$, say $h_i \neq h_1$, and the result of Case B/1 above applies to the pairs of subsystems (H_1, H_2) and (H_1, H_i). Quite similarly to the discussion of Case I above (but referring in it to Case B/1, instead of Case (A)) we get that all average velocities $V_1, \ldots, V_k$ and all velocities of the internal flows $\{S_i^t\}$ $(i = 1, \ldots, k)$ are parallel to the vector $h_2 - h_1 \neq 0$. Recall that $h_2 - h_1$ has a lattice direction, and the shortest (nonzero) lattice vector l_0 parallel to $h_2 - h_1$ has magnitude at most $1/(4r)$, thus completing the proof of Key Lemma 5.4 in the case (3^o). $\square$

4^o **The general case** $1 \leq s < k$**.** In this case the subsystems H_i with $|H_i| \geq 2$ (i. e. $i \leq s$) coexist with the subsystems H_i for which $|H_i| = 1$ $(i > s)$. In order to simplify the notations we assume that $H_i = \{i\}$ for $i = s+1, s+2, \ldots, k$.

Consider the advances $\alpha_j = \alpha(H_j)$ of the subsystems H_j, $1 \leq j \leq s$, with respect to the limiting neutral vector w_∞. Unfortunately, we again have to distinguish between three major situations.

Case I. Not all of $\alpha_1, \ldots, \alpha_s$ **are equal, say,** $\alpha_1 \neq \alpha_2$**.** Let us observe, first of all, that the whole machinery of (3^o) applies to the united subsystem $H_1 \cup H_2 \cup \cdots \cup H_s$, showing that there exists a nozero lattice vector $l_0 \in \mathbb{Z}^2$ so that all velocities of the internal dynamics $\{S_i^t\}$ $(1 \leq i \leq s)$ and all relative velocities of the baricenters $V_i - V_j$ $(1 \leq i, j \leq s)$ are parallel to l_0. (The second part of this statement is clearly equivalent to saying that the average velocities V_i' of the subsystems H_i $(1 \leq i \leq s)$ are parallel to l_0, provided that these average velocities are observed from a reference system attached to the baricenter of $H_1 \cup H_2 \cup \cdots \cup H_s$.)

Let us turn our attention to a one-disk subsystem $H_i = \{i\}$, $s+1 \leq i \leq k$, i is fixed. Just like in (5.18), the advance of the subsystem H_j $(j \leq s)$ with respect to the neutral vector $w_\infty - \alpha_j v_\infty$ is zero, therefore the whole subsystem H_j gets translated by the same vector $h_j \in \mathbb{R}^2$ under the action of $w_\infty - \alpha_j v_\infty$:

$$w_l^\infty - \alpha_j v_l^\infty = h_j, \quad l \in H_j,\ j = 1, \ldots, s. \tag{5.19}$$

Consider now the vectors of displacement $w_i^\infty - \alpha_j v_i^\infty = h_j' \in \mathbb{R}^2$, $j = 1, \ldots, s$, the index i is fixed, $s+1 \leq i \leq k$. If $h_j - h_j' \neq 0$ for at least one $j \leq s$, then the result of Case B/1 of (3^o) applies to the pair of subsystems (H_j, H_i) (see Remark 5.17/a), thus we have that the relative velocity $V_j - V_i$ of the baricenters is parallel to the fixed lattice vector $l_0 \in \mathbb{Z}^2$. This is the most we can prove for the motion of H_i relative to the motion of $H_1 \cup H_2 \cup \cdots \cup H_s$, for if we had such a result for every i $(s+1 \leq i \leq k)$, then the statement of the key lemma would follow.

The unpleasant situation with H_i is when

$$w_i^\infty - \alpha_j v_i^\infty = h_j = w_l^\infty - \alpha_j v_l^\infty,\ l \in H_j,\ j = 1, \ldots, s, \tag{5.20}$$

$s+1 \leq i \leq k$, i is fixed.

With i and j fixed, let us average (5.20) with respect to the weights m_l $(l \in H_j)$ of the subsystem H_j. We obtain

$$W_j = W_i + \alpha_j (V_j - V_i), \quad j = 1, \ldots, s. \tag{5.21}$$

$s+1 \leq i \leq k$, i is fixed. Recall that $V_i = v_i^\infty$ and $W_i = w_i^\infty$ for the one-disk subsystem $H_i = \{i\}$.

We again have to distinguish between two subcases.

Case I/a. Not all average velocities $V_1, \dots, V_s$ are the same.

Sublemma 5.22. *There is a pair of indices $1 \le j_1, j_2 \le s$ for which $\alpha_{j_1} \ne \alpha_{j_2}$ and $V_{j_1} \ne V_{j_2}$.*

Proof. As a matter of fact, this sublemma is trivial. Indeed, if V_{j_1} were equal to V_{j_2} whenever $\alpha_{j_1} \ne \alpha_{j_2}$ $(1 \le j_1, j_2 \le s)$, then we would have, first of all, $V_1 = V_2$, since $\alpha_1 \ne \alpha_2$ by the assumption of Case I. Secondly, for every $j = 3, 4, \dots, s$ either $\alpha_j \ne \alpha_1$ or $\alpha_j \ne \alpha_2$, thus proving $V_j = V_1 = V_2$ for $j = 3, 4, \dots, s$, contradicting to the assumption of I/a. $\square$

By taking the difference of (5.21) for j_1 and j_2, and also using Remark 5.14 for the pair of subsystems (H_{j_1}, H_{j_2}), we get

$$\begin{aligned} c(V_{j_1} - V_{j_2}) = W_{j_1} - W_{j_2} &= \alpha_{j_1}(V_{j_1} - V_i) - \alpha_{j_2}(V_{j_2} - V_i) \\ &= \alpha_{j_1}(V_{j_1} - V_{j_2}) + (\alpha_{j_1} - \alpha_{j_2})(V_{j_2} - V_i), \end{aligned} \tag{5.23}$$

for some scalar c. Since $V_{j_1} - V_{j_2} \parallel l_0$ and $\alpha_{j_1} - \alpha_{j_2} \ne 0$, we obtain that $V_{j_2} - V_i$ is also parallel to the lattice vector l_0, precisely what we wanted to prove in Case I.

Case I/b. $V_1 = V_2 = \cdots = V_s =: V$**.** Now formula (5.21) says that

$$W_j = W_i + \alpha_j(V - V_i), \quad j = 1, \dots, s, \tag{5.24}$$

the index i is fixed, $s+1 \le i \le k$. Take the difference of (5.24) for $j = 1$ and $j = 2$:

$$W_1 - W_2 = (\alpha_1 - \alpha_2) \cdot (V - V_i). \tag{5.25}$$

Recall that $\alpha_1 \ne \alpha_2$ and, by Remark 5.14/a, $W_1 - W_2$ is parallel to l_0. Therefore, the relative velocity $V_i - V$ also proves to be parallel to the lattice vector l_0, the result we just wanted to prove for the subsystem H_i in Case I. Thus Key Lemma 5.4 has been proven in Case I of (4^o).

Case II. $\alpha_1 = \alpha_2 = \cdots = \alpha_s =: \alpha$**, but not all vectors $h_1, \dots, h_s$ in (5.19) are the same.** Assume that $h_1 \ne h_2$. Then the method of Subcase B/1 of (3^o) applies to the subsystem $H_1 \cup \cdots \cup H_s$, showing again that there exists a nonzero lattice vector l_0 such that all velocities of the flows $\{S_i^t\}$ $(1 \le i \le s)$ and all relative velocities $V_{j_1} - V_{j_2}$ $(1 \le j_1, j_2 \le s)$ are parallel to l_0. Just as in Case I above, consider again a one-disk subsystem $H_i = \{i\}$, $s+1 \le i \le k$. We need to show that some (or any) of the velocities $V_i - V_j$ $(j = 1, \dots, s)$ is parallel to l_0. Consider the vector $h_i = w_i^\infty - \alpha v_i^\infty \in \mathbb{R}^2$. This vector should differ from h_1 or h_2. Assume that $h_i \ne h_1$. In this situation the method and result of Subcase B/1 of (3^o) again applies to the pair of subsystems (H_1, H_i) (see Remark 5.17/a), and we obtain that $V_i - V_1$ is parallel to l_0. This step finishes the proof of Key Lemma 5.4 in Case II of (4^o). $\square$

Case III. $\alpha_1 = \alpha_2 = \cdots = \alpha_s =: \alpha$**, and** $h_1 = h_2 = \cdots = h_s =: h$ **in (5.19).** We can assume that $h_1 = \cdots = h_t = h$ and $h_i \ne h$ for $t+1 \le i \le k$. Due to the relation $\sum_{i=1}^k M_i h_i = 0$ (and to the fact that $w_\infty \ne \alpha v_\infty$), we have that $s \le t \le k-1$. Select and fix an arbitrary index $i \in \{t+1, \dots, k\}$, and study the relative motion of the subsystems $H^* =: H_1 \cup H_2 \cup \cdots \cup H_t$ and $H_i = \{i\}$. Since the neutral vector $w_\infty - \alpha v_\infty$ translates the whole subsystem H^* by the same vector h and it translates

the one-disk subsystem $H_i = \{i\}$ by a different vector h_i, the method and result of Subcase B/1 of (3^o) again applies to the pair (H^*, H_i), and we obtain that there exists a nonzero lattice vector $l_0 \in \mathbb{Z}^2$ so that all velocities of the internal dynamics $\{S_j^t\}$ $(j = 1, \dots, s)$, all relative velocities $V_{j_1} - V_{j_2}$ $(j_1, j_2 \in \{1, \dots, t; i\})$, and $h_i - h$ are parallel to l_0. Due to the common presence of the internal dynamics $\{S_1^t\}$, the same thing can be said about any other index $i \in \{t+1, \dots, k\}$ with the same direction vector l_0. This finishes the proof of Key Lemma 5.4 in the last remaining case of (4^o), thus completing the proof of 5.4. We note that every nonzero lattice vector $l_0 \in \mathbb{Z}^2$ that emerged in this proof had the property $||l_0|| \leq 1/(4r)$, thus ensuring the finiteness of the family $\{l_0, l_1, \dots, l_p\}$ in Key Lemma 5.4. $\square$

§6. Non-Existence of J-Manifolds
Part C: The Continuity Argument

The goal of this brief section is to prove the following, strengthened version of Key Lemma 5.4:

Proposition 6.1. *Given a separating manifold J and a phase point $x_0 \in J$ fulfilling all conditions (0)—(3) listed at the beginning of §4, there exists a lattice vector $l_0 \in \mathbb{Z}^2$ $(0 < ||l_0|| \leq \frac{1}{4r})$ such that for every ω-limit phase point $x_\infty = \lim_{n\to\infty} x_{t_n}$ $(t_n \nearrow \infty)$ of x_0 the common velocity direction vector of x_∞ (as described in the statement of Key Lemma 5.4) is the same vector l_0.*

Proof. According to Key Lemma 5.4, the non-empty, compact ω-limit set $\Omega(x_0)$ of x_0 is partitioned into the closed subsets $\Omega(x_0) = C_0 \cup C_1 \cup \dots \cup C_p$ according to the arising (mutually non-parallel) direction vectors $l_0, l_1, \dots, l_p \in \mathbb{Z}^2$. The orbit $S^{\mathbb{R}}x_0$ of x_0 converges to the set $\Omega(x_0)$: $\operatorname{dist}(S^t x_0, \Omega(x_0)) \to 0$, as $t \to \infty$. Taking into account Remark 5.4/a, we see that the components $C_0, \dots, C_p$ of $\Omega(x_0)$ are stable in the following, weak sense: For every $\epsilon_0 > 0$ there exists another $\epsilon_1 > 0$ such that for every $x \in \mathbf{M}$ with $\operatorname{dist}(x, C_j) < \epsilon_1$ $(j = 0, \dots, p)$, and for every $t \in [0, 1]$ we have that $\operatorname{dist}(S^t x, C_j) < \epsilon_0$.

Select an $\epsilon_0 > 0$ with the property that the open ϵ_0-neighbourhoods $U_{\epsilon_0}(C_0)$,..., $U_{\epsilon_0}(C_p)$ of $C_0, \dots, C_p$ are mutually disjoint. Take $\epsilon_1 > 0$ $(\epsilon_1 < \epsilon_0)$ as above, and observe that $\operatorname{dist}(S^t x_0, \Omega(x_0)) < \epsilon_1$ for $t \geq t_0$. For every $t \geq t_0$ there exists a unique index $i = i(t) \in \{0, 1, \dots, p\}$ such that $S^t x_0 \in U_{\epsilon_1}(C_{i(t)})$. By the above stability one has that $i(s) = i(t)$ for $t \leq s \leq t + 1$, thus showing that $i(t) = \text{const} = i_0$ for all $t \geq t_0$, i. e. it is enough to take the single lattice vector l_{i_0} in Key Lemma 5.4. This proves the proposition. $\square$

§7. Non-Existence of J-Manifolds
Part D: The Instability of the ω-limit Set

Given any nonzero lattice vector $l_0 \in \mathbb{Z}^2$, consider the $1 - D$ subtorus $T(l_0) = \{\lambda l_0 \colon \lambda \in \mathbb{R}\}/\mathbb{Z}^2$ of $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$, and define the following, compact subset $L(l_0)$ of

the phase space $\mathbf{M}$:

$$
\begin{aligned}
L(l_0) = \Big\{ & x \in \mathbf{M}\colon\ v_i(x) \parallel l_0 \text{ for } i = 1, \dots, N; \\
& \exists \text{ a partition } \{1, \dots, N\} = H_1 \cup H_2 \cup \cdots \cup H_k \\
& \text{and elements } a_i \in \mathbb{T}^2,\ i = 1, \dots, k, \text{ such that } q_j(x) \in T(l_0) + a_i \\
& \text{for } j \in H_i,\ i = 1, \dots, k, \text{ and the open } r\text{-neighbourhoods of} \\
& \text{the cosets } T(l_0) + a_i,\ i = 1, \dots, k \text{ are mutually disjoint} \Big\}.
\end{aligned} \tag{7.1}
$$

Plainly, the set $L(l_0)$ is S^t-invariant for $t \in \mathbb{R}$. As a consequence of Key Lemma 5.4, Remark 5.4/a, and Proposition 6.1, we have that for any exceptional manifold $J \subset \mathbf{M}$, any typical phase point $x_0 \in J$ (for which $S^{(0,\infty)}x_0$ is non-singular) there exists a lattice vector $l_0 \in \mathbb{Z}^2$ $(0 < ||l_0|| \le \frac{1}{4r})$ such that the ω-limit set $\Omega(x_0)$ is a subset of $L(l_0)$.

We claim that the set $L(l_0)$ is unstable in the following sense:

Proposition 7.2. *Given a nonzero lattice vector $l_0 \in \mathbb{Z}^2$, there exists a suitably small threshold $\epsilon_0 > 0$ such that for every $0 < \epsilon < \epsilon_0$ and for every phase point $z \notin L(l_0)$, with non-singular positive orbit $S^{[0,\infty)}z$, there exists a number $t > 0$ for which $S^t z \notin U_\epsilon(L(l_0))$, where $U_\epsilon(L(l_0))$ denotes the ϵ-neighbourhood of the compact set $L(l_0)$.*

Proof. Given a phase point $x \in L(l_0)$ and a sufficiently small number $\epsilon > 0$, there exists an open and connected neighbourhood $V \subset U_\epsilon(x)$ of x in $\mathbf{M}$ with the following properties:

(i) $V \cong V_1 \times V_2$, where V_1 is an open neighbourhood of x in $L(l_0)$;
(ii) For any $y \in V_1$ the tangent space $\mathcal{T}_y(\{y\} \times V_2)$ of the leaf $\{y\} \times V_2$ is

$$
\{(\delta q, \delta v) \in \mathcal{T}_y\mathbf{M}\colon\ \delta q_i \perp l_0,\ \delta v_i \perp l_0 \text{ for } i = 1, \dots, N\}\,. \tag{7.3}
$$

The billiard flow $\{S^t\}_{t\in\mathbb{R}}$ is easily seen to be uniformly expanding on the leaves $\{y\} \times V_2$ $(y \in V_1)$, provided that the upper bound ϵ of V $(V \subset U_\epsilon(x))$ is small enough, i. e. $\epsilon < \epsilon_0$ for some threshold $\epsilon_0 > 0$. (The threshold ϵ_0 can be chosen uniformly for all $x \in L(l_0)$.) Therefore, if ϵ_0 is selected small enough (uniformly for all $x \in L(l_0)$), then for any $z \in \{y\} \times V_2$ with non-singular positive orbit, for $y \in V_1$ and $z \notin L(l_0)$, there exists a number $t > 0$ such that $\operatorname{dist}(S^t z,\, S^t y) > \epsilon$ and, what is more, $\operatorname{dist}(S^t z,\, L(l_0)) > \epsilon$. This finishes the proof of Proposition 7.2. □

Finishing the Indirect Proof of the Non-Existence of J-Manifolds

Assume the existence of a J-manifold $J \subset \mathbf{M}$. According to Key Lemma 5.4, Remark 5.4/a, and Proposition 6.1, there exists a lattice vector $l_0 \in \mathbb{Z}^2$ $(0 < \|l_0\| \le \frac{1}{4r})$ such that

$$
\mu_J\left(\left\{x \in J\colon\ \lim_{t\to\infty} \operatorname{dist}(S^t x, L(l_0)) = 0\right\}\right) > 0. \tag{7.4}
$$

(Here $\mu_J(\,.\,)$ denotes the hypersurface measure on J induced by the restriction of the Riemannian metric on the phase space $\mathbf{M}$.) Take $\epsilon_0 > 0$, $0 < \epsilon < \epsilon_0$, and the

open neighbourhood $U_\epsilon(L(l_0))$ of $L(l_0)$ provided by Proposition 7.2. It follows from (7.4) that there exists a suitably big threshold $t_0 > 0$ such that

$$\mu_J\left(\left\{x \in J\colon \operatorname{dist}(S^t x, L(l_0)) < \epsilon \text{ for } t \geq t_0\right\}\right) > 0. \tag{7.5}$$

Set

$$A(l_0, \epsilon, t_0) := \left\{x \in J\colon \operatorname{dist}(S^t x, L(l_0)) < \epsilon \text{ for } t \geq t_0\right\}. \tag{7.6}$$

The image $S^{t_0}\left(A(l_0, \epsilon, t_0)\right)$ contains a positively measured subset of a codimension-one submanifold of $\mathbf{M}$. However, due to Proposition 7.2, we have that

$$S^{t_0}\left(A(l_0, \epsilon, t_0)\right) \subset L(l_0),$$

and the set $L(l_0)$ obviously has at least $N - 1$ (≥ 2) codimensions. These two assertions clearly contradict to each other. The obtained contradiction proves that exceptional manifolds J, as described at the beginning of §4, do not exist. □

§8. Concluding Remarks

8.1 Irrational Mass Ratio.

Due to the natural reduction $\sum_{i=1}^N m_i v_i = 0$ (which we always assume), in §1 we had to factorize out the configuration space with respect to spatial translations: $(q_1, \ldots, q_N) \sim (q_1 + a, \ldots, q_N + a)$ for all $a \in \mathbb{T}^2$. It is a remarkable fact, however, that (despite the reduction $\sum_{i=1}^N m_i v_i = 0$) even without this translation factorization the system still retains the Bernoulli mixing property, provided that the masses $m_1, \ldots, m_N$ are rationally independent. (We note that dropping the above mentioned configuration factorization obviously introduces 2 zero Lyapunov exponents.) For the case $N = 2$ (i. e. two disks) this was proven in [S-W(1989)] by successfully applying D. Rudolph's theorem on the B-property of isometric group extensions of Bernoulli shifts [R(1978)].

Suppose that we are given a dynamical system (M, T, μ) with a probability measure μ and an automorphism T. Assume that a compact metric group G is also given with the normalized Haar measure λ and left invariant metric ρ. Finally, let $\varphi\colon M \to G$ be a measurable map. Consider the skew product dynamical system $(M \times G, S, \mu \times \lambda)$ with $S(x, g) = (Tx, \varphi(x) \cdot g)$, $x \in M$, $g \in G$. We call the system $(M \times G, S, \mu \times \lambda)$ an isometric group extension of the base (or factor) (M, T, μ). (The phrase "isometric" comes from the fact that the left translations $\varphi(x) \cdot g$ are isometries of the group G.) Rudolph's mentioned theorem claims that the isometric group extension $(M \times G, S, \mu \times \lambda)$ enjoys the B-mixing property as long as it is at least weakly mixing and the factor system (M, T, μ) is a B-mixing system.

But how do we apply this theorem to show that the system of N hard disks on $\mathbb{T}^2$ with $\sum_{i=1}^N m_i v_i = 0$ is a Bernoulli flow, even if we do not make the factorization (of the configuration space) with respect to spatial translations? It is simple. The base system (M, T, μ) of the isometric group extension $(M \times G, S, \mu \times \lambda)$ will be the time-one map of the factorized (with respect to spatial translations) hard disk

system. The group G will be just the container torus $\mathbb{T}^2$ with its standard Euclidean metric ρ and normalized Haar measure λ. The second component g of a phase point $y = (x, g) \in M \times G$ will be just the position of the center of the (say) first disk in $\mathbb{T}^\nu$. Finally, the governing translation $\varphi(x) \in \mathbb{T}^\nu$ is quite naturally the total displacement

$$\int_0^1 v_1(x_t)dt \qquad (\text{mod } \mathbb{Z}^2)$$

of the first particle while unity of time elapses. In the previous sections the B-mixing property of the factor map (M, T, μ) has been proven successfully for typical geometric parameters $(m_1, \ldots, m_N; r)$. Then the key step in proving the B-property of the isometric group extension $(M \times G, S, \mu \times \lambda)$ is to show that the latter system is weakly mixing. This is just the essential contents of the article [S-W(1989)], and it takes advantage of the assumption of rational independence of the masses. Here we are only presenting to the reader the outline of that proof. As a matter of fact, we not only proved the weak mixing property of the extension $(M \times G, S, \mu \times \lambda)$, but we showed that this system has in fact the K-mixing property by proving that the Pinsker partition π of $(M \times G, S, \mu \times \lambda)$ is trivial. (The Pinsker partition is, by definition, the finest measurable partition of the dynamical system with respect to which the factor system has zero metric entropy. A dynamical system is K-mixing if and only if its Pinsker partition is trivial, i. e. it consists of only sets with measure zero and one, see [K-S-F(1980)].) In order to show that the Pinsker partition is trivial, in [S-W(1989)] we constructed a pair of measurable partitions (ξ^s, ξ^u) for $(M \times G, S, \mu \times \lambda)$ made up by open and connected submanifolds of the local stable and unstable manifolds, respectively. It followed by standard methods (see [Sin(1968)]) that the partition π is coarser than each of ξ^s and ξ^u. Due to the S-invariance of π, we have that π is coarser than

$$(*) \qquad \bigwedge_{n\in\mathbb{Z}} S^n\xi^s \wedge \bigwedge_{n\in\mathbb{Z}} S^n\xi^u.$$

In the final step, by using now the rational independence of the masses, we showed that the partition in $(*)$ is, indeed, trivial.

8.2 The role of Proposition 3.1. By taking a look at §3, we can see that Proposition 3.1 (with its rather involved algebraic proof) was only used to prove the so-called Chernov-Sinai Ansatz, an important, necessary condition of the Theorem on Local Ergodicity. It is exactly the algebraic proof of Proposition 3.1 that necessitates the dropping of a null set of geometric parameters $(m_1, \ldots, m_N; r)$ in such an implicit way that for any given $(N+1)$-tuple $(m_1, \ldots, m_N; r)$ one cannot tell (based upon the presented methods) if that $(N+1)$-tuple belongs to the exceptional null set, or not. This is a pity, indeed, since we cannot make it sure (for any specified $(N+1)$-tuple $(m_1, \ldots, m_N; r)$) that the billiard flow $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ is ergodic. Thus, it would be really pleasant to find any other way of proving the Ansatz in order to avoid the necessary dropping of a null set of parameters. Most experts are absolutely convinced that, in fact, this exceptional null set is actually empty, i. e. $(\mathbf{M}, \{S^t\}_{t\in\mathbb{R}}, \mu)$ is ergodic for every $(N+1)$-tuple $(m_1, \ldots, m_N; r)$).

Without Proposition 3.1, the results of §4–7 (the non-existence of the exceptional J-manifold) are easily seen to yield the following, relaxed version of the Chernov-Sinai Ansatz:

Proposition 8.3 (Ansatz, relaxed version). *The closed set $B \subset \mathcal{SR}^+$ of phase points $x \in \mathcal{SR}^+$ with non-sufficient semi-orbit $S^{(0,\infty)}x$ is of first category in any $(2d-3)$-dimensional cell C of $\mathcal{SR}^+$, which is now equivalent to saying that B has an empty interior in C.*

References

[A-P(1995)] P. K. Agarwal, J. Pach, *Combinatorial Geometry*, Wiley, New York.

[B-L-P-S(1992)] L. Bunimovich, C. Liverani, A. Pellegrinotti, Yu. Sukhov, *Special Systems of Hard Balls that Are Ergodic*, Commun. Math. Phys. **146**, 357-396.

[B-S(1973)] L. A. Bunimovich, Ya. G. Sinai, *The fundamental theorem of the theory of scattering billiards*, Math. USSR-Sb. **19**, 407-423.

[B-F-K(1998)] D. Burago, S. Ferleger, A. Kononenko, *Uniform estimates on the number of collisions in semi-dispersing billiards*, Annals of Mathematics **147**, 695-708.

[C-H(1996)] N. I. Chernov, C. Haskell, *Nonuniformly hyperbolic K-systems are Bernoulli*, Ergod. Th. & Dynam. Sys. **16**, 19-44.

[E(1978)] R. Engelking, *Dimension Theory*, North Holland.

[G(1981)] G. Galperin, *On systems of locally interacting and repelling particles moving in space*, Trudy MMO **43**, 142-196.

[K-S(1986)] A. Katok, J.-M. Strelcyn, *Invariant Manifolds, Entropy and Billiards; Smooth Maps with Singularities*, Lecture Notes in Mathematics **1222**, Springer Verlag.

[K(1948)] L. M. Kelly, American Math. Monthly **55**.

[K-S-F(1980)] I. P. Kornfeld, Ya. G. Sinai, S. V. Fomin, *Ergodic Theory*, Nauka, Moscow 1980.

[K-S-Sz(1989)] A. Krámli, N. Simányi, D. Szász, *Ergodic Properties of Semi–Dispersing Billiards I. Two Cylindric Scatterers in the 3–D Torus*, Nonlinearity **2**, 311–326.

[K-S-Sz(1990)] A. Krámli, N. Simányi, D. Szász, *A "Transversal" Fundamental Theorem for Semi-Dispersing Billiards*, Commun. Math. Phys. **129**, 535–560.

[K-S-Sz(1991)] A. Krámli, N. Simányi, D. Szász, *The K–Property of Three Billiard Balls*, Annals of Mathematics **133**, 37–72.

[K-S-Sz(1992)] A. Krámli, N. Simányi, D. Szász, *The K–Property of Four Billiard Balls*, Commun. Math. Phys. **144**, 107-148.

[K(1942)] N. S. Krylov, *The Processes of Relaxation of Statistical Systems and the Criterion of Mechanical Instability*, Thesis, Moscow; *Republished in English by Princeton University Press*, Princeton N. J. (1979).

[L-W(1995)] C. Liverani, M. Wojtkowski, *Ergodicity in Hamiltonian systems*, Dynamics Reported **4**, 130-202.

[O-W(1998)] D. Ornstein, B. Weiss, *On the Bernoulli Nature of Systems with Some Hyperbolic Structure*, Ergod. Th. & Dynam. Sys. **18**, 441-456.

[P(1977)] Ya. Pesin, *Characteristic Exponents and Smooth Ergodic Theory*, Russian Math. surveys **32**, 55-114.

[R(1978)] D. J. Rudolph, *Classifying the isometric extensions of a Bernoulli shift*, J. d'Anal. Math. **34**, 36-50.

[Sim(1992-A)] N. Simányi, *The K-property of N billiard balls I*, Invent. Math. **108**, 521-548.

[Sim(1992-B)] N. Simányi, *The K-property of N billiard balls II. Computation of neutral linear spaces*, Invent. Math. **110**, 151-172.

[Sim(2000)] N. Simányi, *The Complete Hyperbolicity of Cylindric Billiards*, Ergod. Th. & Dynam. Sys. **To appear**.

[S-Sz(1994)] N. Simányi, D. Szász, *The K-property of 4-D Billiards with Non-Orthogonal Cylindric Scatterers*, J. Stat. Phys. **76, Nos. 1/2**, 587-604.

[S-Sz(1995)] N. Simányi, D. Szász, *The K-property of Hamiltonian systems with restricted hard ball interactions*, Mathematical Research Letters **2, No. 6**, 751-770.

[S-Sz(1999)] N. Simányi, D. Szász, *Hard Ball Systems Are Completely Hyperbolic*, Annals of Math. **149**, 35-96.

[S-Sz(2000)] N. Simányi, D. Szász, *Non-integrability of Cylindric Billiards and Transitive Lie Group Actions*, Ergod. Th. & Dynam. Sys. **20**, 593-610.

[S-W(1989)] N. Simányi, M. P. Wojtkowski, *Two-particle billiard system with arbitrary mass ratio*, Ergod. Th. & Dynam. Sys. **9**, 165-171.

[Sin(1963)] Ya. G. Sinai, *On the Foundation of the Ergodic Hypothesis for a Dynamical System of Statistical Mechanics*, Soviet Math. Dokl. **4**, 1818-1822.

[Sin(1968)] Ya. G. Sinai, *Dynamical systems with countably multiple Lebesgue spectrum II*, Amer. Math. Soc. Transl. **68 (2)**, 34-38.

[Sin(1970)] Ya. G. Sinai, *Dynamical Systems with Elastic Reflections*, Russian Math. Surveys **25:2**, 137-189.

[S-Ch(1987)] Ya. G. Sinai, N.I. Chernov, *Ergodic properties of certain systems of 2–D discs and 3–D balls*, Russian Math. Surveys **(3) 42**, 181-207.

[Sz(1993)] D. Szász, *Ergodicity of classical billiard balls*, Physica A **194**, 86-92.

[Sz(1994)] D. Szász, *The K-property of 'Orthogonal' Cylindric Billiards*, Commun. Math. Phys. **160**, 581-597.

[Sz(1996)] D. Szász, *Boltzmann's Ergodic Hypothesis, a Conjecture for Centuries?*, Studia Sci. Math. Hung **31**, 299-322.

[V(1979)] L. N. Vaserstein, *On Systems of Particles with Finite Range and/or Repulsive Interactions*, Commun. Math. Phys. **69**, 31-56.

[W(1988)] M. Wojtkowski, *Measure theoretic entropy of the system of hard spheres*, Ergod. Th. & Dynam. Sys. **8**, 133-153.

[W(1990)] M. Wojtkowski, *Linearly stable orbits in 3-dimensional billiards*, Commun. Math. Phys. **129, no. 2**, 319-327.