TIAN'S INVARIANT OF THE GRASSMANN MANIFOLD

by Julien Grivaux

ABSTRACT. — We prove that Tian's invariant on the complex Grassmann manifold $G_{p,q}(\mathbb{C})$ is equal to $1/(p+q)$. The method introduced here uses a Lie group of holomorphic isometries which operates transitively on the considered manifolds and a natural imbedding of $\left(\mathbb{P}^1(\mathbb{C})\right)^p$ in $G_{p,q}(\mathbb{C})$.

RÉSUMÉ. — On prouve que l'invariant de Tian sur la grassmannienne $G_{p,q}(\mathbb{C})$ est $1/(p+q)$. La méthode présentée dans cet article utilise un groupe de Lie d'isométries holomorphes qui opère transitivement sur les variétés considérées ainsi qu'un plongement naturel de $\left(\mathbb{P}^1(\mathbb{C})\right)^p$ dans $G_{p,q}(\mathbb{C})$.

1. Introduction

On a complex manifold, an hermitian metric h is characterized by the *1-1 symplectic* form ω defined by $\omega = i\, g_{\lambda\overline{\mu}}\, dz^\lambda \wedge d\overline{z}^\mu$, where $g_{\lambda\overline{\mu}} = h_{\lambda\overline{\mu}}/2$.
The metric is a *Kähler metric* if ω is closed, i. e. $d\omega = 0$; then M is a *Kähler manifold.*
On a Kähler manifold, we can define the *Ricci form* by $R = i\, R_{\lambda\overline{\mu}}\, dz^\lambda \wedge d\overline{z}^\mu$, where $R_{\lambda\overline{\mu}} = -\partial_{\lambda\overline{\mu}} \log |g|$.
A Kähler manifold is *Einstein with factor* k if $R = k\omega$. For instance, choosing a local coordinate system $Z = (z_1, \ldots, z_m)$, the projective space $\mathbb{P}_m(\mathbb{C})$ with the Fubini-Study metric $\omega = i\partial\overline{\partial} \log\left(1 + ||Z||^2\right)$ is Einstein with factor $m+1$.
On a Kähler manifold M, the *first Chern class* $C^1(M)$ is the cohomology class of the Ricci tensor, that is the set of the forms $R + i\partial\overline{\partial}\varphi$, where φ is C^∞ on M. If there is

11th December 2005

• *E-mail :* julien.grivaux@free.fr JULIEN GRIVAUX, Université Pierre et Marie Curie

2000 Mathematics Subject Classification. — 53C55, 32M10.
Key words and phrases. — Kähler manifold, Einstein-Kähler metric, first Chern class, admissible functions, Tian's invariant, grassmannian.

a form in $C^1(M)$ which is positive (resp. negative, zero), then $C^1(M)$ is *positive* (resp. *negative*, *zero*). If a Kähler manifold is Einstein, then $C^1(M)$ and k are both positive (resp. negative, zero). In the negative case, it was proved by Aubin ([**Au1**], see also [**Au4**]), that there exists a unique Einstein-Kähler metric (E.K. metric) on M . It is so for the zero case too ([**Au1**], [**Ya**]). The question for the positive case is still open: some manifolds, such as the complex projective space blown up at one point, do not admit an E.K. metric (for obstructions, see [**Li**] and [**Fu**]). Aubin [**Au2**] and Tian [**Ti**] have shown that for suitable values of holomorphic invariants of the metric, there exists an E.K. metric on M.

For $\omega/2\pi$ in $C^1(M)$, *Tian's invariant* $\alpha(M)$ is the supremum of the set of the real numbers α satisfying the following: there exists a constant C such that the inequality $\displaystyle\int_M e^{-\alpha\varphi} \leq C$ holds for all the C^∞ functions φ with $\omega+i\partial\overline{\partial}\varphi > 0$ and $\sup\varphi \geq 0$, where $\omega = i\, g_{\lambda\overline{\mu}}\, dz^\lambda \wedge d\overline{z}^\mu$ is the metric form. Such functions φ are said *ω-admissible*.

In [**Ti**], Tian established that if $\alpha(M) > m/(m+1)$, m being the dimension of M, there exists an E.K. metric on M. This condition is not necessary: it does not hold on the projective space, where Tian's invariant is $1/(m+1)$.

In the same paper, Tian introduces a more restrictive invariant $\alpha_G(M)$, considering only the admissible functions φ invariant by the action of a compact group G of holomorphic isometries. The sufficient condition for the existence of an E.K. metric on M remains $\alpha_G(M) > m/(m+1)$; it is more easily satisfied if the group G is rich enough.

In many cases, the group G is a non-discrete Lie group. The invariant $\alpha_G(M)$ can be computed using subharmonic functions methods and the maximum principle (for effective examples, see [**Be1**], [**Be2**], [**Be-Ch1**], [**Be-Ch2**], [**Re**]).

In this paper, we prove the following theorem:

THEOREM 1.1. — *Tian's invariant on $G_{p,q}(\mathbb{C})$ is given by $\alpha\left(G_{p,q}(\mathbb{C})\right) = 1/(p+q)$.*

This generalizes the known result on $\mathbb{P}^m(\mathbb{C})$ ([**Ti**], see also [**Au3**]). Let us also mention that Tian's invariant has been computed on $\mathbb{P}^m(\mathbb{C})$ blown up at one point and on certain Fermat hypersurfaces using Hörmander L^2 estimates for the $\overline{\partial}$-equation ([**Ti**]).

We first compute the volume element of the metric $\mathscr{G}_{p,q}$; then we will establish some general preliminary results concerning Tian's invariant as well as imbeddings of $\left\{\mathbb{P}^1(\mathbb{C})\right\}^p$ in $G_{p,q}(\mathbb{C})$ which allow us to deduce $\alpha\left(G_{p,q}(\mathbb{C})\right)$ from $\alpha\left(\mathbb{P}^1(\mathbb{C})\right)$.

2. Basic properties of the Grassmann manifold

We propose here a short survey of the properties of Grassmann manifold (for more details, see [**Ko-No**]). We denote by $G_{p,q}(\mathbb{C})$ the set of the subspaces of dimension p in $\mathbb{C}^{p+q}$; in particular, $G_{1,m}(\mathbb{C})$ is the complex projective space of dimension m. It is known (see [**Au3**]) that on $\mathbb{P}_m(\mathbb{C})$, the Fubini-Study metric is Einstein with factor $m+1$ and that Tian's invariant is $1/(m+1)$. Now, let $M^*(p+q,p)$ be the set of the matrices of rank p in $M_{p+q,p}(\mathbb{C})$. The group $Gl_p(\mathbb{C})$ acts by multiplication on the right on $M^*(p+q,p)$. More precisely $\big(M^*(p+q,p),\ \pi,\ G_{p,q}(\mathbb{C})\big)$ is a principal fiber bundle with group $Gl_p(\mathbb{C})$. The group $Gl_{p+q}(\mathbb{C})$ acts by multiplication on the left on $M^*(p+q,p)$ and induces an action on $G_{p,q}(\mathbb{C})$; so does the unitary group $U(p+q)$. These groups act transitively on $G_{p,q}(\mathbb{C})$, which shows that $G_{p,q}(\mathbb{C})$ is compact.

We denote by $\mathcal{I}$ the set of all increasing-ordered subsets of p elements in $\{1,\dots,p+q\}$. Let P be an element of $M^*(p+q,p)$, $P=\big(p_{ij}\big)_{\substack{1\le i\le p+q\\ 1\le j\le p}}$. By Cauchy-Binet formula we get: $\det\big({}^tP\overline{P}\big)=\sum_{I\in\mathcal{I}}|\det m_I(P)|^2$, where $m_I(P)$ is the matrix $\big(p_{ij}\big)_{\substack{i\in I\\ 1\le j\le p}}$. The form ω, where $\omega=i\,\partial\overline{\partial}\log\det\big({}^tP\overline{P}\big)$, is invariant by the action of $Gl_p(\mathbb{C})$ on $M^*(p+q,p)$, and so it projects onto a form $\mathscr{G}_{p,q}$. The metric $\mathscr{G}_{p,q}$ is a Kähler metric form on $G_{p,q}(\mathbb{C})$. For $p=1$, this metric on $G_{1,m}(\mathbb{C})$ is the Fubini-Study metric on the complex projective space. The action of the unitary group $U(p+q)$ on $G_{p,q}(\mathbb{C})$ preserves the metric $\mathscr{G}_{p,q}$ so that $U(p+q)$ is a group of holomorphic isometries which operates transitively on $G_{p,q}(\mathbb{C})$. For I in $\mathcal{I}$, let U_I be the set of the matrices P in $M^*(p+q,p)$ such that $\det(m_I(P))$ is non-zero. Then $\pi(U_I)$ is a coordinate open set on $G_{p,q}(\mathbb{C})$, the matrix Z_I in $M_{p,q}(\mathbb{C})$ is the coordinate, the inverse of the chart φ_I sends $M^*(p+q,p)$ onto $\pi(U_I)$ and we have $m_I\big(\varphi_I^{-1}(Z_I)\big)=I^{(p)}$ where $I^{(p)}$ is the $p\times p$ identity matrix, and $m_{I^c}\big(\varphi_I^{-1}(Z_I)\big)=Z_I$.

LEMMA 2.1. — *For I in $\mathcal{I}$, let λ_I be the map from $\pi(U_I)$ to $\mathbb{R}_+$ defined by*

$$\lambda_I(Z_I)=\big|\det(\mathrm{Id}+{}^tZ_I\overline{Z}_I)\big|^{-(p+q)}.$$

Then $\big(\lambda_I\big)_{I\in\mathcal{I}}$ are the components of a maximal differential form η on $G_{p,q}(\mathbb{C})$, namely:

$$\eta=\lambda_I\big(i/2\big)^{pq}\big(dZ\wedge d\overline{Z}\big)_I.$$

Proof. — It suffices to show that the following transformation rule holds:

$$\text{for every } I,\widetilde{I} \text{ in } \mathcal{I},\ \lambda_I \text{ is equal to } \lambda_{\widetilde{I}}\times\left|\det\ \frac{\partial\, Z_{\widetilde{I}}}{\partial\, Z_I}\right|^2 \text{ on } \pi(U_I)\cap\pi(U_{\widetilde{I}}).$$

Let P_I be the matrix $\varphi_I^{-1}(Z_I)$. Then $P_I\big\{m_{\widetilde{I}}(P_I)\big\}^{-1}=P_{\widetilde{I}}$, so $Z_{\widetilde{I}}=m_{\widetilde{I}^c}(P_I)\,\big\{m_{\widetilde{I}}(P_I)\big\}^{-1}$. The differential of the map which sends Z_I on P_I is the map which sends H on $\breve{H}$, where

$m_{I^c}(\breve{H}) = H$ and $m_I(\breve{H}) = 0$. The change of charts sending Z_I on $Z_{\widetilde{I}}$, we obtain

$$\begin{aligned} D\,Z_{\widetilde{I}}(H) &= m_{\widetilde{I}^c}(\breve{H})\left\{m_{\widetilde{I}}(P_I)\right\}^{-1} - m_{\widetilde{I}^c}(P_I)\left\{m_{\widetilde{I}}(P_I)\right\}^{-1} m_{\widetilde{I}}(\breve{H})\left\{m_{\widetilde{I}}(P_I)\right\}^{-1} \\ &= \left(m_{\widetilde{I}^c}(\breve{H}) - \gamma\, m_{\widetilde{I}}(\breve{H})\right)\alpha^{-1}, \\ \text{where} \quad \alpha &= m_{\widetilde{I}}(P_I),\ \beta = m_{\widetilde{I}^c}(P_I) \text{ and } \gamma = \beta\,\alpha^{-1}. \end{aligned}$$

Let us define a map u from $M_{q,p}(\mathbb{C})$ to $M_{q,p}(\mathbb{C})$ by $u(H) = m_{\widetilde{I}^c}(\breve{H}) - \gamma\, m_{\widetilde{I}}(\breve{H})$. We can choose $I = \{q+1, \ldots, q+p\}$ and $\widetilde{I} = \{1, \ldots, r\} \cup \{q+1+r, \ldots, q+p\}$, where $0 \leq r \leq \inf(p,q)$. We define the $k \times l$ matrix $E_{i,j}^{(k\times l)}$ by $\left(E_{i,j}^{(k\times l)}\right)_{\lambda\mu} = \delta_{i\lambda}\,\delta_{j\mu}$. We have

$$\begin{aligned} m_{\widetilde{I}}\left(\breve{E}_{i,j}^{(q\times p)}\right) &= E_{i,j}^{(p\times p)} \quad \text{if } i \leq r, \text{ and 0 if } i > r, \\ \text{and} \quad m_{\widetilde{I}^c}\left(\breve{E}_{i,j}^{(q\times p)}\right) &= E_{i-r,j}^{(q\times p)} \quad \text{if } i > r, \text{ and 0 if } i \leq r. \text{ Hence} \\ \left(\gamma\, m_{\widetilde{I}}(\breve{E}_{i,j}^{(q\times p)})\right)_{\alpha\beta} &= \gamma_{\alpha i}\, m_{\widetilde{I}}(\breve{E}_{i,j}^{(q\times p)})_{ij}\,\delta_{j\beta} = \gamma_{\alpha i}\,\delta_{j\beta} \quad \text{if } i \leq r, \text{ and 0 elsewhere.} \end{aligned}$$

Now the map which sends H to $\gamma\, m_{\widetilde{I}}(\breve{H})$ can be restricted if $1 \leq j \leq p$ to the span B_j of the $\left(E_{i,j}\right)_{1\leq i\leq q}$. The r first columns of its matrix are those of γ, the others are 0. The map which sends H to $\gamma\, m_{\widetilde{I}^c}(\breve{H})$ maps also B_j into itself. The right upper block of its matrix is $I^{(q-r)}$, the other elements are 0. This allows us to compute the matrix of the restriction of u to B_j, whose determinant is $(-1)^{r\times(q-r)} \det\left(\gamma_{ij}\right)_{\substack{q-r+1\leq i\leq q \\ 1\leq j\leq r}}$. So $\det u = (-1)^{p\times r\times(q-r)} \left[\det\left(\gamma_{i\,j}\right)_{\substack{q-r+1\leq i\leq q \\ 1\leq j\leq r}}\right]^p$. For $1 \leq i \leq q$, let C_i be the span of the $(E_{i,j})_{1\leq j\leq p}$. Each C_i is stable by the map from $M_{q,p}(\mathbb{C})$ to $M_{q,p}(\mathbb{C})$ which sends H to $H\,\alpha^{-1}$. The matrix of the restriction is α^{-1}, so the determinant of the map is $(\det\alpha)^{-q}$. Hence

$$\left|\det DZ_{\widetilde{I}}(H)\right|^2 = \left|\det\left(\gamma_{i,j}\right)_{\substack{q-r+1\leq i\leq q \\ 1\leq j\leq r}}\right|^{2p} \times \left|\det\alpha\right|^{-2q}.$$

Let A be the right $r \times r$ upper block of α. The left $(p-r)\times(p-r)$ lower block of α is $I^{(p-r)}$ and the right $(p-r)\times r$ lower block is 0, so $\det\alpha = (-1)^{r(p-r)}\det A$. The left $r\times(p-r)$ lower block of β is 0, the right $r\times r$ block is $I^{(r)}$ so that the left $r\times r$ lower block of γ is A^{-1}.
From this we deduce $\left|\det\, DZ_{\widetilde{I}}(H)\right|^2 = \left|\det\alpha\right|^{-2(p+q)}$. Since $P_I\,\alpha^{-1} = P_{\widetilde{I}}$, we have

$$\lambda_{\widetilde{I}} = \left|\det\left({}^tP_{\widetilde{I}}\overline{P}_{\widetilde{I}}\right)\right|^{-(p+q)} = \left|\det\alpha\right|^{2(p+q)}\lambda_I = \left|\det\frac{\partial Z_{\widetilde{I}}}{\partial Z_I}\right|^{-2}\lambda_I.$$

□

LEMMA 2.2. — *The unitary group $U(p+q)$ preserves η.*

Proof. — We call I the set $\{q+1,\dots,q+p\}$. We define P_I in $\pi(U_I)$ by $P_I=\varphi_I^{-1}(Z_I)$. Let U be an element in $U(p+q)$ such that $m_I(UP_I)$ is invertible. Let $\tilde{P}_I=UP_I\left\{m_I(UP_I)\right\}^{-1}$ and $\tilde{Z}_I=m_{I^c}(\tilde{P}_I)$. We have $\tilde{Z}_I=m_{I^c}(U)\,P_I\left\{m_I(U)P_I\right\}^{-1}$. So

$$D\tilde{Z}_I(H)=m_{I^c}(U)\left[\breve{H}\left\{m_I(U)P_I\right\}^{-1}-P_I\left\{m_I(U)P_I\right\}^{-1}m_I(U)\,\breve{H}\left\{m_I(U)P_I\right\}^{-1}\right].$$

Thus $D\tilde{Z}_I(H)=X\breve{H}\delta^{-1}$, where $\delta=m_I(U)P_I$ and $X=m_{I^c}(U)\big[I^{(p+q)}-P_I\,\delta^{-1}\,m_I(U)\big]$. Let X_1 be the $q\times q$ matrix of the q first columns of X. Then, $X\breve{H}=X_1H$ and we get $D\tilde{Z}_I(H)=X_1H\delta^{-1}$. The determinant of the map from $M_{q,p}(\mathbb{C})$ to $M_{q,p}(\mathbb{C})$ which sends H to $H\delta^{-1}$ is $(\det\delta)^{-q}$. The determinant of the map from $M_{q,p}(\mathbb{C})$ to $M_{q,p}(\mathbb{C})$ which sends H to X_1H is $(\det X_1)^p$, so $\det D\tilde{Z}_I=(\det X_1)^p\,(\det\delta)^{-q}$. We divide U into four blocks:

$$U=\begin{pmatrix}U_q & U_{q,p}\\ U_{p,q} & U_p\end{pmatrix},\quad U_q\in M_q(\mathbb{C}),\ U_p\in M_p(\mathbb{C}),\ U_{p,q}\in M_{p,q}(\mathbb{C}),\ U_{q,p}\in M_{q,p}(\mathbb{C}).$$

Then $\delta=U_{p,q}\,Z_I+U_p$, so $X_1=U_q-\big(U_q\,Z_I+U_{q,p}\big)\big(U_{p,q}\,Z_I+U_p\big)^{-1}U_{q,p}$. Let Z in $M_{p+q,p+q}(\mathbb{C})$ be the matrix with blocks $Z_q=I^{(q)}$, $Z_{p,q}=0$, $Z_{q,p}=Z_I$, $Z_p=I^{(p)}$, the notations being the same as above. Writing $\det U=\det(UZ)$ and using the column transformation $C_1\leftarrow C_1-C_2\left(U_{p,q}\,Z_I+U_p\right)^{-1}U_{p,q}$ where C_1 is made of the first q columns and C_2 of the remaining ones, we get

$$\det U=\det\Big[U_q-\big(U_q\,Z_I+U_{q,p}\big)\left(U_{p,q}\,Z_I+U_p\right)^{-1}U_{p,q}\Big]\times\det\big(U_{p,q}Z_I+U_p\big).$$

Hence $\left|\det\,D\tilde{Z}_I\right|^2=\left|\det\delta\right|^{-2(p+q)}$. We have $\tilde{P}_I=AP_I\delta^{-1}$, so

$$\lambda_{\tilde{I}}=\det\big({}^t\tilde{P}_I\overline{\tilde{P}}_I\big)^{-(p+q)}=\det\big({}^tP_I\overline{P}_I\big)^{-(p+q)}\times\left|\det\delta\right|^{2(p+q)}=\lambda_I\left|\det D\tilde{Z}_I\right|^{-2},$$

which proves the result. □

Proposition 2.3. — 1. $dV\left(\mathscr{G}_{p,q}\right)=\eta$.

2. *If* $I\in\mathcal{I}$, $\left|\mathscr{G}_{p,q}\right|_I=\left\{\det\big(I^{(p)}+{}^t\,Z_I\overline{Z}_I\big)\right\}^{-(p+q)}$.

3. $\mathcal{R}\left(\mathscr{G}_{p,q}\right)=(p+q)\mathscr{G}_{p,q}$.

Proof. — 1. Let I in $\mathcal{I}$. It is easy to compute $\mathscr{G}_{p,q}$ at the point $Z_I=0$: $\mathscr{G}_{p,q}(H,K)=Tr(H\overline{K})$. Then $dV(\mathscr{G}_{p,q})_{\big|_{Z_I=0}}=\big(i/2\big)^{pq}\left(dZ\wedge d\overline{Z}\right)_I=\eta_{\big|_{Z_I=0}}$. Since $dV(\mathscr{G}_{p,q})$ and η are invariant by the transitive action of $U(p+q)$, we have $dV(\mathscr{G}_{p,q})=\eta$.

2. Since $dV(\mathscr{G}_{p,q})=\left|\mathscr{G}_{p,q}\right|_I(i/2)^{pq}\left(dZ\wedge d\overline{Z}\right)_I$, property 1 gives the result.

3. Remark that $\mathscr{G}_{p,q} = i\,\partial\overline{\partial}\,\log\bigl\{\det(I^{(p)} + {}^tZ_I\overline{Z}_I)\bigr\}$. Since $\mathcal{R}\left(\mathscr{G}_{p,q}\right) = -i\,\partial\overline{\partial}\,\log\left|\mathscr{G}_{p,q}\right|_I$, we obtain $\mathcal{R}\left(\mathscr{G}_{p,q}\right) = (p+q)\mathscr{G}_{p,q}$, which expresses that $\mathscr{G}_{p,q}$ is Einstein, with factor $p+q$. □

3. Some general results about Tian's invariant

3.1. Tian's invariant with a normalization on a finite set. — If X is a manifold, we will denote by μ_X a measure on X compatible with the manifold structure.

THEOREM 3.1. — *Let M be a compact Kähler manifold. We suppose that there exists a compact Lie group G of holomorphic isometries. Let $\Delta_n = \{P_1, \ldots, P_n\}$ be a finite subset of M. Let $\alpha(\omega)$ (resp. $\alpha_{\Delta_n}(\omega)$) be the supremum of the set of the nonnegative real numbers α satisfying the condition: there exists a constant C such that the inequality $\int_M e^{-\alpha\varphi} \leq C$ holds for all the ω-admissible functions φ with $\sup\varphi \geq 0$ (resp. with $\varphi(P_i) \geq 0$ for $1 \leq i \leq n$). Suppose in addition that the orbit of each P_i under the action of G has positive measure. Then $\alpha(\omega) = \alpha_{\Delta_n}(\omega)$.*

We first establish a few lemmas which will be useful for the proof.

LEMMA 3.2. — *Let $\left(\varphi_n\right)_{n\geq 0}$ be a sequence of admissible functions with nonnegative maxima. Then there exists a subset Ω of M, with $\mu_M(\Omega) = \mu_M(M)$, and a subsequence φ_{n_k} of φ_n, such that for every ω in Ω, $\left(\varphi_{n_k}(\omega)\right)_{k\geq 0}$ is bounded from below.*

Proof. — Let Q_n be a point such that $\varphi_n(Q_n)$ vanishes. Green's formula runs as follows:

$$\varphi_n(Q_n) = \frac{1}{V}\int_M \varphi_n + \int_M G(Q_n, R)\,\Delta\varphi_n(R)\,dV(R),$$

with $G(Q,R) \leq 0$ and $\int_M G(Q,R)\,dV(R) = -C$, where C is a positive constant. Since φ_n is admissible, $\Delta\varphi_n$ is greater than $-m$, m being the dimension of M. Thus $\int_M |\varphi_n| \leq C\,m\,V$. The sequence $\left(\varphi_n\right)_{n\geq 0}$ is bounded in $L^1(M)$. We can therefore extract a subsequence $\left(\varphi_{n_k}\right)_{n_k\in I}$ which converges almost everywhere to a function φ of $L^1(M)$. Since φ is finite almost everywhere, we get the result. □

LEMMA 3.3. — *Let $\left(\varphi_n\right)_{n\geq 0}$ be a sequence of admissible functions with nonnegative maxima and suppose that there exists a compact group G of holomorphic isometries of M such that the orbit of each P_i has positive measure. Let $\Phi : G \to \mathbb{R}\cup\{-\infty\}$ be the map defined by $\Phi(g) = \inf_{\Delta_n}\inf_{k\geq 0}(\varphi_k \circ g)$. Then there exists g in G such that $\Phi(g)$ is finite.*

Proof. — Suppose that $\Phi \equiv -\infty$. For $i = 1, \ldots, n$, let A_i be the set of the g in G such that $\inf_{k\geq 0}(\varphi_k \circ g)(P_i) = -\infty$. The sets A_i are measurable and $\cup_{i=1}^n A_i = G$, so there exists i such that A_i has positive measure. From Lemma (3.2), $A_i.P_i$ is a subset of Ω^c. Since Ω and M have the same measure, the measure of $A_i.P_i$ vanishes. Let u_i be the map from G to M which sends g to $g(P_i)$. Then u_i has constant rank on G. Indeed, $u_i \circ L(g) = \sigma_g \circ u_i$, where $L(g)$ is the left translation by g and σ_g the map from M to M which sends x to $g.x$. Since $G.P_i$ has positive measure, u_i is a submersion on G, so that $u_i(A_i)$ has positive measure. This is a contradiction since $u_i(A_i) = A_i.P_i$. □

We can now prove Theorem (3.1).

Proof. — It is clear that $\alpha(\omega) \leq \alpha_{\Delta_n}(\omega)$. Conversely, let $\varepsilon > 0$. There exists a sequence $\big(\varphi_n\big)_{n\geq 0}$ of admissible functions with positive maxima such that $\int_M e^{-(\alpha(\omega)+\varepsilon)\varphi_k}$ goes to infinity as k goes to infinity. Replacing φ_n by $\varphi_n - \sup \varphi_n$, we can take $\sup \varphi_n = 0$. First we apply Lemma(3.2). For the sake of simplicity, we take $\varphi_{n_k} = \varphi_k$. From Lemma (3.3), there exists an element g in G such that $\Phi(g)$ is finite; we define Ψ_k by $\Psi_k = \varphi_k \circ g - \Phi(g)$. Since g is an isometry, Ψ_k is ω-admissible, and from the very definition of Φ, $\Psi_k(P_i)$ is nonnegative. Furthermore, $\int_M e^{-(\alpha(\omega)+\varepsilon)\Psi_k} = e^{(\alpha(\omega)+\varepsilon)\Phi(g)} \int_M e^{-(\alpha(\omega)+\varepsilon)\varphi_k}$. This proves that $\int_M e^{-(\alpha(\omega)+\varepsilon)\Psi_k}$ goes to infinity as k goes to infinity. Then, $\alpha_{\Delta_n}(\omega) \leq \alpha(\omega) + \varepsilon$. This inequality holds for every positive ε, and so $\alpha_{\Delta_n}(\omega) \leq \alpha(\omega)$. □

3.2. Tian's invariant on a product. — For a Kähler form ω on a compact Kähler manifold M, $\alpha(\omega)$ is defined as in Theorem (3.1).

Proposition 3.4. — *Let $\big(M_i\big)_{1\leq i\leq n}$ be compact Kähler manifolds with metric forms $\big(\omega_i\big)_{1\leq i\leq n}$. We endow the product $M_1 \times \cdots \times M_n$ with the metric $\omega_1 \oplus \cdots \oplus \omega_n$. Then $\alpha(\omega_1 \oplus \cdots \oplus \omega_n) = \inf_{1\leq i\leq n} \alpha(\omega_i)$.*

Proof. — It suffices to make the proof when $n = 2$, the general result will follow by induction.

(1) Suppose that $\alpha(\omega_1) \leq \alpha(\omega_2)$, and let $\varepsilon > 0$. There exists a sequence $\big(\varphi_n\big)_{n\geq 0}$ of ω_1-admissible functions on M_1 with positive maxima such that $\int_{M_1} e^{-\left(\alpha(\omega_1)+\varepsilon\right)\varphi_n}$ goes to infinity when n goes to infinity. We define ψ_n on $M_1 \times M_2$ by $\psi_n(m_1, m_2) = \varphi_n(m_1)$. Thus ψ_n is $(\omega_1 \oplus \omega_2)$-admissible on $M_1 \times M_2$, with positive maximum, and $\int_{M_1\times M_2} e^{-\left(\alpha(\omega_1)+\varepsilon\right)\psi_n} =$

$V(M_2)\int_{M_1} e^{-\left(\alpha(\omega_1)+\varepsilon\right)\varphi_n}$, so that $\int_{M_1\times M_2} e^{-\left(\alpha(\omega_1)+\varepsilon\right)\psi_n}$ goes to infinity when n goes to infinity. We have therefore $\alpha(\omega_1\oplus\omega_2)\leq\alpha(\omega_1)+\varepsilon$. This yields $\alpha(\omega_1\oplus\omega_2)\leq\alpha(\omega_1)$.

(2) Let us now prove the opposite inequality. Let α be a real number such that $\alpha<\inf\big(\alpha(\omega_1),\alpha(\omega_2)\big)$ and φ an $(\omega_1\oplus\omega_2)$-admissible function on $M_1\times M_2$. If m_2 is in M_2, the function which sends m_1 to $\varphi(m_1,m_2)$ is ω_1-admissible. The same holds for M_1. Let (u,v) in $M_1\times M_2$ be such that $\varphi(u,v)\geq 0$. Then

$$\begin{aligned}\int_{M_1\times M_2} e^{-\alpha\varphi(m_1,m_2)}dV_1\,dV_2 &= \int_{M_1} e^{-\alpha\varphi(m_1,v)}\left(\int_{M_2} e^{-\alpha\left[\varphi(m_1,m_2)-\varphi(m_1,v)\right]}dV_2\right)dV_1\\ &\leq C_2\int_{M_1} e^{-\alpha\varphi(m_1,v)}dV_1\leq C_1\,C_2.\end{aligned}$$

Thus, $\alpha\leq\alpha\big(\omega_1\oplus\omega_2\big)$ and we get $\inf\big(\alpha(\omega_1),\alpha(\omega_2)\big)\leq\alpha\big(\omega_1\oplus\omega_2\big)$. □

3.3. Tian's invariant on $G_{p,q}(\mathbb{C})$. — Since there is a natural duality isomorphism between $G_{p,q}(\mathbb{C})$ and $G_{q,p}(\mathbb{C})$, we can assume that $p\leq q$ without loss of generality.

3.3.1. Imbedding of $\big\{\mathbb{P}^1(\mathbb{C})\big\}^p$ into $G_{p,q}(\mathbb{C})$ when $p\leq q$. — For w in $\mathbb{C}^{p(q-1)}$, $w=\big(w_{i,j}\big)_{\substack{1\leq i\leq q\\ 1\leq j\leq p\\ i\neq j}}$, we define the map $\tilde\rho_w$ from $\Big\{\mathbb{C}^2\setminus(0,0)\Big\}^p$ to $M_{p+q,p}(\mathbb{C})$ by

$$\tilde\rho_w\Big((\lambda_i,\mu_i)_{1\leq i\leq p}\Big)=\begin{cases}\lambda_i\,\delta_{ij} & \text{if}\quad i\leq p\\ w_{i-p,j}\,\lambda_j & \text{if}\quad i>p\text{ and }i\neq j+p\\ \mu_i & \text{if}\quad i>p\text{ and }i=j+p\end{cases}$$

We make, for $p+1\leq i\leq p+q$, the following row transformations: $L_i\leftarrow L_i-\sum_{\substack{1\leq j\leq p\\ i\neq j+p}} w_{i-p,j}\,L_j$. We get a matrix $\big(c_{ij}\big)_{\substack{1\leq i\leq p+q\\ 1\leq j\leq p}}$ with $c_{ij}=\delta_{ij}\,\lambda_i$ if $1\leq i\leq p$ and $c_{ij}=\delta_{i-p,j}\,\mu_j$ if $p+1\leq i\leq p+q$, which has rank p. $\tilde\rho_w$ induces a map from $\big\{\mathbb{P}^1(\mathbb{C})\big\}^p$ into $G_{p,q}(\mathbb{C})$ as shown on the following diagram, where γ is the projection of the principal fiber bundle $\Big\{\mathbb{C}^2\setminus(0,0)\Big\}^p$ onto $\big\{\mathbb{P}^1(\mathbb{C})\big\}^p$. Remark that $\tilde\rho_w$ sends $[0,1]\times\cdots\times[0,1]$ onto $\pi(A)$, where $m_{\{p+1,\ldots,2p\}}(A)=I^{(p)}$ and $m_{\{p+1,\ldots,2p\}^c}(A)=0^{(q\times p)}$.

$$\begin{array}{ccc}\Big\{\mathbb{C}^2\setminus(0,0)\Big\}^p & \xrightarrow{\ \tilde\rho_w\ } & M^*(p+q,p)\\ \Big\downarrow{\scriptstyle\gamma} & & \Big\downarrow{\scriptstyle\pi}\\ \big\{\mathbb{P}^1(\mathbb{C})\big\}^p & \xrightarrow[\ \rho_w\ ]{} & G_{p,q}(\mathbb{C})\end{array}$$

We have

$$\begin{aligned}\big(\pi\circ\tilde\rho_w\big)^*\big(\mathscr{G}_{p,q}\big) &= i\,\partial\overline\partial\,\log\Big(\det\big({}^t\tilde\rho_w\overline{\tilde\rho}_w\big)\Big)\\ &= i\,\partial\overline\partial\,\log\left(\frac{\det\big({}^t\tilde\rho_w\overline{\tilde\rho}_w\big)}{\prod_{k=1}^{p}\big(|\lambda_k|^2+|\mu_k|^2\big)}\right)+\sum_{k=1}^{p} i\,\partial\overline\partial\,\log\big(|\lambda_k|^2+|\mu_k|^2\big)\\ &= i\,\partial\overline\partial\,\log\tilde\Phi+\gamma^*\big(FS_1\oplus\cdots\oplus FS_1\big),\end{aligned}$$

where FS_1 is the Fubini-Study metric on $\mathbb{P}^1(\mathbb{C})$. $\tilde\Phi$ is invariant by the action of the structural group $\mathbb{C}^*\times\cdots\times\mathbb{C}^*$, so it induces a map Φ from $\big\{\mathbb{P}^1(\mathbb{C})\big\}^p$ into $\mathbb{C}$. Note that $\Phi\big([0,1]\times\cdots\times[0,1]\big)=1$. Then $\big(\pi\circ\tilde\rho_w\big)^*\big(\mathscr{G}_{p,q}\big)=\pi^*\big(i\,\partial\overline\partial\,\log\Phi+FS_1\oplus\cdots\oplus FS_1\big)$, so that $\rho_w^*\big(\mathscr{G}_{p,q}\big)=i\,\partial\overline\partial\,\log\Phi+FS_1\oplus\cdots\oplus FS_1$.

3.3.2. Lower bound of $\alpha(\mathscr{G}_{p,q})$. — For I in $\mathcal{I}$, we define P_I by $m_I(P_I)=I^{(p)}$ and $m_{I^c}(P_I)=0^{(q\times p)}$. If $n=\binom{p+q}{p}$, we set $\Delta_n=\big\{P_I\big\}_{I\in\mathcal{I}}$. Since $U(p+q)$ is a transitive group of holomorphic isometries of $G_{p,q}(\mathbb{C})$, we know from proposition (3.1), that $\alpha\big(\mathscr{G}_{p,q}\big)=\alpha_{_{\Delta_n}}\big(\mathscr{G}_{p,q}\big)$. We set $I=\{p+1,\dots,2p\}$. Let φ be an admissible function on $G_{p,q}(\mathbb{C})$, nonnegative on Δ_n. The last equality of the precedent section shows that the function $\varphi\circ\rho_w+\log\Phi$ is $\big(FS_1\oplus\cdots\oplus FS_1\big)$-admissible for every w in $\mathbb{C}^{p(q-1)}$. Furthermore, $\big(\varphi\circ\rho_w+\log\Phi\big)$ sends $[0,1]\times\cdots\times[0,1]$ to the nonnegative number $\varphi(P_{_I})$. It is known that $\alpha(FS_1)=1$ (see [**Au3**]). Proposition (3.4) yields $\alpha\big(FS_1\oplus\cdots\oplus FS_1\big)=1$.
Let α be a real number such that $\alpha<1$. There exists a constant C, independent of φ, such that $\displaystyle\int_{\left\{\mathbb{P}^1(\mathbb{C})\right\}^p} e^{-\alpha\varphi\circ\rho_w}\,\Phi^{-\alpha}\le C$. We define the map F_I from $\pi(U_I)$ to $\mathbb{R}_+$ by $F_I(Z_I)=\det\big(Id+{}^tZ_I\overline{Z}_I\big)$. On $\big\{\mathbb{P}^1(\mathbb{C})\big\}^p$, we work with the coordinates $\mu_1,\dots,\mu_p$ in the chart $\lambda_1=\cdots=\lambda_p=1$. Thus

$$\Phi(\mu)=\frac{F_I\circ\rho_w(\mu)}{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)},\quad\text{so that}\quad\int_{\mu\in\mathbb{C}^p} e^{-\alpha\varphi\circ\rho_w(\mu)}\,\frac{dV_\mu\big(\mathbb{C}^p\big)}{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)^{2-\alpha}\,\big(F_{_I}\circ\rho_w(\mu)\big)^{\alpha}}\le C.$$

We have the inequality $\sum_{i=1}^{q}\sum_{j=1}^{p}\big|Z_{ij}\big|^2 \leq F_I\big(P_I\big)$. In particular, for every k in $\{1,\ldots,p\}$, $1+|\mu_k|^2 \leq f_I\circ\rho_w(\mu)$, and $f_I\circ\rho_w(\mu) \geq 1+\sum_{\substack{1\leq i\leq q\\ 1\leq j\leq p\\ i\neq j}}\big|w_{ij}\big|^2$. Thus, for $\kappa>0$ and $w\in\mathbb{C}^{p(q-1)}$,

$$\frac{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)^{2-\alpha}}{\big(F_I\circ\rho_w(\mu)\big)^{\kappa+p+q-\alpha}} \leq \frac{1}{\big(F_I\circ\rho_w(\mu)\big)^{\kappa-p+q+\alpha(p-1)}} \leq \frac{1}{\Big(1+\sum_{\substack{1\leq i\leq q\\ 1\leq j\leq p\\ i\neq j}}\big|w_{ij}\big|^2\Big)^{\kappa}} = \frac{1}{\Big(1+\big\|w\big\|^2\Big)^{\kappa}}.$$

We have, according to Proposition (2.3),

$$\int_{\pi(U_I)}\frac{e^{-\alpha\varphi}}{F_I^{\kappa}} = \int_{w\in\mathbb{C}^{p(q-1)}}\int_{\mu\in\mathbb{C}^p}\frac{e^{-\alpha\varphi\circ\rho_w(\mu)}}{\big(F_I\circ\rho_w(\mu)\big)^{\kappa+p+q}}\,dV_\mu\big(\mathbb{C}^p\big)\,dV_w\big(\mathbb{C}^{p(q-1)}\big)$$

$$= \int_{w\in\mathbb{C}^{p(q-1)}}\int_{\mu\in\mathbb{C}^p}\left(\frac{e^{-\alpha\varphi\circ\rho_w(\mu)}}{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)^{2-\alpha}\,\big(F_I\circ\rho_w(\mu)\big)^{\alpha}}\right) \times\frac{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)^{2-\alpha}}{\big(F_I\circ\rho_w(\mu)\big)^{\kappa+p+q-\alpha}}\,dV_\mu\big(\mathbb{C}^p\big)\,dV_w\big(\mathbb{C}^{p(q-1)}\big)$$

$$= \int_{w\in\mathbb{C}^{p(q-1)}}\left(\int_{\mu\in\mathbb{C}^p}\frac{e^{-\alpha\varphi\circ\rho_w(\mu)}}{\prod_{k=1}^{p}\big(1+|\mu_k|^2\big)^{2-\alpha}\,\big(F_I\circ\rho_w(\mu)\big)^{\alpha}}\,dV_\mu\big(\mathbb{C}^p\big)\right)\times\frac{dV_w\big(\mathbb{C}^{p(q-1)}\big)}{\Big(1+\big\|w\big\|^2\Big)^{\kappa}}$$

$$\leq C\int_{w\in\mathbb{C}^{p(q-1)}}\frac{dV_w\big(\mathbb{C}^{p(q-1)}\big)}{\Big(1+\big\|w\big\|^2\Big)^{\kappa}} \leq C' \qquad \text{if } \kappa>p(q-1).$$

Thus, we obtain that for all I in $\mathcal{I}$, $\int_{\pi(U_I)}\frac{e^{-\alpha\varphi}}{F_I^{\kappa}}\leq C$, where C is independent of φ.

Since $G_{p,q}(\mathbb{C})$ is compact, there exists a family $\big(V_I\big)_{I\in\mathcal{I}}$ of open sets of $G_{p,q}(\mathbb{C})$ such that V_I is relatively compact in $\pi(U_I)$ for every $I\in\mathcal{I}$, and $\bigcup_{I\in\mathcal{I}}V_I = G_{p,q}(\mathbb{C})$. There exists $M>0$ such that $F_I\leq M$ on V_I for every $I\in\mathcal{I}$. Thus

$$\int_{G_{p,q}(\mathbb{C})}e^{-\alpha\varphi} \leq \sum_{I\in\mathcal{I}}\int_{V_I}e^{-\alpha\varphi} \leq \sum_{I\in\mathcal{I}}M^{\kappa}\int_{V_I}\frac{e^{-\alpha\varphi}}{F_I^{\kappa}} \leq M^{\kappa}\sum_{I\in\mathcal{I}}\int_{\pi(U_I)}\frac{e^{-\alpha\varphi}}{F_I^{\kappa}} \leq C\,M^{\kappa}\binom{p+q}{p}.$$

We deduce that $\alpha\big(\mathscr{G}_{p,q}\big)\geq 1$.

3.3.3. Upper bound of $\alpha(\mathscr{G}_{p,q})$. — We use here a method which can be found in [**Re**] for the complex projective space. Let I in $\mathcal{I}$. We define $\tilde{K}$ from $M^*(p+q,p)$ to $\mathbb{P}^1(\mathbb{R})$ by the relation $\tilde{K}(M) = \big[\,|\det m_{_I}(H)\,|^2, \det {}^tM\overline{M}\,\big]$. $\tilde{K}$ is invariant by the action of the structural group $G_p(\mathbb{C})$, so it induces a C^∞ map K from $G_{p,q}(\mathbb{C})$ to $\mathbb{P}^1(\mathbb{R})$. Remark that $\psi = \log K$ is a Kähler potential on U_I for the metric $\mathscr{G}_{p,q}$.

LEMMA 3.5. — *There exists a decreasing sequence $\big(\varphi_n\big)_{n\geq 0}$ of admissible functions with positive maxima which converges pointwise to $-\psi$ on $\pi\big(U_I\big)$.*

Proof. — We construct a decreasing sequence $\big(f_n\big)_{n\geq 0}$ of C^∞ convex functions on $\mathbb{R}_+$ satisfying the conditions $1 + f_n' > 0$, $f_n(x) = -\big(1 - 1/n\big)x$ for x in $[0,n]$ and $f_n(x) = -n$ for $x \geq 2n$. Let y be an element of $\pi(U_I)^c$ and Ω_n the set of the elements x in $\pi(U_I)$ such that $\psi(x) > 2n$. Since $F_I(y) = [0,1]$, there exists a neighborhood V of y such that the inequality $z > e^{2n}$ holds for every point $[1,z]$ in $F_I(V)$. Thus $V \cap \pi\big(U_I\big)$ is included in Ω_n. We have proved that $W_n = \Omega_n \cup \pi\big(U_I\big)^c$, so that W_n is an open neighborhood of $\pi\big(U_I\big)^c$. We define φ_n by $\varphi_n = f_n \circ \psi$ on $\pi\big(U_I\big)$ and $\varphi_n = -n$ on W_n. Thus φ_n is well defined and $\varphi_n(0) = 0$. It remains to show that φ_n is admissible on $\pi\big(U_I\big)$. We have

$$\Big(\mathscr{G}_{p,q} + i\,\partial\overline{\partial}\,\varphi_n\Big)_{\lambda\overline{\mu}} = \partial_{\lambda\overline{\mu}}\psi + \partial_\lambda\big(f_n' \circ \psi\big)\partial_{\overline{\mu}}\psi = \big(1 + f_n' \circ \psi\big)\partial_{\lambda\overline{\mu}}\psi + f_n'' \circ \psi\,\partial_\lambda\psi\,\partial_{\overline{\mu}}\psi.$$

Hence the matrix of the metric $\mathscr{G}_{p,q} + i\,\partial\overline{\partial}\,\varphi_n$ is of the form $A + T$ where A is positive definite and T has rank one and positive trace. So $A + T$ is positive definite and we get the result. □

LEMMA 3.6. — *Let n in $\mathbb{N}^*$ and r a positive real number. Then*

$$\int_{||X||\leq r} \frac{dV_X(M_n(\mathbb{C}))}{\big|\det X\big|^2} = +\infty.$$

Proof. — We can write

$$\int_{||X||\leq r} \frac{dV_X(M_n(\mathbb{C}))}{\big|\det X\big|^2} = \sum_{k=0}^{\infty}\ \int_{r/2^{k+1}\leq||X||\leq r/2^k} \frac{dV_X(M_n(\mathbb{C}))}{\big|\det X\big|^2}.$$

We put $Y = 2^k X$, so

$$\int_{r/2^{k+1}\leq||X||\leq r/2^k} \frac{dV_X(M_n(\mathbb{C}))}{\big|\det X\big|^2} = \int_{1/2\leq||Y||\leq 1} \frac{dV_Y(M_n(\mathbb{C}))}{\big|\det Y\big|^2}.$$

The terms in the series are strictly positive and independent of k. The sum is therefore infinite. □

We can now prove that $\alpha\big(\mathscr{G}_{p,q}\big)$ is upper bounded by 1. Suppose that $\alpha\big(\mathscr{G}_{p,q}\big) > 1$. Then there exists a positive C such that for every integer n, $\displaystyle\int_{\pi(U_I)} e^{-\varphi_n} \leq C$. Using Lemma (3.5) and monotonous convergence, $\displaystyle\int_{\pi(U_I)} F_I \leq C$. Since $\pi(U_I)^c$ has zero measure, $\displaystyle\int_{G_{p,q}(\mathbb{C})} F_I \leq C$. Let $\tilde{I}$ in $\mathcal{I}$ be such that $I \cap \tilde{I} = \emptyset$ (this is possible since $p \leq q$). We have $P_{\tilde{I}}\big\{m_I\big(P_{\tilde{I}}\big)\big\}^{-1} = P_I$. Remark that $m_I\big(P_{\tilde{I}}\big) = m_I\big(Z_{\tilde{I}}\big)$. Thus $\det\big(Id + {}^t Z_I \overline{Z}_I\big) = \det\big({}^t P_{\tilde{I}} \overline{P}_{\tilde{I}}\big)\,\big|\det m_I\big(Z_{\tilde{I}}\big)\big|^{-2}$. For $\big\|Z_{\tilde{I}}\big\| \leq r$, $\det\big({}^t P_{\tilde{I}} \overline{P}_{\tilde{I}}\big) \leq M$, so that $\displaystyle\int_{||Z_{\tilde{I}}||\leq r} \frac{dV_{Z_{\tilde{I}}}(M_{q,p}(\mathbb{C}))}{\big|\det m_I\big(Z_{\tilde{I}}\big)\big|^2} < +\infty$. Integrating over the remaining variables $\big(Z_{ij}\big)_{i\in\tilde{I}^c\cap I^c}$ yields $\displaystyle\int_{||Z||\leq r} \frac{dV_Z(M_p(\mathbb{C}))}{\big|\det Z\big|^2} < +\infty$, which is in contradiction with the result of Lemma (3.6). Thus we obtain $\alpha\big(\mathscr{G}_{p,q}\big) \leq 1$.

References

[Au1] T. Aubin, *Équations du type Monge-Ampère sur les variétés kählériennes compactes*, Bull. Sci. Math. **102** (1978), 63-95.

[Au2] T. Aubin, *Réduction du cas positif de l'équation de Monge-Ampère sur les variétés Kählériennes compactes à la démonstration d'une inégalité*, J. Funct. Anal. **57** (1984), 143-153.

[Au3] T. Aubin, *Métriques d'Einstein-Kähler et exponentiel de fonctions admissibles*, J. Func. Anal. **88** (1990), 385-394.

[Au4] T. Aubin, *Some Non-linear Problems in Riemannian Geometry*, Springer-Verlag, Berlin, 1998.

[Be1] A. Ben Abdesselem, *Lower bound of admissible functions on sphere*, Bull. Sci. Math **126** (2002), 675-680.

[Be2] A. Ben Abdesselem, *Enveloppes inférieures de fonctions admissibles sur l'espace projectif complexe. Cas symétrique*, To appear in Diff. Geom. and its App.

[Be-Ch1] A. Ben Abdesselem, P. Cherrier, *Einstein-Kähler metrics on a class of bundles involving integral weights*, J. Math. Pures Appl. **3**, vol 81, (2002), 259-281.

[Be-Ch2] A. Ben Abdesselem, P. Cherrier, *Estimations of Ricci tensor on certain Fano manifolds*, Math. Z. **233**, 2000, 481-505.

[Fu] A. Futaki, *An obstruction to the existence of Kähler-Einstein metrics*, Invent. Math. **73**, 1983, 437-443.

[Ko-No] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry*, Volume II John Wiley (1969).

[Li] A. Lichnerowicz, *Sur les transformations analytiques des variétés kählériennes*, Cr. Acac. Sci **244**, 1957, 3011-3014.

[Ma] Y. Matsushima, *Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145-150.

[Re] C. Real, *Métriques d'Einstein-Kähler sur des variétés à première classe de Chern positive*, J. Func. Anal. **106** (1992), 145-188.

[Ti] G. Tian, *On Kähler-Einstein metrics on certain Kähler manifolds with* $C^1(M) > 0$, Invent. Math. **89** (1987), 225-246.

[Ya] S. T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equations*, I. Comm. Pure Appl. Math. **31** (1978), 339-411.