Coupled Map Lattices with Asynchronous Updatings

Torsten Fischer*
Niels Bohr Institute
DK-2100 Copenhagen

July 10, 1999

Abstract

We consider on $M = (S^1)^{\mathbb{Z}^d}$ a family of continuous local updatings, of finite range type or Lipschitz-continuous in all coordinates with summable Lipschitz-constants. We show that the infinite-dimensional dynamical system with independent identically Poisson-distributed times for the individual updatings is well-defined. In the setting of analytically coupled uniformly expanding, analytic circle maps with weak, exponentially decaying interaction, we define transfer operators for the infinite-dimensional system, acting on Banch-spaces that include measures whose finite-dimensional marginals have analytic, exponentially bounded densities. We prove existence and uniqueness (in the considered Banach space) of a probability measure and its exponential decay of correlations.

0 Introduction

In this paper we study coupled map lattices with indepentent identically (i.i.) Poisson-distributed updatings at the individual sites.
A deterministic coupled map lattice (CML) is given by a $\mathbb{Z}^d$-lattice with a copy of the same Riemannian manifold at each lattice point (i.e. the state space is the product of these manifolds with index set $\mathbb{Z}^d$) and a map on the infinite space that can be decomposed into an uncoupled map that acts individually on each component and an 'interaction step' where the change of each coordinate depends also on the other sites.
L.A. Bunimovich and Y.G. Sinai prove in [8] (cf. also the remarks on this in [5]) the existence and uniqueness of an invariant measure and its exponential decay of correlations for a one-dimensional lattice of interval maps with weak coupling. By constructing a Markov partition they relate the system to a two-dimensional spin system whose Gibbs measure corresponds to the invariant measure of the CML.
G. Keller and M. Künzle prove in [21] the existence and uniqueness of an invariant measure for periodic or infinite one-dimensional lattices of weakly coupled interval maps by studying the transfer operators on the space of measures whose finite-dimensional marginals have densities of bounded variation. For small perturbation of the uncoupled map any invariant measure is 'close' to the one they found.
J. Bricmont and A. Kupiainen extend in [4] and [5, 6] the result of Bunimovich and Sinai [8] to $\mathbb{Z}^d$-lattices of weakly coupled circle maps with analytic and Hölder-continuous interaction, respectively.
They represent the iterates of the Perron-Frobenius operator for finite-dimensional subsystems (over $\Lambda \subset \mathbb{Z}^d$) by a 'polymer'- or 'cluster'-expansion that gives rise to a representation of the corresponding invariant measure in terms of a $(d+1)$-dimensional spin system. The weak limit (as $\Lambda \to \mathbb{Z}^d$) of these measures is the unique (in a certain class) invariant probability measure and it is exponentially mixing with respect to (wrt) spatio-temporal shifts.
C. Maes and A. Van Moffaert introduce in [25] for a similar setting as in [4] a simplified 'cluster'-expansion for the truncated Perron-Frobenius operator and show stochastic stability of the CML under stochastic perturbation.

*The largest part of work on this paper was supported by the EC via TMR-Fellowship ERBFMBICT-961157

In [2] V. Baladi, M. Degli Esposti, S. Isola, E. Järvenpää and A. Kupiainen define Frechet spaces, and, for $d = 1$, a Banach space and transfer operators for the infinite-dimensional systems, considered by Bricmont and Kupiainen in [4], and study the spectral properties of these operators.
In [13] we consider analytically coupled circle maps (uniformly expanding and analytic) on the $\mathbb{Z}^d$-lattice with exponentially decaying interaction and introduce Banach spaces for the infinite-dimensional system that include measures whose finite-dimensional marginals have analytic, exponentially bounded densities. We define transfer operators on these spaces, get a unique (in the considered Banach spaces) probability measure and prove its exponential decay of correlations.
CMLs with multi-dimensional local systems of hyperbolic type have been studied by Ya.B. Pesin and Ya.G. Sinai [26], M. Jiang [17, 18], M. Jiang and A. Mazel [19], M. Jiang and Ya.B. Pesin [20] and D.L. Volevich [29, 30].
For detailed reviews on mathematical results on CMLs we refer to [2], [5], [7] and [20].

An interacting particle system (IPS) is given by an infinite lattice with a copy of the same state space (that is usually a finite or countable set but can also be a Riemannian manifold) at each site. The updating at an individual site is a deterministic or stochastic map (e.g. in the case of finite local state spaces it is given by a stochastic matrix with transition probabilities as its coefficients) that is applied with 'exponential waiting times', i.e. like the waiting times for jumps in a Poisson process. The jump rates for the updating depends also on the other sites. R.J. Glauber introduces in [14] (a special case of) the stochastic Ising model as a model for magnetism. The total state space $\{-1, +1\}^{\mathbb{Z}^d}$ represents the spins of the atoms at all sites. The rate for a flip of an individual spin depends on the states of the neighbour sites. F. Spitzer [27, 28] and R.L. Dobrushin [9, 10] study more general systems where the individual jump rates do not only depend on the nearest neighbours.
A basic problem is to establish the existence of infinite systems with asynchronous updatings. The infinitely many jumps in a finite time-interval cannot be 'listed', i.e. there is no order preserving bijection between the time-ordered set of jumps and $\mathbb{N}$.
R.L. Dobrushin obtains in [9] the infinite system as the limit of subsystems over finitely many sites.
By using a percolation argument T.E. Harris proves in [15] that for systems of finite range interaction and a sufficiently small time interval the history of an individual particle depends on only finitely many sites, and so he provides a natural definition of the infinite system. With probability 1 the set $\mathbb{Z}^d$ splits into finite clusters such that each site is affected at most by sites in the same cluster.
R. Holley shows in [16] for generators, corresponding to one-dimensional models, and T.M. Liggett in [23] for the d-dimensioanal case, that these operators generate, in fact, a semigroup, acting on continuous functions.
Here we have only mentioned different methods to establish the existence of the infinite systems. For detailed reviews on IPSs and results on invariant measures, mixing properties, phase transitions and applications to physics and other sciences we refer to [11] and [24].

In this paper we consider the infinite topological product $M = (S^1)^{\mathbb{Z}^d}$ and continuous updating maps for the individual coordinates that are of finite range or Lipschitz-continuous wrt all coordinates with a summable family of Lipschitz constants (cf. Section 2.2 for the definition). The times for the updatings at the individual sites are independently Poisson-distributed with the same constant rate $\lambda > 0$. For the finite range case we show that with probability 1 the updatings at any finite set of sites and any finite time-interval depend on only finitely many sites. Our proof uses time- and space-oriented percolation and is different from the one in [15]. This result provides a natural definition of the infinite dynamical system.
For the systems with infinite range interaction we show that with probability 1 it is well-defined as the net-limit of its finite-dimensional subsystems with arbitrary boundary conditions. We combine standard estimates for error growth with ideas from percolation theory. The limit of the corresponding Markov kernels, acting on continuous functions, exists and provides a definition of the infinite process, different from the widely used generator approach.
Our proofs still work if we replace S^1 by any compact Riemannian manifold or stochastic systems with

finite state spaces. The assumption of having the same constant jump rate at all sites is by no means essential and can be weakened to the case of upper bounded individual jump rates that depend on other states as well. However, we do not consider these generalizations in this paper.
In a setting similar to that of [13], i.e. for analytically coupled circle maps (uniformly expanding and analytic) on the $\mathbb{Z}^d$-lattice with weak, exponentially decaying interaction but with asynchronous updatings as described above, we define transfer operators for the Markov kernels of the infinite system. The operators act on the Banach space $\mathcal{H}_\vartheta$ (introduced in [13]) that includes measures whose finite-dimensional marginals have analytic, exponentially bounded densities. Using 'cluster-expansion'-like techniques, we represent these integral operators in terms of configurations and prove the existence and uniqueness (in $\mathcal{H}_\vartheta$) of an invariant probability measure and its exponential decay of correlations.
The paper is organized as follows. Section 1 provides definitions, notation and some propositions about stochastic processes and metric spaces. In Section 2 we define the infinite-dimensional systems for finite range (Section 2.1) and infinite range interaction (Section 2.2) and the corresponding Markov kernels (Section 2.3). In Section 3 we study the transfer operators for a specific class of interactions. In Section 4 we prove the mixing properties of the invariant measure (found in Section 3) wrt spatio-temporal shifts.

1 Basic Definitions and Examples

In this section we present definitions from probability theory and topology and also introduce most of the notation used in this paper. We have taken most definitions and statements on probability theory from [3].

Definition 1.1 $\mathbb{N}$ denotes the set of natural numbers including zero. Let $(E, \mathcal{A}_2)$ be a measurable space, $(\Omega, \mathcal{A}_1, P)$ a probability space and $(X_t)_{t\in I}$ a family (with index set $I \neq \emptyset$) of random variables on $(\Omega, \mathcal{A}_1, P)$ with values in E.

- Then $(\Omega, \mathcal{A}_1, P, (X_t)_{t\in I})$ is called a **stochastic process with values in** $(\mathbf{E}, \mathcal{A}_\mathbf{2})$.
- If $I = \mathbb{R}^{\geq 0}$, $[0, T]$ or $[0, T)$ for some $T > 0$ the process is called a **continuous time stochastic process**.
- For fixed $\omega \in \Omega$ the map $t \mapsto X_t(\omega)$ is called the **trajectory** of ω. It is also denoted by $X_\bullet(\omega)$.
- We consider the set $\mathbb{N}$ as measurable space with the discrete σ-algebra. For any set Λ we denote by $\mathbb{N}^\Lambda$ the product space, established with the product σ-algebra.

  A continuous time stochastic process with values in $\mathbb{N}^\Lambda$ and with index set I and P-a.a. of whose trajectories are non-decreasing (i.e. the functions $t \mapsto \pi_q \circ X_t(\omega)$ are non-decreasing for all $q \in \Lambda$ and P-a.a. $\omega \in \Omega$. 'π_q' denotes the projection on the qth coordinate.), is called a **counting process** with values in $\mathbb{N}^\Lambda$ We say that such a process is **of finite expectation** if for all $t \in I$ the random variable $\omega \mapsto \sum_{q\in\Lambda} \pi_q \circ X_t(\omega)$ has finite expectation.

Remark 1.1
1. We will also use the short-hand-notation $X_\bullet$ for a stochastic process if Ω, $\mathcal{A}_1$ and P are obvious from the context.
2. The term *path* seems to be more common than *trajectory* but we will denote something else later on by *path*.
3. Finite expectation means that with probability 1 there are only finitely many jumps (cf. Definition 1.2 below) in every finite time-interval.

Definition 1.2 (cf. [3]) Let $(\Omega, \mathcal{A}_1, P, (X_t)_{t\in I})$ be a continuous time counting process with values in $\mathbb{N}$ as in Definition 1.1 and $\omega \in \Omega$. We define

$$X_t^+(\omega) \stackrel{\text{def}}{=} \begin{cases} X_t(\omega) & \text{if} \quad I = [0,t] \\ \lim_{s \searrow t} X_s(\omega) & \text{otherwise} \end{cases} \tag{1}$$

$$X_t^-(\omega) \stackrel{\text{def}}{=} \begin{cases} \lim_{s \nearrow t} X_s(\omega) & \text{if} \quad t > 0 \\ X_0(\omega) & \text{if} \quad t = 0 \end{cases} \tag{2}$$

We say that $X_\bullet(\omega)$ **jumps** at time $t \geq 0$ if $X_t^-(\omega) < X_t^+(\omega)$. Then $X_t^+(\omega) - X_t^-(\omega)$ is called the **size of the jump**.
Let $X_\bullet(\omega)$ be a continuous time counting process with values in $\mathbb{N}^\Lambda$ and $\omega \in \Omega$. We say that $X_\bullet(\omega)$ jumps at time t and site $q \in \Lambda$ if $\pi_q \circ X_\bullet(\omega)$ jumps at t. Then we also say that ω jumps at (q,t).
We define the **jump set $\mathbf{\Lambda}(\omega, \mathbf{t})$ of ω at time $\mathbf{t}$** as the set of all $q \in \Lambda$ such that ω jumps at (q,t).

Definition 1.3 (cf. [3]) Let $I = \mathbb{R}^{\geq 0}$ or $I = [0,T)$ for some $T > 0$. A stochastic process $(\Omega, \mathcal{A}, P, (X_t)_{t \in I})$ with values in $\mathbb{N}$ is called **(normalized) Poisson process with parameter $\lambda > 0$** if the following holds:

1. The process has stationary and independent increments which for all $s < t \in I$ satisfy

$$P(\{\omega : X_t(\omega) - X_s(\omega) = n\}) = p_\lambda(t-s, n) \tag{3}$$

with

$$p_\lambda(t,n) \stackrel{\text{def}}{=} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \tag{4}$$

2. P-almost every trajectory $X_\bullet(\omega)$ is a right-continuous, increasing function having at most jumps of size 1.

3. At time 0 P-a.a. trajectories have value 0:

$$P(\omega : X_0(\omega) = 0) = 1 \tag{5}$$

Theorem 1.1 *(cf. [3], Satz 41.2) For any $\lambda > 0$ and I as in Definition 1.3 there exists a (normalized) Poisson process with parameter λ. Any two such processes are equivalent (i.e. if $X_\bullet^1$ and $X_\bullet^2$ are two such processes then for any finite sequence $t_1 < \ldots < t_n$ in I the projections $(X_{t_1}^1, \ldots, X_{t_n}^1)$ and $(X_{t_1}^2, \ldots, X_{t_n}^2)$ have the same distribution.)*

□

Definition 1.4 Let Λ be a nonempty set and $(\Omega_q, \mathcal{A}_q, P_q, (X_t^q)_{t \in I})_{q \in \Lambda}$ be a family of stochastic processes with values in $(E_q, \mathcal{A}^q)$, respectively. We set

$$\Omega \stackrel{\text{def}}{=} \prod_{q \in \Lambda} \Omega_q, \tag{6}$$

$$\tilde{\mathcal{A}} \stackrel{\text{def}}{=} \bigotimes_{q \in \Lambda} \mathcal{A}_q, \tag{7}$$

$$\tilde{P} \stackrel{\text{def}}{=} \bigotimes_{q \in \Lambda} P_q, \tag{8}$$

$$\mathcal{A} \stackrel{\text{def}}{=} \text{completion of } \tilde{\mathcal{A}} \text{ wrt } \tilde{P}, \tag{9}$$

$$P \stackrel{\text{def}}{=} \text{extension of } \tilde{P} \text{ to } \mathcal{A} \tag{10}$$

$$\text{and} \quad X_t \stackrel{\text{def}}{=} (X_t^q)_{q \in \Lambda}. \tag{11}$$

In (6) we mean the cartesian product of spaces, in (7) the product sigma-algebra and in (8) the produc measure.
Then the process $(\Omega, \mathcal{A}, P, (X_t)_{t\in I}$ with values in $(\prod_{q\in\Lambda} E_q, \bigotimes_{q\in\Lambda} \mathcal{A}^q)$ is called the **product of the family of processes**.

Remark 1.2 1. Products of stochastic processes as in Definition 1.4 exist. For example the existence of the non-completed product measure follows from standard measure theory (cf. [3].)

2. For non-empty, at most countable Λ and a family (indexed by Λ) of Poisson processes two such products $X^1_\bullet$ and $X^2_\bullet$ are equivalent because for all $q \in \Lambda$ the Poisson processes $\pi_q \circ X^1_\bullet$ and $\pi_q \circ X^2_\bullet$ are equivalent (cf. Theorem 1.1). It follows from the definition of the product σ-algebra $\bigotimes_{q\in\Lambda} \mathcal{A}^q$ that $X^1_\bullet$ and $X^2_\bullet$ are equivalent.

Definition 1.5 Let $\lambda > 0$ and Λ a nonempty, at most countable set. A **Poisson process on Λ with parameter λ** is the product of a family, indexed by Λ, of Poisson processes with parameter λ.

Remark 1.3 1. For $\lambda > 0$ the Poisson process on $\mathbb{Z}^d$ with parameter λ is clearly not of finite expectation. In fact, for any $t > 0$ there are P-almost surely infinitely many jumps in $[0, t]$, i.e.

$$P(\{\omega : \sum_{q\in\mathbb{Z}^d} \pi_q \circ X_t(\omega) = \infty\}) = 1. \tag{12}$$

2. But if $\Lambda_1 \subset \mathbb{Z}^d$ is finite then $\pi_{\Lambda_1} \circ X_\bullet(\omega)$ has finitely many jumps in $[0, t]$ for P-a.a. $\omega \in \Omega$ and any $t > 0$.

3. There are P-almost surely no simultaneous jumps at two different sites:

$$P(\{\omega : \exists q_1 \neq q_2 \in \mathbb{Z}^d, t \geq 0 \text{ such that } \omega \text{ jumps at } (q_1, t) \text{ and } (q_2, t)\}) = 0. \tag{13}$$

4. For $0 \leq t_0 < t$

$$P(\{\omega : \omega \text{ jumps at } t_0\}) = 0. \tag{14}$$

Proof of Remark 1.3 We only show (13). The proofs of the other statements are similar. We set

$$A(q_1, q_2, T) \stackrel{\text{def}}{=} \{\omega : \exists t \in [0, T) \text{ such that } \omega \text{ jumps at } (q_1, t) \text{ and } (q_2, t)\}. \tag{15}$$

We have to prove that the set

$$\bigcup_{T\in\mathbb{N}} \bigcup_{q_1,q_2\in\mathbb{Z}^d} A(q_1, q_2, T) \tag{16}$$

has P-measure zero and it is sufficient to show that

$$P(A(q_1, q_2, T)) = 0 \tag{17}$$

for fixed $q_1 \neq q_2 \in \mathbb{Z}^d$ and $T > 0$. For this we set

$$I_{N,k} \stackrel{\text{def}}{=} \left[(k-1)\frac{T}{N}, k\frac{T}{N}\right) \tag{18}$$

for $N \in \mathbb{N} \setminus \{0\}$ and $1 \leq k \leq N$. By (4) we have for $i = 1, 2$:

$$P\left(\{\omega : \text{ jumps at } (q_i, t) \text{ for some } t \in I_{N,k}\}\right) = 1 - e^{-\lambda\frac{T}{N}} \tag{19}$$

and so, using the estimate $e^x \geq 1 + x$:

$$\begin{aligned} & P\left(\{\omega : \exists k; t_1, t_2 \in I_{N,k} \text{ such that } \omega \text{ jumps at } (q_1, t_1) \text{ and } (q_2, t_2)\}\right) \\ & \leq \quad N \left(1 - e^{-\lambda \frac{T}{N}}\right)^2 \\ & \leq \quad \lambda^2 T^2 \frac{1}{N} \end{aligned} \tag{20}$$

which converges to 0 as $N \to \infty$.

□

The following two definitions prepare Definition 1.8 that we will need in Section 2.

Definition 1.6 In view of Definition 1.3 and Remark 1.3 we define (for a given Poisson process like in that remark) the set $\mathcal{N}_1$ of P-measure zero:

$$\begin{aligned} \mathcal{N}_1 \quad \stackrel{\text{def}}{=} \quad & \{\omega : X_\bullet(\omega) \text{ is not non-decreasing, has jumps at 0,} \\ & \qquad \text{simultaneous jumps or jumps of size greater than 1}\}. \end{aligned} \tag{21}$$

Definition 1.7 For $q = (q_1, \ldots, q_n) \in \mathbb{Z}^d$ we define

$$\|q\| \stackrel{\text{def}}{=} |q_1| + \ldots + |q_n|. \tag{22}$$

For $R \geq 0$

$$B_R(q) \stackrel{\text{def}}{=} \{\tilde{q} \in \mathbb{Z}^d : \|q - \tilde{q}\| \leq R\} \tag{23}$$

is the set of points that have distance at most R from q.

Definition 1.8 Let $a, b \in \mathbb{Z}^d$ and $n \geq 0$. A **path from a to b** is a finite sequence $Q = (q_0 = a, q_1, \ldots, q_n = b)$ of points $q_i \in \mathbb{Z}^d$. We call $\max_{0 \leq i \leq n-1} \|q_{i+1} - q_i\|$ the **step size** of Q. Note the special case of a path $Q = (q_0)$. It is called the **empty path at site $\mathbf{q_0}$** and we define its step size to be 0.

Definition 1.9 Let $(\Omega, \mathcal{A}, P, (X_t)_{t \geq 0})$ be a Poisson process with parameter $\lambda > 0$ and with values in $\mathbb{N}^{\mathbb{Z}^d}$. Let $T > 0$, $\omega \in \Omega$ and $Q = (q_0 = a, q_1, \ldots, q_n = b)$ a path. We extend Q to the infinite sequence $\widetilde{Q} = (q_0, q_1, \ldots, q_n, q_{n+1} = q_n, \ldots)$ in which q_n is repeated.
We define a process $(\Omega, \mathcal{A}, P, (Z_t)_{t \in [0,T]})$ with values in $\mathbb{N}$ as follows.

$$\begin{aligned} Z : [0, t] \times \Omega \quad & \to \quad \mathbb{N} \\ (t, \omega) \quad & \mapsto \quad Z_t(\omega) \end{aligned} \tag{24}$$

If $\omega \in \mathcal{N}_1$ or it does not jump at (q_0, t) for any $t \in (0, T)$ we set $Z_\bullet(\omega) = 0$ on $[0, T]$.
Otherwise there is a maximal sequence

$$t_{-1} \stackrel{\text{def}}{=} T > t_0 > t_1 > \ldots > t_{m(\omega)} \tag{25}$$

such that

$$t_i \stackrel{\text{def}}{=} \max\{t \in (0, t_{i-1}) : \omega \text{ jumps at } (q_i, t)\} \quad \text{for} \quad 0 \leq i \leq m(\omega). \tag{26}$$

'Maximal' means that ω does not jump at $q_{m(\omega)+1}$ in the time interval $(0, t_{m(\omega)})$ and so the sequence (25) cannot be extended. (Intuitively one can think that one sits at time T at site q_0 and, going *backwards*

in time, waits for the next jump of ω at q_0 (which happens at time t_0), then jumps (instantly) to q_1 and waits (*backwards* in time) for the next jump of ω at q_1, then jumps to q_2 etc. After n jumps (should this occur) one does not change the sites any more, but possibly jumps from q_n to q_n. $m(\omega)$ is the total number of jumps. It is P-a.s. finite because P-a.a. ω have only finitely many jumps at q_n.)
We set for $t \in [0,T]$:

$$\widetilde{Z}_t(\omega) \stackrel{\text{def}}{=} \begin{cases} i & \text{for} \quad t \in [t_i, t_{i-1}) \\ m(\omega) & \text{for} \quad t \in [0, t_{m(\omega)}] \end{cases} \tag{27}$$

And $Z_\bullet(\omega)$ is the (uniquely defined) right-continuous function, such that $Z_\bullet(\omega) = \widetilde{Z}_{T-\bullet}(\omega)$ everywhere, except possibly where these functions jump. Then $(\Omega, \mathcal{A}, P, (Z_t)_{t\in[0,T]})$ is a Poisson process with parameter λ. (A precise proof of this uses that the constructed process is 'made of' independent Poisson processes and that these have independent increments.) We call it the **Poisson process induced by the path Q**.

Definition 1.10 In the setting of Definition 1.9 we call Q a **causal path wrt** $(\mathbf{t}, \omega)$ if $Z_T(\omega) \geq n$ and a **maximal causal path wrt** $(\mathbf{t}, \omega)$ if $Z_T(\omega) = n$. (The latter means that $Q = (q_0, \ldots, q_n)$ cannot be extended to any causal path $(q_0, \ldots, q_n, q_{n+1})$.)
We define:

- $\mathbf{Path(q, n, R)}$ to be the set of paths that start at q, have exactly n steps and are of step size at most R.
- $\mathbf{Path(q \to \Lambda)}$ for any $\emptyset \neq \Lambda \in \mathbb{Z}^d$ to be the set of paths starting at q and ending in Λ.
- $\mathbf{Path_C(t, \omega, q, \Lambda)}$ for $q \in \Lambda$ to be the set of causal wrt (t, ω) paths $Q = (q_0 = q, \ldots, q_n)$ such that
  1. Q is maximal causal and $q_0, \ldots, q_n \in \Lambda$
  or 2. $q_0, \ldots, q_{n-1} \in \Lambda$ and $q_n \in \Lambda^C$.
- $\mathbf{Path_C(t, \omega, q \to \Lambda^C)}$ for $q \in \Lambda$ to be the set of causal paths $(q_0 = q, \ldots, q_n)$ such that $q_0, \ldots, q_{n-1} \in \Lambda$ and $q_n \in \Lambda^C$. (So this is the subset of elements in $\mathbf{Path_C}(t, \omega, q, \Lambda)$ for which case 2. applies.)

Remark 1.4
1. We have defined the property of being causal for general paths and not related this definition to any kind of interaction. When we study finite range interaction, of range R say, we will consider only causal paths of step size at most R.
2. A term like *inverse causal path* from a to b instead of *causal path* would actually be more appropiate as it corresponds to b affecting a (cf. Definition 2.1) but not necessarily the other way around. However, we prefer the shorter notion.

Definition 1.11 (cf. [3]) Let $(\Omega_1, \mathcal{A}_1)$ and $(\Omega_2, \mathcal{A}_2)$ be measurable spaces. A map $K : \Omega_1 \times \mathcal{A}_2 \to [0,1]$ is called a **Markov kernel from** $(\mathbf{\Omega_1}, \mathcal{A}_1)$ **to** $(\mathbf{\Omega_2}, \mathcal{A}_2)$ if the two following conditions are satisfied:

MK1 $\omega_1 \mapsto K(\omega_1, A_2)$ is $\mathcal{A}_1$-measurable for all $A_2 \in \mathcal{A}_2$.
MK2 $A_2 \mapsto K(\omega_1, A_2)$ is a probability measure on $\mathcal{A}_2$ for all $\omega_1 \in \mathcal{A}_1$.

If $(\Omega_1, \mathcal{A}_1) = (\Omega_2, \mathcal{A}_2)$ then K is called a **Markov kernel on** $(\mathbf{\Omega_1}, \mathcal{A}_1)$.

Example 1.1 Let (Y, ϱ_Y) be a metric space and $\mathcal{B}_Y$ its Borel σ-algebra. $\mathcal{C}^0(Y,Y)$ is the space of continuous maps from Y to Y. It has a uniform metric, defined by $\varrho_{\mathcal{C}^0(Y,Y)}(g_1, g_2) = \sup_{\mathbf{y}\in\mathbf{Y}} \varrho_Y(g_1(\mathbf{y}), g_2(\mathbf{y}))$ and the Borel σ-algebra $\mathcal{B}_{\mathcal{C}^0(Y,Y)}$ wrt this metric. Further let $(\Omega, \mathcal{A}, P)$ be a probability space and

$$\begin{array}{rcl} S : \Omega & \to & \mathcal{C}^0(Y,Y) \\ \omega & \mapsto & S_\omega \end{array} \tag{28}$$

a measurable (wrt the σ-algebras $\mathcal{A}$ and $\mathcal{B}_{\mathcal{C}^0(Y,Y)}$) map.
Then

$$K_S(y, Y_1) \stackrel{\text{def}}{=} P(\{\omega : S_\omega(y) \in Y_1\}) \tag{29}$$

for all $y \in Y, Y_1 \in \mathcal{B}_Y$, defines a Markov kernel on $(Y, \mathcal{B}_Y)$.

Proof To verify **MK1** we fix an $Y_1 \in \mathcal{B}_Y$ and show that the map $y \mapsto K_S(y, Y_1)$ is measurable. First we note that S can be seen as a measurable map from $\Omega \times Y$ to Y. We write it as the composite of measurable maps $S \times \mathrm{id}_Y$ and the 'evaluation map':

$$(\omega, y) \mapsto (S_\omega, y) \mapsto S_\omega(y). \tag{30}$$

The map $S \times \mathrm{id}_Y$ is measurable by assumption and the definition of the product σ-algebra of $\mathcal{C}^0(Y, Y) \times Y$. The evaluation map is continuous (wrt to the product topology), hence measurable wrt the Borel σ-algebras. So the composite in (30) is measurable in $\Omega \times M$. It follows that the map $y \mapsto P(\{\omega : S_\omega(y) \in Y_1\})$ is measurable (cf. Lemma 8.1 on p. 159 in [22]) and so **MK1** holds.
Next we show **MK2**. Consider for fixed $y \in Y$ the composite of measurable maps

$$\omega \mapsto (\omega, y) \mapsto S_\omega(y) \tag{31}$$

that maps Ω to Y. We see that $K(y, \cdot)$ is the image of P wrt this map and so a probability measure which was to be shown.

□

Definition 1.12 (cf. [3]) Let K be a Markov kernel from $(\Omega_1, \mathcal{A}_1)$ to $(\Omega_2, \mathcal{A}_2)$ and $E^*(\mathcal{A}_i)$ $(i = 1, 2)$ the set of $\mathcal{A}_i$-measurable functions with values in $[0, \infty]$. Then K defines a map from $E^*(\mathcal{A}_2)$ to $E^*(\mathcal{A}_1)$ as follows:

$$(Kf)(\omega_1) \stackrel{\text{def}}{=} \int_{\Omega_2} K(\omega_1, d\omega_2)\, f(\omega_2) \tag{32}$$

for any $f \in E^*(\mathcal{A}_2)$.

Example 1.2 (cf. [3]) For the characteristic function χ_{A_2} of an $\mathcal{A}_2$-measurable set A_2 we get

$$K\chi_{A_2}(\omega_1) = K(\omega_1, A_2). \tag{33}$$

Now we consider a special case of Example 1.1.

Example 1.3 Let $S : Y \to Y$ be a continuous map map on (Y, ϱ_Y) and let $(\Omega, \mathcal{A}, P, (X_t)_{t \in I})$ be a counting process with values in $\mathbb{N}$ and $t \in I$.
The map

$$\begin{aligned} S_\omega^t : Y &\to Y \\ y &\mapsto S^{X_t(\omega)}(y), \end{aligned} \tag{34}$$

where $S^{X_t(\omega)}$ denotes the $X_t(\omega)$-th iterate of S, is well-defined for all $\omega \in \Omega$. Further $S_\omega(y)$ is measurable wrt (ω, y). In fact, S_ω depends just on $X_t(\omega)$ and so we get a countable, measurable partition of Ω:

$$\Omega = \bigcup_{n \in \mathbb{N}} U(n) \tag{35}$$

$$\text{with} \quad U(n) \stackrel{\text{def}}{=} \{\omega \in \Omega : X_t(\omega) = n\} \tag{36}$$

We define a Markov kernel by

$$\begin{aligned} K_S^t(y, Y_1) &\stackrel{\text{def}}{=} P(\{\omega : S_\omega^t(y) \in Y_1\}) \\ &= \sum_{n: S^n(y) \in Y_1} P(U(n)) \\ &= \int_\Omega dP(\omega)\, \chi_{Y_1} \circ S_\omega^t(y) \end{aligned} \tag{37}$$

for $y \in Y$ and $Y_1 \in \mathcal{B}_Y$.

We prepare a generalization of Example 1.3 with a definition and a technical lemma.

Definition 1.13 Let $\mathcal{F}$ be the set of finite subsets of $\mathbb{Z}^d$. Consider a fixed $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$. We define $\mathcal{J}$ to be the union of a one-point set $\{j_\infty\}$ and the set of finite sequences $(\Lambda_1, \ldots, \Lambda_n)$ of subsets of Λ. Then $\mathcal{J}$ is countable and we consider it as a measurable space, established with the discrete σ-algebra.
Let $(\Omega, \mathcal{A}, P, (X_t)_{t \in I})$ be a discrete or continuous time counting process with values in $\mathbb{N}^\Lambda$ and index-set $I = \{1, \ldots, N\}$ or $[0, T]$, respectively. We define a map

$$\begin{aligned} \mathbf{j} : \Omega &\to \mathcal{J} \\ \omega &\mapsto \mathbf{j}(\omega) \end{aligned} \tag{38}$$

If $X_\bullet(\omega)$ is non-decreasing, has only finitely many jumps and at most jumps of size 1 then we define $\mathbf{j}(\omega)$ to be the (time-ordered) sequence of jump sets of ω. Otherwise we set $\mathbf{j}(\omega) = j_\infty$. We define for $j \in \mathcal{J}$:

$$U(j) \stackrel{\text{def}}{=} \{\omega : \mathbf{j}(\omega) = j\} \tag{39}$$

Lemma 1.1 *Let $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$ be fixed and $(\Omega, \mathcal{A}, P, (X_t)_{t \in I})$ a continuous time counting process with index-set $I = [0, T]$ and values in $\mathbb{N}^\Lambda$ such that for P-a.a. ω the trajectory $X_\bullet(\omega)$ is non-decreasing, has only finitely many jumps and at most jumps of size 1. Then the map $\mathbf{j}$, as defined in Definition 1.13, is measurable.*

Proof We consider the case $I = [0, T]$. By assumption $\mathcal{N} = U(j_\infty)$ is measurable and has measure zero. We have to show that $U(j)$ is measurable for any $j = (\Lambda_1, \ldots, \Lambda_n)$. For any $q_1, q_2 \in \Lambda$, $n_1, n_2 \in \mathbb{N} \setminus \{0\}$ and $t \in [0, T]$ we define $A_1(q_1, n_1, q_2, n_2)$ to be the set of all $\omega \in \Omega \setminus \mathcal{N}$ that have at least n_1 jumps at q_1 and at least n_2 jumps at q_2 and the n_1th jump at q_1 happens at the same time as the n_2th jump at q_2. Similarly $A_2(q_1, n_1, q_2, n_2)$ is the set of all $\omega \in \Omega \setminus \mathcal{N}$ that have at least n_1 jumps at site q_1 and the n_1th jump at q_1 occurs before the n_2th jump (if any) at q_2. We only show the measurability of the sets $A_2(\cdot)$. The proof of the measurability of the sets $A_1(\cdot)$ uses similar arguments.

$$A^{\geq}(q_1, n_1, t) \stackrel{\text{def}}{=} \{\omega \in \Omega \setminus \mathcal{N} : \pi_{q_1} \circ X_t(\omega) \geq n_1\} \tag{40}$$

is the set of all $\omega \in \Omega \setminus \mathcal{N}$ that have at least n_1 jumps at site q_1 and the n_1th of these jumps happens at the latest at time t. We define the set $A^{<}(q_2, n_2, t)$ analogously.
$A^{\geq}(q_1, n_1, t)$ and $A^{<}(q_2, n_2, t)$ are measurable, and so is $A_2(q_1, n_1, q_2, n_2)$ since

$$A_2(q_1, n_1, q_2, n_2) = \bigcup_{t \in [0,T] \cap \mathbb{Q}} \left(A^{\geq}(q_1, n_1, t) \cap A^{<}(q_2, n_2, t) \right). \tag{41}$$

Now ω belongs to $U(j)$ if and only if, for all $1 \leq k \leq n$ and $q_1, q_2 \in \Lambda_k$ and $q_3 \in \Lambda \setminus \Lambda_k$ the following holds:

- If for exactly n_1 indices $1 \leq i \leq k$ the point q_1 belongs to Λ_i and for exactly n_2 indices $1 \leq j \leq k$ the point q_2 belongs to Λ_j then $\omega \in A_1(q_1, n_1, q_2, n_2)$.

- If for exactly n_1 indices $1 \leq i \leq k$ the point q_1 belongs to Λ_i and for exactly $n_3 - 1$ indices $1 \leq j < k$ the point q_3 belongs to Λ_j then $\omega \in A_2(q_1, n_1, q_3, n_3)$.
- If for exactly $l \geq 0$ indices $1 \leq n_1 < n_2 < \ldots < n_l \leq n$ a point $q \in \Lambda$ belongs to Λ_{n_i} then $\omega \in \{\tilde{\omega} \in \Omega \setminus \mathcal{N} : \pi_q \circ X_T(\tilde{\omega}) = l\}$.

We see that $U(j)$ is the intersection of finitely many measurable sets and hence measurable.

□

Example 1.4 We consider a generalization of Example 1.3. Let (Y, ϱ_Y) be a measurable space, Λ a non-empty finite set and $(\Omega, \mathcal{A}, P, (X_t)_{t\in I})$ a counting process with values in $\mathbb{N}^\Lambda$ that has finite expectation and with P-almost surely only jumps of size at most 1. Let $S = (S_{\Lambda_1})_{\Lambda_1 \subset \Lambda}$ be a family of continuous maps on Y^Λ, such that S_{Λ_1} changes at most the Λ_1-coordinates, i.e. if $\mathbf{y}_\Lambda \in Y^\Lambda$ and $q \in \Lambda \setminus \Lambda_1$ we have for the qth coordinate $\pi_q \circ S_{\Lambda_1}(\mathbf{y}_\Lambda) = y_q$.
For $t \in I$ and P-a.a. $\omega \in \Omega$ with $X_t(\omega) \in \mathbb{N}^\Lambda$ we have a finite sequence of jump-sets $\mathbf{j}(\omega) = (\Lambda_1, \ldots, \Lambda_n)$, as defined in Definition 1.13, and it depends measurably on ω, as was shown in Lemma 1.1. We define

$$S^t_\omega : \quad Y^\Lambda \quad \to Y^\Lambda \tag{42}$$

$$\mathbf{y}_\Lambda \quad \mapsto S_{\mathbf{j}(\omega)}(\mathbf{y}_\Lambda) \stackrel{\text{def}}{=} S_{\Lambda_n} \circ \ldots \circ S_{\Lambda_1}(\mathbf{y}_\Lambda) \tag{43}$$

We get a representation of $K^t_S(\mathbf{y}_\Lambda, Y_1)$, similar to the one in (37):

$$\begin{aligned} K^t_S(\mathbf{y}_\Lambda, Y_1) &= P(\{\omega : S^t_\omega(\mathbf{y}_\Lambda) \in Y_1\}) \\ &= \int_\Omega dP(\omega)\, \chi_{Y_1} \circ S^t_\omega(\mathbf{y}_\Lambda) \\ &= \sum_{j \in \mathcal{J} : S_j(\mathbf{y}_\Lambda) \in Y_1} P(U(j)) \end{aligned} \tag{44}$$

for $\mathbf{y}_\Lambda \in Y^\Lambda$ and $Y_1 \in \bigotimes_{q\in\Lambda} \mathcal{Y}$.

We have seen in Example 1.4 that S^t_ω depends on $\mathbf{j}(\omega)$ only.
As we are interested in spatially extended systems we need some definitions and facts about infinite-dimensional systems.

Definition 1.14 S^1 is the one-dimensional sphere. We define it to be isometric as Riemannian manifold to $\mathbb{R}/2\pi\mathbb{Z}$. This defines in particular a metric ϱ_{S^1} on S^1 and also the normalized Lebesgue measure on the (completed) Borel σ-algebra.
The diameter of S^1 is

$$c_s \stackrel{\text{def}}{=} \operatorname{diam}_{\varrho_{S^1}}(S^1) = \pi. \tag{45}$$

(It seems a bit redundant to introduce the constant c_S instead of using π in the following. But we indicate that the proofs in Section 2 work if S^1 is replaced by any compact Riemannian manifold or more general by a bounded metric space with a Borel probability measure. Further we use the letter 'π' as notation for projections.)
We set

$$M \stackrel{\text{def}}{=} (S^1)^{\mathbb{Z}^d} \tag{46}$$

and give it the product topology and product Lebesgue measure on the (completed) Borel σ-algebra.

For $\Lambda \subset \mathbb{Z}^d$ we denote by π_Λ the projection on the Λ-coordinates.
Note that the product of the Borel σ-algebras is the same as the Borel σ-algebra for the product space. M is compact and metrizable in the following way:

Definition 1.15 Let $(b(q))_{q\in\mathbb{Z}^d}$ be a family of positive numbers such that

$$\lim_{R\to\infty} \sup_{\|q\|\geq R} b(q) = 0. \tag{47}$$

Then the metric ϱ_M on M, associated to $(b(q))_{q\in\mathbb{Z}^d}$, is defined by

$$\varrho_M(\mathbf{x},\mathbf{y}) \stackrel{\text{def}}{=} \sup_{q\in\mathbb{Z}^d} b(q)\, \varrho_{S^1}(x_q, y_q) \tag{48}$$

for $\mathbf{x},\mathbf{y} \in M$.

Remark 1.5

1. One can easily show that ϱ_M, as defined in Definition 1.15, is in fact a metric and also compatible with the product topology.
2. A sequence $(\mathbf{x}^{(n)})_{n\in\mathbb{N}}$ in M converges wrt the product topology iff it converges wrt to each coordinate, i.e. $(x_q^{(n)})_{n\in\mathbb{N}}$ converges for every $q \in \mathbb{Z}^d$. The same holds also for nets $(\mathbf{x}^\Lambda)_{\Lambda\in\mathcal{F}}$.
3. The product topology does not distinguish any particular sites despite the the fact that the weights $b(q)$ depend on q. Spatial shifts, like $\mathbf{x} \mapsto \tilde{\mathbf{x}}$ with $\tilde{x}_q = x_{q-r}$ for some $r \in \mathbb{Z}^d$, are homeomorphisms.
4. The space $\mathcal{C}^0(M,M)$ of continuous maps on (M,ϱ_M) is complete wrt the metric defined by

$$\varrho_{\mathcal{C}^0(M,M)}(f,g) \stackrel{\text{def}}{=} \sup_{\mathbf{x}\in M} \varrho_M(f(\mathbf{x}), g(\mathbf{x})). \tag{49}$$

We denote by $\mathcal{B}_{\mathcal{C}^0(M,M)}$ the Borel σ-algebra wrt this metric.

Lemma 1.2 *Let $(\Omega,\mathcal{A})$ be a measurable space and $(f^\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ be a net of measurable maps*

$$\begin{aligned} f^\Lambda : \Omega &\to \mathcal{C}^0(M,M) \\ \omega &\mapsto f^\Lambda_\omega \end{aligned} \tag{50}$$

such that for all $\Lambda_1 \in \mathcal{F}\setminus\{\emptyset\}$ and $\omega \in \Omega$ the net $(\pi_{\Lambda_1}\circ f^\Lambda_\omega)_{\Lambda_1\subset\Lambda\in\mathcal{F}}$ converges (as $\Lambda\to\mathbb{Z}^d$) in $\mathcal{C}^0(M,(S^1)^{\Lambda_1})$, say to $\pi_{\Lambda_1}\circ f_\omega$.
Then

$$f_{\omega,q}(\mathbf{x}) \stackrel{\text{def}}{=} \lim_{\Lambda\to\mathbb{Z}^d} \pi_q \circ f^\Lambda_\omega(\mathbf{x}) \tag{51}$$

defines a measurable map

$$\begin{aligned} f : \Omega &\to \mathcal{C}^0(M,M) \\ \omega &\mapsto f_\omega \end{aligned} \tag{52}$$

whose qth coordinate function is given by (51).

Proof Fix $\omega \in \Omega$, $\mathbf{x} \in M$ and a metric ϱ_M like in Definition 1.15 . We show that f_ω is continuous in $\mathbf{x}$. For that let $\epsilon > 0$ and choose $R_0 \in \mathbb{N}$ such that

$$c_s \, b(q) < \epsilon \tag{53}$$

for all q with $\|q\| > R_0$. We note that the qth coordinate function of $\pi_{\Lambda_1} \circ f_\omega \in \mathcal{C}^0(M, (S^1)^{\Lambda_1})$ is the same as the qth coordinate function $f_{\omega,q}$ of f_ω.
By continuity of $\pi_{B_{R_0}} \circ f_\omega$ we can choose a $\delta > 0$ such that for all $\mathbf{y} \in B_\delta(\mathbf{x})$ and all q with $\|q\| \leq R_0$:

$$c_s \, b(q) \, \varrho_{S^1}(f_{\omega,q}(\mathbf{x}), f_{\omega,q}(\mathbf{y})) < \epsilon. \tag{54}$$

From (53) and (54) we conclude that for all $\mathbf{y} \in B_\delta(\mathbf{x})$

$$\varrho_M(f_\omega(\mathbf{x}), f_\omega(\mathbf{y})) < \epsilon \tag{55}$$

which was to be shown. Finally f depends measurably on ω because it is pointwise limit of measurable functions with values in a metric space (cf. [22], p 117, for example).

□

Remark 1.6

1. Lemma 1.2 is in particular based on the compactness on M wrt the product toplology. M is not compact wrt the different metric, defined by

$$\tilde{\varrho}_M(\mathbf{x}, \mathbf{y}) \stackrel{\text{def}}{=} \sup_{q \in \Lambda} \varrho_{S^1}(x_q, y_q).$$

   In this case the conclusion from 'local' to 'global' does not hold.

2. As f in (52) is $(\mathcal{A}, \mathcal{B}_{\mathcal{C}^0(M,M)})$-measurable, the map $(\omega, \mathbf{x}) \mapsto f_\omega(\mathbf{x})$ is $(\mathcal{A} \times \mathcal{B}_M, \mathcal{B}_M)$-measurable. We have proved this fact in Example 1.1.

2 Infinite-Dimensional Systems

In Example 1.4 we used a counting process with values in $\mathbb{N}^\Lambda$ (for finite Λ) and a family of updating-maps on Y^Λ to define Markov kernels on the product Y^Λ. These kernels act on the product space $\mathcal{C}^0(Y^\Lambda)$ of continuous functions (cf. Definition 1.12 and Proposition 2.4). In view of spatially extended systems like coupled map lattices or interacting particle systems we would like to define analogous operators for infinite-dimensional systems ($\Lambda = \mathbb{Z}^d$). As counting process we take the Poisson process $(\Omega, \mathcal{A}, P, (X_t)_{t \geq 0})$ with parameter $\lambda > 0$ and values in $\mathbb{N}^{\mathbb{Z}^d}$.
Recall that the set $\mathcal{N}_1$, defined in Definition 1.6, of all $\omega \in \Omega$ such that $X_\bullet(\omega)$ is not nondecreasing, jumps at time 0, has simultaneous jumps or jumps of size greater than one, has P-measure zero. So we have to consider updatings only at single sites. They are given by a family of continuous maps $(S_q)_{q \in \mathbb{Z}^d}$ such that $S_q : M \to M$ changes only the qth coordinate (cf. Example 1.4 for a definition.) We remark that such a family naturally gives rise to updatings at more than one point at the same time.
A problem is obviously that the Poisson process, restricted to any finite interval $[0, t]$ of length $t > 0$ is not of finite expectation (cf. Definition 1.1 and Remark 1.3.1). P-a.s. there are infinitely many jumps and it is even impossible to define an order preserving bijection between them and $\mathbb{N}$. However, in Subsection 2.1 we will show for systems with finite range interaction that for P-a.a. $\omega \in \Omega$, any $q \in \mathbb{Z}^d$ and $t > 0$ the site q is affected in $[0, t]$ (cf. Definition 2.1) by only finitely many sites, so that maps '$\pi_q \circ S^t_\omega$' from M to $(S^1)^{\{q\}}$ and then also 'S^t_ω' from M to M can be defined in a natural way. The proof is based on a percolation argument. Percolation techniques, but different from the ones presented here, were already used by Harris in [15] for proving the existence of certain interacting particle systems of finite range. It follows in particular that $\pi_\Lambda \circ S^t_\omega : M \to (S^1)^\Lambda$ for finite $\Lambda \neq \emptyset$ is the limit (as $\tilde{\Lambda} \to \mathbb{Z}^d$) of maps

that are constructed by using the 'cut offs' $\pi_\Lambda \circ S^t_{\tilde\Lambda,\xi,\omega}$, corresponding to a finite $\tilde\Lambda \supset \Lambda$ and boundary conditions ξ. In fact, this limit also exists and is independent of the boundary conditions for a huge class of infinite range interactions as we will show in Subsection 2.2. It gives rise to a natural definition of the system. But we also note that for infinite range interaction each site is with positive probability affected by infinitely many other sites. So we cannot use the same definition as for finite range interaction.
In Section 2.3 we define Markov kernels K^t_S for the infinite system S^t and $K^t_{S,\tilde\Lambda}$ for the system $S^t_{\tilde\Lambda}$ that fixes the $\tilde\Lambda^C$-coordinates for a finite $\tilde\Lambda$. We show that K^t_S is the weak limit of $K^t_{S,\tilde\Lambda}$ (as $\tilde\Lambda \to \mathbb{Z}^d$), i.e. the corresponding operators on continuous functions converge weakly.

2.1 Finite Range Interaction

Now we are considering an interaction of range $R \in \mathbb{N} \setminus \{0\}$, i.e. $\pi_q \circ S_q(\mathbf{x})$ depends only on $\mathbf{x}_{B_R(q)}$. (Recall that $B_R(q)$ was defined in (23).)

Definition 2.1 Given R as above, $q, \tilde q \in \mathbb{Z}^d$, $T > 0$, $\omega \in \Omega$. We say that $\tilde{\mathbf{q}}$ **affects** $\mathbf{q}$ **wrt** $(\mathbf{R}, \mathbf{t}, \omega)$ if there is a causal path from q to $\tilde q$ of step size at most R. (Recall that we defined *path* etc. in Definitions 1.8 to 1.10). If $\emptyset \neq \Lambda \subset \mathbb{Z}^d$ we say that $\tilde{\mathbf{q}}$ **affects** $\mathbf{\Lambda}$ **wrt** $(\mathbf{R}, \mathbf{t}, \omega)$ if $\tilde q$ affects at least one point in Λ wrt (R, t, ω).
We set

$$\mathrm{Aff}_{(R,t,\omega)}(\Lambda) \stackrel{\text{def}}{=} \{\tilde q \in \mathbb{Z}^d : \tilde q \text{ affects } \Lambda \text{ wrt } (R,t,\omega)\} \tag{56}$$

$$\text{and} \quad \Omega_R \stackrel{\text{def}}{=} \{\omega : \exists t > 0, q \in \mathbb{Z}^d \text{ such that } |\mathrm{Aff}_{(R,t,\omega)}(q)| = \infty\} \tag{57}$$

where $|\cdot|$ denotes the cardinality.



Figure 1: The history of q_0

Figure 1 is a picture of $\mathrm{Aff}_{(1,T,\omega)}(q_0)$. We consider the finite time-interval $(0, T]$ and nearest neighbour interaction and a particular ω. For each jump we draw a cross at the particular point (q, t). There are jumps at (q_2, t_6), (q_0, t_5), (q_{-2}, t_4), (q_1, t_3), (q_0, t_2) and (q_3, t_1) The last jump at q_0 is at time t_2. We draw a thick horizontal line between (q_0, t_2) and (q, t_2) for all nearest neighbours q of q_0 because the updating of q_2 depends also on these sites. So we have to consider the 'histories' of q_0 *and* its nearest neighbours *before* time t_2. Note that $q_3 \in \mathrm{Aff}_{(1,T,\omega)}(q_0)$ and it is updated at time t_1 (and so affected by q_4 for example) but that updating has no influence on q_0 (at time T). We also note that, for example, q_{-1} affects q_0 (wrt $(1, T, \omega)$) but not the other way around. So we have to consider only the time- and space-ordered percolation.

Proposition 2.1 *Ω_R has P-measure zero:*

$$P(\Omega_R) = 0. \tag{58}$$

Proof $\mathrm{Aff}_{(R,t,\omega)}(q)$ is increasing in t and so

$$\Omega_R = \bigcup_{t\in\mathbb{N}} \bigcup_{q\in\mathbb{Z}^d} \{\omega : |\mathrm{Aff}_{(R,t,\omega)}(q)| = \infty\}. \tag{59}$$

So it is sufficient to show that for fixed $q \in \Lambda$ and $t > 0$ the set $\{\omega : |\mathrm{Aff}_{(R,t,\omega)}(q)| = \infty\}$ has P-measure zero. If we set

$$A_N \stackrel{\mathrm{def}}{=} \{\omega : \mathrm{Aff}_{(R,t,\omega)}(q) \not\subset B_N(q)\} \tag{60}$$

it is sufficient to show that

$$\lim_{N\to\infty} P(A_N) = 0. \tag{61}$$

If q is affected by some $\tilde{q} \notin B_N(q)$ wrt (R, t, ω) then there is a maximal causal path of step size at most R from q to $\tilde{q}$ with at least N_0 steps, where N_0 is the smallest integer greater than $\frac{N}{R}$.
Consider any maximal causal path $Q = (q_0 = q, \ldots, q_n)$ of step size at most R and with $n \geq N_0$. Q is a maximal causal path wrt (t, ω) iff the trajectory of ω wrt the Poisson process induced by Q (cf. Definition 1.9) has exactly n jumps. The probability of this is $p_\lambda(n, t)$ (which was defined in (4).)
We set

$$c_{d,R} \stackrel{\mathrm{def}}{=} |B_R(q)|. \tag{62}$$

(Recall that $B_R(q)$ was defined in (23) and $|\cdot|$ denotes the cardinality.)
Then

$$|\mathrm{Path}(q, n, R)| = c_{d,R}^n \tag{63}$$

because at each step in the path one can choose between $c_{d,R}$ lattice-points.
So we have

$$A_N \subset \bigcup_{n>N_0} \bigcup_{Q\in\mathrm{Path}(q,n,R)} \{\omega : Q \text{ is maximal causal wrt}(R, t, \omega)\} \tag{64}$$

and so

$$\begin{aligned} P(A_N) &\leq \sum_{n\geq N_0} c_{d,R}^n \, e^{-\lambda t} \, \frac{(\lambda t)^n}{n!} && (65)\\ &\leq e^{(c_{d,R}-1)\lambda t} \, (c_{d,R}\lambda t)^{N_0} \, \frac{1}{N_0!} && (66) \end{aligned}$$

which converges to 0 as $N_0 \to \infty$ which was to show. For the last inequality we have used the estimate for the Lagrange remainder in Taylor's formula.

□

Definition 2.2 Let a finite range interaction (i.e. a family of updatings) be given by $(S_q)_{q\in\mathbb{Z}^d}$. Fix $\omega \in \Omega \setminus (\Omega_R \cup \mathcal{N}_1)$, $\emptyset \neq \Lambda \subset \tilde{\Lambda} \in \mathcal{F}$, $\xi \in M$ and $t > 0$. Then ω has only finitely many jumps in $\tilde{\Lambda} \times (0, t)$, say at $(q_1, t_1), \ldots, (q_n, t_n)$ with $0 < t_1 < \ldots < t_n < t$.
We denote by $\mathbf{x}_{\tilde{\Lambda}} \vee \xi_{\tilde{\Lambda}^C}$ the point in M that has the same $\tilde{\Lambda}$-coordinates as $\mathbf{x}$ and the same $\tilde{\Lambda}^C$-coordinates as ξ.

We define

$$\begin{aligned} S_{q,\bar\Lambda,\xi} : (S^1)^{\bar\Lambda} &\to (S^1)^{\bar\Lambda} \\ S_{q,\bar\Lambda,\xi}(\mathbf{x}_{\bar\Lambda}) &\stackrel{\text{def}}{=} \pi_{\bar\Lambda} \circ S_q(\mathbf{x}_{\bar\Lambda} \vee \xi_{\bar\Lambda^C}) \end{aligned} \tag{67}$$

and

$$\begin{aligned} \Omega \setminus (\mathcal{N}_1 \cup \Omega_R) \ni \omega &\mapsto S^t_{\bar\Lambda,\xi,\omega} \in \mathcal{C}^0(M,M) \\ S^t_{\bar\Lambda,\xi,\omega}(\mathbf{x}) &\stackrel{\text{def}}{=} S_{q_n,\bar\Lambda,\xi} \circ \ldots \circ S_{q_1,\bar\Lambda,\xi}(\mathbf{x}_{\bar\Lambda}) \vee \xi_{\bar\Lambda^C}. \end{aligned} \tag{68}$$

The maps $S^t_{\bar\Lambda,\xi,\omega}$ are continuous as composites of continuous maps. Furthermore $S^t_{\bar\Lambda,\xi,\omega}$ depends only on $\omega_{\bar\Lambda}$ (i.e. on $\pi_{\bar\Lambda} \circ X_\bullet(\omega)$) and (68) gives rise to a countable, measurable partition of $\Omega \setminus (\mathcal{N}_1 \cup \Omega_R)$: ω and $\tilde\omega$ belong to the same set of this partition if they have the same list of jump sites $(q_1, \ldots, q_n)$ (ordered wrt the jump times).
Now let $\tilde\Lambda \supset \mathrm{Aff}_{(R,t,\omega)}(\Lambda)$ and $\xi \in M$ and define

$$\pi_\Lambda \circ S^t : \Omega \setminus (\mathcal{N}_1 \cup \Omega_R) \to \mathcal{C}^0(M, (S^1)^\Lambda), \tag{69}$$

$$\pi_\Lambda \circ S^t_\omega(\mathbf{x}) \stackrel{\text{def}}{=} \pi_\Lambda \circ S^t_{\bar\Lambda,\xi,\omega}(\mathbf{x}_{\bar\Lambda}). \tag{70}$$

The definition does not depend on the choice of $\tilde\Lambda$ or ξ because the right-hand side (rhs) of (70) depends, by definition, on the $\mathrm{Aff}_{(R,t,\omega)}(\Lambda)$-coordinates of $\mathbf{x}$ only.
Further the family $(\pi_\Lambda \circ S^t_\omega(\mathbf{x}))_{\Lambda \in \mathcal{F}\setminus\{\emptyset\}}$ is consistent in the sense that for any $\emptyset \neq \Lambda_1 \subset \Lambda_2 \in \mathcal{F}$:

$$\pi_{\Lambda_1}\left(\pi_{\Lambda_2} \circ S^t_\omega(x)\right) = \pi_{\Lambda_1} \circ S^t_\omega(x), \tag{71}$$

and so defines a map

$$\begin{aligned} S^t_\omega : M &\to M \\ (S^t_\omega(x))_q &\stackrel{\text{def}}{=} \pi_q \circ S^t_\omega(x) \end{aligned} \tag{72}$$

Finally we set $S^t_\omega = \mathrm{id}_M$ for $\omega \in \Omega_R \cup \mathcal{N}$.

Proposition 2.2 *The map S^t_ω, defined in (70) and (72) is continuous and depends measurably on ω.*

Proof The net $(\tilde S^t_{\bar\Lambda,\xi,\omega})_{\bar\Lambda \in \mathcal{F}\setminus\{\emptyset\}}$ satisfies the assumptions in Lemma 1.2 and so all statements of Proposition 2.2 follow.

□

2.2 Infinite Range Interaction

We extend our notion of 'S^t_ω' to interactions that are not necessarily of finite range.
Consider a family $(S_q)_{q\in\mathbb{Z}^d}$ of maps $S_q : M \to M$ such that S_q does not change the $\mathbb{Z}^d \setminus \{q\}$-coordinates and $\pi_q \circ S_q : M \to S^1$ is *Lipschitz-continuous wrt all coordinates* and the Lipschitz constants depend only on the relative positions of the sites, i.e. there are constants $w(r)$ for all $r \in \mathbb{Z}^d$ such that for all $q, \tilde q \in \mathbb{Z}^d$ and $\mathbf{x}, \mathbf{y} \in M$ with $\mathbf{x}_{\mathbb{Z}^d\setminus\{\tilde q\}} = \mathbf{y}_{\mathbb{Z}^d\setminus\{\tilde q\}}$ (i.e. $\mathbf{x}$ and $\mathbf{y}$ differ at most in their $\tilde q$-coordinates.)

$$\varrho_{S^1}(\pi_q \circ S_q(\mathbf{x}), \pi_q \circ S_q(\mathbf{y})) \le w(\tilde q - q)\, \varrho_{S^1}(y_{\tilde q}, z_{\tilde q}). \tag{73}$$

We further assume summability of the Lipschitz-constants, i.e.

$$\sum_{q\in\mathbb{Z}^d} w(q) = c_1 \tag{74}$$

with a positive constant c_1.
We need the following technical lemma.

Lemma 2.1 *If $(w(q))_{q\in\mathbb{Z}^d}$ is a family of non-negative real numbers satisfying (74) then there are families $(w_1(q))_{q\in\mathbb{Z}^d}$ and $(w_2(q))_{q\in\mathbb{Z}^d}$ of non-negative and positive numbers, respectively, such that*

$$\begin{aligned} w(q) &= w_1(q)\, w_2(q) \quad \text{for all } q\in\mathbb{Z}^d, && (75)\\ \sum_{q\in\mathbb{Z}^d} w_1(q) &\le 2c_1+1 && (76)\\ \text{and} \quad \lim_{R\to\infty} a(R) &= 0, && (77) \end{aligned}$$

where the positive function $a(\cdot)$ is defined by

$$a(R) \stackrel{\text{def}}{=} \sup_{\|r_1\|+\ldots+\|r_n\|=R} w_2(r_1)\cdot\ldots\cdot w_2(r_n) \tag{78}$$

(The empty product is defined to be equal to 1.)

Proof We can choose $r_0 = 0 < r_1 < \ldots \in \mathbb{N}$ such that

$$\sum_{\|q\|<r_i} w(q) \ge c_1 - 4^{-(i+1)} \quad \text{for } i\ge 1. \tag{79}$$

Then we have

$$\begin{aligned} \sum_{\|q\|<r_1} w(q) &\le c_1 && (80)\\ \text{and} \quad \sum_{r_i\le\|q\|<r_{i+1}} w(q) &\le 4^{-(i+1)} \quad \text{for } i\ge 1. \end{aligned}$$

We set for $i\ge 1$ and $r_{i-1}\le\|q\|<r_i$:

$$\begin{aligned} w_2(q) &\stackrel{\text{def}}{=} 2^{-i} && (81)\\ w_1(q) &\stackrel{\text{def}}{=} 2^i\, w(q). && (82) \end{aligned}$$

Then (75) is obviously satisfied. To prove (76) we use (80) and (82):

$$\begin{aligned} \sum_{q\in\mathbb{Z}^d} w_1(q) &= \sum_{i=0}^{\infty} \sum_{r_i\le\|q\|<r_{i+1}} w_1(q) && (83)\\ &\le 2c_1 + \sum_{i=1}^{\infty} 2^{-i}\\ &= 2c_1+1. \end{aligned}$$

Now we prove (77). We show by induction (wrt i) that for every $i\ge 1$ there is an n_i such that

$$a(R) < 2^{-i} \quad \text{for all } R \geq n_i. \tag{84}$$

For $i = 1$ the statement is true with $n_1 = 1$ because $a(R) \leq \frac{1}{2}$ for every $R \geq 1$ as there is at least one factor on the right-hand-side in (78) and each such factor is at most $\frac{1}{2}$.
Now we assume now that that the statement holds for i and n_i. We set

$$n_{i+1} \stackrel{\text{def}}{=} r_i + 2n_i. \tag{85}$$

Then every path $(q_0, \ldots, q_n)$ of length $R \geq n_{i+1}$ has at least one step of size at least r_i (i.e. there is an $1 \leq l \leq n$ such that $\|q_l - q_{l-1}\| \geq r_i$) or it can be divided into two paths both of length at least n_i (i.e. there is an $1 \leq l \leq m-1$ such that $\|q_0 - q_1\| + \ldots \|q_{l-1} + q_l\| \geq n_i$ and $\|q_{l+1} - q_l\| + \ldots \|q_n + q_{n-1}\| \geq n_i$). So each product on the right-hand side of (78) has at least one factor less than or equal to $2^{-(i+1)}$ or two factors less than or equal to 2^{-i}. As the other factors are smaller than 1 the product is bounded by $2^{-(i+1)}$ as was to be shown.

□

Now we fix (like in Lemma 2.1) a choice of $(w_1(q))_{q\in\mathbb{Z}^d}$ and $(w_2(q))_{q\in\mathbb{Z}^d}$ and so the function a.

Definition 2.3 We fix the metric ϱ_M on M by

$$\varrho_M(\mathbf{x}, \mathbf{y}) \stackrel{\text{def}}{=} \sup_{r\in\mathbb{Z}^d} a(\|r\|)\, \varrho_{S^1}(x_r, y_r). \tag{86}$$

Remark 2.1 It follows from Remark 1.5.1 and (77) that ϱ_M is a metric and compatible with the product topology.

Lemma 2.2 *The maps $S_q : M \to M$ are continuous (wrt the product topology on M).*

Proof According to Remark 1.5.2 and the uniform choice of the Lipschitz-constants (cf. (73)) we only have to show that the maps $\pi_q \circ S_0 : M \to S^1$ are continuous.
If $q \neq 0$ then the qth coordinate is not changed by S_0 and

$$\begin{aligned} a(\|q\|)\, \varrho_{S^1}(\pi_q \circ S_0(\mathbf{x}), \pi_q \circ S_0(\mathbf{y})) &= a(\|q\|)\, \varrho_{S^1}(x_q, y_q) \\ &\leq a(\|q\|)\, \frac{1}{a(\|q\|)}\, \varrho_M(\mathbf{x}, \mathbf{y}) \\ &\leq \varrho_M(\mathbf{x}, \mathbf{y}). \end{aligned} \tag{87}$$

If $q = 0$ we estimate

$$\begin{aligned} & a(0)\, \varrho_{S^1}(\pi_0 \circ S_0(\mathbf{x}), \pi_0 \circ S_0(\mathbf{y})) \\ &\leq a(0) \sum_{r\in\mathbb{Z}^d} w(r)\, \varrho_{S^1}(x_r, y_r) \\ &\leq a(0) \sum_{r\in\mathbb{Z}^d} w(r)\, \frac{1}{a(\|r\|)}\, \varrho_M(\mathbf{x}, \mathbf{y}) \\ &\leq a(0)\, (2c_1 + 1)\, \varrho_M(\mathbf{x}, \mathbf{y}) \end{aligned} \tag{88}$$

where we have used (73) for the first, the definition of ϱ_M for the second and (75) for the third inequality. So $\pi_q \circ S_0$ is continuous for all $q \in \mathbb{Z}^d$.

□

In the following we estimate the distance (wrt the uniform norm) between $\pi_0 \circ S^t_{\Lambda,\xi,\omega}$ and $\pi_0 \circ S^t_{\Lambda,\tilde{\xi},\omega}$ for different boundary conditions ξ_{Λ^C} and $\tilde{\xi}_{\Lambda^C}$ (that might even depend on the time) at the Λ^C-sites. Conditions (73) and (74) allow us to apply standard estimates for the 'error-growth' for composites of maps. Using the linear nature of the 'Lipschitz-condition' (73), we write the products of sums (over all coordinates, like in (73)) as sums (over paths) of products (corresponding to the particular paths).
We fix $t > 0$, $\Lambda \in \mathcal{F}$ and $\omega \in \Omega \setminus \mathcal{N}_1$. By definition of $\mathcal{N}_1$ (cf. (21)) ω has no jumps at 0, no simultaneous jumps and only finitely many jumps in $\Lambda \times (0,t)$, say at $(q_1,t_1),\ldots,(q_N,t_N)$ with $0 < t_1 < \ldots < t_N < t$. We set $t_0 \stackrel{\text{def}}{=} 0$ and fix arbitrary $\xi = (\xi(t_0),\ldots,\xi(t_N))$, $\tilde{\xi} = (\tilde{\xi}(t_0),\ldots,\tilde{\xi}(t_N)) \in M^{N+1}$ and $\mathbf{x},\mathbf{y} \in M$. We set $\mathbf{x}(0) \stackrel{\text{def}}{=} \mathbf{x}_\Lambda \vee \xi_{\Lambda^C}(0)$, $\mathbf{y}(0) \stackrel{\text{def}}{=} \mathbf{y}_\Lambda \vee \tilde{\xi}_{\Lambda^C}(0)$ and define for $1 \le i \le N$ recursively:

$$x_q(t_i) \stackrel{\text{def}}{=} \begin{cases} \pi_q \circ S_q(\mathbf{x}(t_{i-1})) & \text{for} \quad q = q_i \\ x_q(t_{i-1}) & \text{for} \quad q \in \Lambda \setminus \{q_i\} \\ \xi_q(t_i) & \text{for} \quad q \in \Lambda^C \end{cases} . \tag{89}$$

We define $\mathbf{y}(t_i)$ analogously, using $\mathbf{y}$ and $\tilde{\xi}$ instead of $\mathbf{x}$ and ξ, respectively.
Two points in S^1 can have distance at most $c_s = \mathrm{diam}_{\varrho_{S^1}}(S^1)$. For estimating the distance between $x_q(t_i)$ and $y_q(t_i)$ we define

$$\triangle_q(0) \stackrel{\text{def}}{=} \tilde{\triangle}_q(0) \stackrel{\text{def}}{=} \begin{cases} \varrho_{S^1}(x_q(0),y_q(0)) & \text{for} \quad q \in \Lambda \\ c_s & \text{for} \quad q \in \Lambda^C \end{cases} , \tag{90}$$

and for $1 \le i \le N$

$$\triangle_q(i) \stackrel{\text{def}}{=} \begin{cases} \sum_{r\in\mathbb{Z}^d} w(r-q)\,\triangle_r(i-1) & \text{for} \quad q = q_i \\ \triangle_q(i-1) & \text{for} \quad q \in \Lambda \setminus \{q_i\} \\ c_s & \text{for} \quad q \in \Lambda^C \end{cases} \tag{91}$$

$$\tilde{\triangle}_q(i) \stackrel{\text{def}}{=} \begin{cases} \max\{c_s, \sum_{r\in\mathbb{Z}^d} w(r-q)\,\tilde{\triangle}_r(i-1)\} & \text{for} \quad q = q_i \\ \tilde{\triangle}_q(i-1) & \text{for} \quad q \in \Lambda \setminus \{q_i\} \\ c_s & \text{for} \quad q \in \Lambda^C \end{cases}$$

The functions $\triangle_q$ and $\tilde{\triangle}_q$ depend on $\mathbf{x}$, $\mathbf{y}$ and Λ but we do not refer to this in our notation. We have introduced them for estimating the difference between $x_q(t_i)$ and $y_q(t_i)$ (cf. (92)) and so the difference between $x_q(t)$ and $y_q(t)$. This difference depends also on ω and so do the corresponding estimates for $\triangle_q$ and $\tilde{\triangle}_q$. In Definition 2.4 we will relate $\triangle_q$ and $\tilde{\triangle}_q$ to families of random variables $(Y_\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ and $(\tilde{Y}_\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$, respectively. For $\triangle_q$ we find a particularly nice expansion (cf. (93)). From this follows the convergence of Y_Λ to zero in expectation (as $\Lambda \to \mathbb{Z}^d$). We will show that $\tilde{Y}_\Lambda$ is bounded by Y_Λ and decreasing and so converges P-almost surely to zero by the Monotone Convergence Theorem (cf. Theorem 2.1).

Proposition 2.3 *The following holds for $0 \le i \le N$:*

1.
$$\varrho_{S^1}(x_q(t_i),y_q(t_i)) \le \tilde{\triangle}_q(i) \le \triangle_q(i) \tag{92}$$

2.
$$\triangle_q(i) = \sum_{\substack{(r_0=q_i,r_1,\ldots,r_n)\\ \in \mathrm{Path}_{\mathbf{C}}(t_i,\omega,q,\Lambda)}} w(r_1-r_0)\cdot\ldots\cdot w(r_n-r_{n-1})\,\triangle_{r_n}(0) \tag{93}$$

3. If in particular $\mathbf{x}_\Lambda = \mathbf{y}_\Lambda$ *then*

$$\begin{aligned} &\triangle_0(N(\omega)) \\ &\leq \quad c_S\, a(\mathrm{dist}_{\mathbb{Z}^d}(q, \Lambda^C)) \\ &\qquad \sum_{\substack{(r_0=q, r_1, \ldots, r_n) \\ \in \mathrm{Path}_{\mathrm{C}}(t,\omega,q\to\Lambda^C)}} w_1(r_1 - r_0) \cdot \ldots \cdot w_1(r_n - r_{n-1}) \end{aligned} \tag{94}$$

where $N(\omega)$ *is the number of jumps of* ω *in* $\Lambda \times (0, t)$.

Proof We prove (92) and (93) by induction wrt i.
$\underline{i = 0}$: (92) holds by definition of $\triangle_q(0)$ and $\tilde{\triangle}_q(0)$ (cf. (90)). At time 0 no jump has happened and the only summand on the right-hand-side in (93) corresponds to the empty path at site q and so the equality in (93) holds.
$\underline{i - 1 \to i}$: (93) holds obviously for i and $q \neq q_i$ as there is no updating at site q and

$$\mathrm{Path}_{\mathrm{C}}(t_i, \omega, q, \Lambda) = \mathrm{Path}_{\mathrm{C}}(t_{i-1}, \omega, q, \Lambda) \tag{95}$$

At site q_i there is a jump at time t_i and so we have

$$\triangle_{q_i}(i) = \sum_{r \in \mathbb{Z}^d} w(r - q_i)\, \triangle_r(i - 1) \tag{96}$$

Using the representation (93) for $\triangle_{r-q}(i-1)$ and the fact that every $(q_i, r_1, \ldots, r_n) \in \mathrm{Path}_{\mathrm{C}}(t_i, \omega, q_i, \Lambda)$ can be (uniquely) split into (q_i, r_1) and $(r_1, \ldots, r_n) \in \mathrm{Path}_{\mathrm{C}}(t_{i-1}, \omega, r_1, \Lambda)$, we see that (93) holds for i.
Next we show the first inequality in (92) for i. For $q \in \Lambda^C$ the distances between $x_q(t_i) = \xi_q(t_i)$ and $y_q(t_i) = \tilde{\xi}_q(t_i)$ is bounded by c_s and for $q \in \Lambda \setminus \{q_i\}$ we have $x_q(t_i) = x_q(t_{i-1})$ and $y_q(t_i) = y_q(t_{i-1})$. So in both cases the first inequality in (92) holds.
Now we consider the site q_i where a jump happens at time t_i. Using (73), assumption (92), for $i-1$, and (96), we get

$$\begin{aligned} \varrho_{S^1}(x_q(t_i), y_q(t_i)) &\leq \sum_{r \in \mathbb{Z}^d} w(r - q_i)\, \varrho_{S^1}(x_q(t_{i-1}), y_q(t_{i-1})) \\ &\leq \sum_{r \in \mathbb{Z}^d} w(r - q_i)\, \triangle_r(i - 1) \\ &\leq \triangle_{q_i}(i) \end{aligned} \tag{97}$$

So the first inequality in (92) is proved for i. The second follows immediately from (91). So statements 1 and 2 are proved.
Finally (94) follows from (93)): $\triangle_q(0) = 0$ for $q \in \Lambda$. So we only have to sum over paths $(r_0 = 0, \ldots, r_n)$ that end in $r_n \in \Lambda^C$.
In particular, if we set $R \stackrel{\mathrm{def}}{=} \|r_n\|$, then

$$\tilde{\triangle}_{r_n}(0) = c_s, \tag{98}$$

$$\mathrm{dist}_{\mathbb{Z}^d}(q, \Lambda^C) \leq R, \tag{99}$$

$$R \leq \|r_n - r_{n-1}\| + \ldots + \|r_1 - r_0\|, \tag{100}$$

and so by the choice of w_1, w_2 and a, made before Definition 2.3, we get

$$
\begin{aligned}
& w(r_1 - r_0) \cdot \ldots \cdot w(r_n - r_{n-1}) \\
\leq \quad & w_1(r_1 - r_0) \cdot \ldots \cdot w_1(r_n - r_{n-1})\, a(R) \\
\leq \quad & w_1(r_1 - r_0) \cdot \ldots \cdot w_1(r_n - r_{n-1})\, a(\mathrm{dist}_{\mathbb{Z}^d}(q, \Lambda^C)).
\end{aligned} \tag{101}
$$

Using (93), (98) and (101), we get (94) .

□

Remark 2.2 The summing over causal paths in Proposition 2.3 reflects that the result of an updating depends only on what has happened before.

Definition 2.4 We define two families $(Y_\Lambda)_{\Lambda \in \mathcal{F} \setminus \{\emptyset\}}$ and $(\tilde{Y}_\Lambda)_{\Lambda \in \mathcal{F} \setminus \{\emptyset\}}$ of random variables on $\Omega \setminus \mathcal{N}_1$. Let $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$ and $\omega \in \Omega \setminus \mathcal{N}_1$, say with exactly $N(\omega)$ jumps in $\Lambda \times [0,t]$. If we choose $\mathbf{x}, \mathbf{y} \in M$ with $\mathbf{x}_\Lambda = \mathbf{y}_\Lambda$ the value of $\triangle_0(N(\omega))$ (as defined by (90) and (91)) does not depend on $\mathbf{x}$ or $\mathbf{y}$. We define $Y_\Lambda(\omega)$ to be equal to this value:

$$
Y_\Lambda(\omega) \stackrel{\mathrm{def}}{=} \triangle_0(N(\omega)) \tag{102}
$$

$\tilde{Y}_\Lambda$ is defined analogously, using $\tilde{\triangle}_0(N(\omega))$ instead of $\triangle_0(N(\omega))$.

Remark 2.3

1. We remark that Y_Λ depends measurably on ω. In fact there is a countable, measurable partition of $\Omega \setminus \mathcal{N}_1$ such that ω and $\tilde{\omega}$ belong to the same set (of that partition) if the sums for $\triangle_0(t_{N(\omega)})$ and $\triangle_0(t_{N(\tilde{\omega})})$ (cf. (93)) are over the same paths.

2. From (92) we see that

$$
\tilde{Y}_\Lambda \leq Y_\Lambda. \tag{103}
$$

Now we fix $\xi, \mathbf{x} \in M$ and define the map $S^t_{\Lambda,\xi,\omega}$ like in (68).

Theorem 2.1 *1. There is a set $\mathcal{N}_2$ of P-measure zero such that*

$$
\lim_{\Lambda \to Z^d} \tilde{Y}_\Lambda = 0 \qquad \text{for } \omega \in \Omega \setminus (\mathcal{N}_1 \cup \mathcal{N}_2). \tag{104}
$$

2. The limit

$$
\pi_0 \circ S^t_\omega \stackrel{\mathrm{def}}{=} \lim_{\Lambda \to \mathbb{Z}^d} \pi_0 \circ S^t_{\Lambda,\xi,\omega} \tag{105}
$$

exists in $\mathcal{C}^0(M, S^1)$ for all $\omega \in \Omega \setminus (\mathcal{N}_1 \cup \mathcal{N}_2)$. It is measurable in ω and does not depend on ξ.

3. There is a set $\mathcal{N} \subset \Omega$ of P-measure zero such that we can define maps

$$
\pi_q \circ S^t_\omega \stackrel{\mathrm{def}}{=} \lim_{q \in \Lambda \to \mathbb{Z}^d} \pi_q \circ S^t_{\Lambda,\xi,\omega} \tag{106}
$$

for all $q \in \mathbb{Z}^d$ and $\omega \in \Omega \setminus \mathcal{N}$.

Further we can define a map $S^t_\omega \in \mathcal{C}^0(M, M)$ by

$$
\left(S^t_\omega(\mathbf{x})\right)_q \stackrel{\mathrm{def}}{=} \pi_q \circ S^t_\omega(\mathbf{x}). \tag{107}
$$

S^t_ω depends measurably on ω.

Proof First we show that

$$\lim_{\Lambda\to\mathbb{Z}^d} E(Y_\Lambda) = 0 \tag{108}$$

We set $R \stackrel{\text{def}}{=} \text{dist}_{\mathbb{Z}^d}(q, \Lambda^C)$. Using (94) we get

$$\begin{aligned} & E(Y_\Lambda) \qquad\qquad (109)\\ \leq\ & \int_\Omega dP(\omega)\, c_S\, a(R) \\ & \times \sum_{\substack{(r_0=0,r_1,\ldots,r_n)\\ \in \text{Path}_C(t,\omega,q\to\Lambda^C)}} w_1(r_1-r_0)\cdot\ldots\cdot w_1(r_n-r_{n-1}) \\ =\ & c_s\, a(R) \sum_{Q\in\text{Path}(0\to\Lambda^C)} w_1(r_1-r_0)\cdot\ldots\cdot w_1(r_n-r_{n-1}) \\ & \times P(\{\omega : Q\in \text{Path}_C(t,\omega,0\to\Lambda^C)\}) \end{aligned}$$

A path $Q = (q_0, q_1, \ldots, q_n)$ with $q_n \in \Lambda^C$ is causal wrt (t,ω) (i.e. $Q \in \text{Path}_C(t,\omega,0\to\Lambda^C)$) iff the Poisson process induced by Q has at least n jumps. So we can estimate the probability

$$\begin{aligned} P(\{\omega : Q\in \text{Path}_C(t,\omega,0\to\Lambda^C)\}) &= \sum_{m\geq n} e^{-\lambda t}\frac{(\lambda t)^m}{m!} && (110)\\ &\leq \frac{(\lambda t)^n}{n!} && (111) \end{aligned}$$

For the last line we have used Taylor's formula, as we did in (66). So we get, using (76),

$$\begin{aligned} E(Y_\Lambda) &\leq c_s\, a(R) \sum_{n=1}^{\infty} \frac{(\lambda t)^n}{n!} \left(\sum_{r\in\mathbb{Z}^d} w_1(r)\right)^n && (112)\\ &\leq c_2\, a(R) && (113) \end{aligned}$$

whith $c_2 = c_s\, e^{\lambda t(2c_1+1)}$. (Recall that we consider a fixed t at the moment, so c_2 is a constant.) By (77) we get

$$\lim_{\Lambda\to Z^d} E(Y_\Lambda) = 0 \tag{114}$$

and, using (103),

$$\lim_{\Lambda\to Z^d} E(\tilde{Y}_\Lambda) = 0. \tag{115}$$

$\tilde{Y}_\Lambda(\omega)$ is decreasing for all $\omega \in \Omega\setminus\mathcal{N}_1$: For a fixed ω and $0\in\Lambda_1\subset\Lambda_2\in\mathcal{F}$ we consider the (time-ordered) sequence of jumps $(q_1,t_1),\ldots,(q_n,t_n)$ of ω at sites $q_1,\ldots,q_2\in\Lambda_1$. It is a subsequence of the sequence of jumps $(\overline{q}_1,\overline{t}_1),\ldots,(\overline{q}_m,\overline{t}_m)$ of ω at sites $\overline{q}_1,\ldots,\overline{q}_m\in\Lambda_2$. The jumps (q_i,t_i) in the first sequence correspond to jumps $(\overline{q}_{j(i)},\overline{t}_{j(i)})$ in the second one. Then $q_i=\overline{q}_{j(i)}$ and $t_i=\overline{t}_{j(i)}$ but the indices i and $j(i)$ are not the same in general.
We define $\tilde{\triangle}^1_q(i)$ and $\tilde{\triangle}^2_q(j)$ as in (90) and (91) for the sets Λ_1 and Λ_2, respectively. We show that

$$\tilde{\triangle}^1_q(i) \geq \tilde{\triangle}^2_q(j(i)) \tag{116}$$

If $q \in {\Lambda_1}^C$ then (116) obviously holds because $\tilde{\triangle}^1_q(i) = c_S$ is an upper bound for $\tilde{\triangle}^2_q(j)$. For $q \in \Lambda_1$ we show (116) by induction wrt i.
If $i = 0$ then (116) is true by (90). Now assume that (116) holds for all q and a particular $i < n$. For $q \in \Lambda_1 \setminus \{q_{i+1}\}$ we have

$$\tilde{\triangle}^1_q(i+1) = \tilde{\triangle}^1_q(i) \geq \tilde{\triangle}^2_q(j(i)) = \tilde{\triangle}^2_q(j(i+1)) \tag{117}$$

where the inequality holds by assumption and the equalities by (91). For the site $q = q_{i+1}$ we have by (91)

$$\begin{aligned} \tilde{\triangle}^1_q(i+1) &= \max\{c_S, \sum_{r\in\mathbb{Z}^d} w(r-q)\tilde{\triangle}^1_r(i)\} \\ &\geq \max\{c_S, \sum_{r\in\mathbb{Z}^d} w(r-q)\tilde{\triangle}^2_r(j(i+1)-1)\} \\ &= \tilde{\triangle}^2_q(j(i+1)) \end{aligned} \tag{118}$$

which was to be shown. Here we have used that $\tilde{\triangle}^1_r(i) \geq \tilde{\triangle}^2_r(j(i+1)-1)$. This follows for $r \in \Lambda_1$ from the definition of $\tilde{\triangle}^1_r$ and $\tilde{\triangle}^2_r$ and for $r \in \Lambda_1$ from assumption (116) and the fact that $\tilde{\triangle}^2_r(j(i+1)-1) = \tilde{\triangle}^2_r(j(i))$. Using the definition of $\tilde{Y}_{\Lambda_1}(\omega)$ and $\tilde{Y}_{\Lambda_2}(\omega)$ (cf. Definition 2.4), we conclude

$$\tilde{Y}_{\Lambda_1}(\omega) \geq \tilde{Y}_{\Lambda_2}(\omega) \tag{119}$$

which was to be shown.
We have proved (115) and that $(\tilde{Y}_\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ is decreasing. So we conclude (104), by using the Monotone Convergence Theorem.
Now we prove the second statement in Theorem 2.1, using the first one. First we note that for $\omega \in \Omega \setminus (\mathcal{N}_1 \cup \mathcal{N}_2)$ the map $S^t_{\Lambda,\xi,\omega}$ is continuous since it is the composite of finitely many continuous (cf. Lemma 2.2) updating maps.
For $\Lambda \subset \tilde{\Lambda}$ we have

$$\varrho_{\mathcal{C}^0(M,S^1)}(\pi_0 \circ S^t_{\Lambda,\xi,\omega}, \pi_0 \circ S^t_{\tilde{\Lambda},\xi,\omega}) \leq \tilde{Y}_\Lambda(\omega). \tag{120}$$

So by (104) the net $(\pi_0 \circ S^t_{\Lambda,\xi,\omega})_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ is a Cauchy net with values in $\mathcal{C}^0(M,S^1)$ for $\omega \in \Omega\setminus(\mathcal{N}_1\cup\mathcal{N}_2)$ and so converges. Furthermore it is a pointwise limit, i.e. for each particular ω, and so $\pi_0 \circ S^t_\omega$ is measurable in ω. (The last conclusion uses the theorem that the pointwise limit of measurable functions with values in a metric space is measurable. (cf. for example [22], p. 117)).
As mentioned in Remark 1.5.3 there is no distinction of the point 0 by the product topology. So for all $q \in \mathbb{Z}^d$ we can define $\pi_q \circ S^t_\omega$ for all $\omega \in \Omega \setminus \mathcal{N}^q$ where $P(\mathcal{N}^q) = 0$. In the same way we can define for each $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$ and $\omega \in \Omega \setminus \mathcal{N}^\Lambda$ (with $P(\mathcal{N}^\Lambda) = 0$) maps $\pi_\Lambda \circ S^t_\omega \in \mathcal{C}^0(M, (S^1)^\Lambda)$ that depend measurably on ω, and such that $S^t_\omega(\mathbf{x})$ depends measurably on $(\omega, \mathbf{x})$.
The set

$$\mathcal{N} \stackrel{\text{def}}{=} \bigcup_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}} \mathcal{N}^\Lambda \tag{121}$$

has P-measure zero. So by Lemma 1.2 the map S^t_ω is well-defined for $\omega \in \Omega \setminus \mathcal{N}$ and the statements in 3. hold.

□

Remark 2.4 1. It follows from the proof of Theorem 2.1 that one can define a random dynamical system (cf. [1]), given by the map $R^{\geq} \times \Omega \times M \ni (t, \omega, x) \mapsto S^t_\omega(x)$ and the shifts $\theta(t)$ on Ω, defined

by $X^q_{t_2}(\theta(t_1)\omega) \stackrel{\rm def}{=} X^q_{t_1+t_2}(\omega) - X^q_{t_1}(\omega)$. One can further define for P-a.a. $\omega \in \Omega$ the linear operators '$\circ S^t_\omega$', acting on continuous functions and the corresponding transfer operators '$\mathcal{L}^t_\omega$'. So one has operator-valued random variables. However, in the following we consider only the averaged (wrt ω) operators.

2. We have defined maps $S^t_\omega \in \mathcal{C}^0(M,M)$ for finite range updating in (72) and a special class of infinite range updatings in (107), using (105). Note that the second class does not include the first. It might be interesting to find more general classes of (single site) updating functions for which the limit in (105) exists, or examples where it does not exist.

2.3 Markov Kernels

In Section 1 we defined the Poisson process $(\Omega, \mathcal{A}, P, (X_t)_{t\in[0,T]})$ with parameter λ and values in $\mathbb{N}^\Lambda$, the measure space $(M, \mathcal{B}_M, \mu)$ and the measurable space $(\mathcal{C}^0(M,M), \mathcal{B}_{\mathcal{C}^0(M,M)})$.
We have nets $(S^T_\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ of maps $S^T_\Lambda : \Omega\setminus\mathcal{N} \to \mathcal{C}^0(M,M)$ with limit $S^T \in \mathcal{C}^0(M,M)$, and the following statements hold:

1. S^T_Λ and S^T are $(\mathcal{A}, \mathcal{B}_{\mathcal{C}^0(M,M)})$-measurable.
2. S^T is the pointwise limit of the net $(S^T_\Lambda)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$.
3. For fixed $\mathbf{x} \in M$ the map $S^T_{\Lambda,\bullet}(\mathbf{x}) : \Omega \to M$ is $(\mathcal{A}, \mathcal{B}_M)$-measurable.

More precisely, for finite range interaction (cf. Section 2.1) S^T was defined in (72) and S^T_Λ in (68) (Now we drop the fixed boundary condition ξ and the '$\sim$' in the notation for convenience.) For infinite range interaction (cf. Section 2.2) we define S^T_Λ in the same way as for finite range interaction and the existence of the limit S^T is established in (107). Note that these maps are a priori not defined on a set of P-measure zero. For these exceptional $\omega \in \Omega$ we define S^T_ω and $S^T_{\Lambda,\omega}$ to be equal to the identity on M.
Statement 3. follows from measurability wrt $(\omega, \mathbf{x})$ of $S^T_{\Lambda,\omega}(\mathbf{x})$ (Proposition 2.2 and Remark 1.6.2 for finite range interaction and statement 3. of Theorem 2.1 and Remark 1.6.2 for infinite range interaction), the fact that one-point-sets in M are measurable, and Fubini's Theorem.
Like in Example 1.1 we set

$$\begin{aligned} K^T_S : M \times \mathcal{B}_M &\to [0,1] \\ K^T_S(\mathbf{x}, A) &\stackrel{\rm def}{=} P(\{\omega : S^T_\omega(\mathbf{x}) \in A\}). \end{aligned} \tag{122}$$

The corresponding operator, applied to an $f \in \mathcal{C}^0(M)$, is

$$(K^T_S f)(\mathbf{x}) = \int_M K^T_S(\mathbf{x}, d\mathbf{y})\, f(\mathbf{y}) \tag{123}$$

$$= \int_\Omega dP(\omega)\, f \circ S^T_\omega(\mathbf{x}). \tag{124}$$

(123) is the definition (cf. (32)), and (124) is a consequence of (122).
We define analogously the Markov kernels $K^T_{S,\Lambda}$ and corresponding operators for the Poisson process with values in $\mathbb{N}^\Lambda$.

Proposition 2.4 *K^T_S and $K^T_{S,\Lambda}$ are bounded linear operators on $\mathcal{C}^0(M)$.*

Proof We give the proof for K^T_S. The one for $K^T_{S,\Lambda}$ is analogous. Let $\omega \in \Omega$, $f \in \mathcal{C}^0(M)$ and $(\mathbf{x}^{(n)})_{n\in\mathbb{N}}$ a sequence in M with limit $\mathbf{x}$. Then

$$\lim_{n\to\infty} S_\omega^T(\mathbf{x}^{(n)}) = S_\omega^T(\mathbf{x}) \tag{125}$$

$$\text{and so} \quad \lim_{n\to\infty} f\circ S_\omega^T(\mathbf{x}^{(n)}) = f\circ S_\omega^T(\mathbf{x}). \tag{126}$$

Further

$$\|f\circ S_\omega^T\|_\infty \le \|f\|_\infty \tag{127}$$

Using the Dominated Convergence Theorem, we conclude

$$\begin{aligned}\lim_{n\to\infty}(K_S^T f)(\mathbf{x}^{(n)}) &= \lim_{n\to\infty}\int_\Omega dP(\omega)\, f\circ S_\omega^T(\mathbf{x}^{(n)}) \\ &= \int_\Omega dP(\omega)\, f\circ S_\omega^T(\mathbf{x}) \\ &= (K_S^T f)(\mathbf{x}).\end{aligned} \tag{128}$$

So $K_S^T f$ is continuous. Continuity of the operator follows from (124) and (127).

□

Proposition 2.5 *The net $(K_{S,\Lambda}^T)_{\Lambda\in\mathcal{F}\setminus\{\emptyset\}}$ converges weakly to K_S^T (as $\Lambda\to\mathbb{Z}^d$), i.e. for all $f\in\mathcal{C}^0(M)$:*

$$\lim_{\Lambda\to\mathbb{Z}^d} K_{S,\Lambda}^T f = K_S^T f. \tag{129}$$

Proof We have

$$\|K_S^T f - K_{S,\Lambda}^T f\|_\infty \le \int_\Omega dP(\omega)\, \|f\circ S_\omega^T - f\circ S_{\Lambda,\omega}^T\|_\infty. \tag{130}$$

Because of condition 2 on page 23 and (127) the rhs converges to 0 (as $\Lambda\to\mathbb{Z}^d$).

□

Remark 2.5 It follows from Remark 2.4.1 and the homogeneity of Poisson processes wrt time that $(K_S^t)_{t\ge 0}$ is a semigroup.

3 Transfer Operators

In this section we define transfer operators for the Markov kernels for a special class of updating functions that we have already studied in [13]. First we recall some definitions and notations from [13].
For $\delta>0$ we denote by A_δ the annulus

$$A_\delta \stackrel{\text{def}}{=} \{z\in\mathbb{C} \mid -\delta\le \ln|z|\le\delta\} \tag{131}$$

and by Γ its positively oriented boundary.
For $\emptyset\neq\Lambda\subset\mathbb{Z}^d$ the normalized Lebesgue-measure on $(S^1)^\Lambda$ is denoted by μ^Λ. For finite Λ it is given by

$$d\mu^\Lambda(\mathbf{z}) = \frac{d\mathbf{z}}{(2\pi i)^{|\Lambda|}}\frac{1}{\mathbf{z}} \stackrel{\text{def}}{=} \prod_{p\in\Lambda}\frac{dz_p}{2\pi i}\frac{1}{z_p}. \tag{132}$$

We also use $d\mu^\Lambda(\mathbf{z})$ as a shorthand notation for the right-hand side of (132) for $\mathbf{z}\in A_\delta^\Lambda$.

In **Assumption I** (see below) we will fix a $\delta > 0$. For $\Lambda \in \mathcal{F}$ we denote by $\mathcal{H}_\Lambda$ the space of continuous functions on the polyannulus A_δ^Λ that are holomorphic on its interior and write $\|\cdot\|_\Lambda$ for the uniform norm on $\mathcal{H}_\Lambda$. As a function on A_δ^Λ is also a function on $A_\delta^{\mathbb{Z}^d}$ we can drop the index Λ and mean the uniform norm on the infinite-dimensional polyannulus. $\mathcal{H}$ is the vectorspace of all consistent families $\phi = (\phi_\Lambda)_{\Lambda\in\mathcal{F}}$ of functions $\phi_\Lambda \in \mathcal{H}_\Lambda$. Consistency means

$$\begin{aligned}(\pi_{\Lambda_1}\phi_{\Lambda_2})(\mathbf{z}_{\Lambda_1}) &\stackrel{\text{def}}{=} \int_{(S^1)^{\Lambda_2\setminus\Lambda_1}} d\mu^{\Lambda_2\setminus\Lambda_1}(\mathbf{z}_{\Lambda_2\setminus\Lambda_1})\,\phi(\mathbf{z}_{\Lambda_1}\vee\mathbf{z}_{\Lambda_2\setminus\Lambda_1}) \\ &= \phi_{\Lambda_1}\end{aligned} \tag{133}$$

for all $\Lambda_1 \subseteq \Lambda_2 \in \mathcal{F}$ and $\mathbf{z}_{\Lambda_1} \in A_\delta^{\Lambda_1}$. (Note that we use the same symbol 'π_Λ' for projections of functions and projections of coordinates, for example from M to $(S^1)^\Lambda$.)
For $0 < \vartheta < 1$ and $\phi \in \mathcal{H}$ we define

$$\|\phi\|_\vartheta = \sup_{\Lambda\in\mathcal{F}} \vartheta^{|\Lambda|}\,\|\phi_\Lambda\|_\Lambda \tag{134}$$

$$\|\phi\|_{var} \stackrel{\text{def}}{=} \lim_{\Lambda\to\mathbb{Z}^d} \int_{(S^1)^\Lambda} d\mu^\Lambda\,|\phi_\Lambda|. \tag{135}$$

We set

$$\mathcal{H}_\vartheta \stackrel{\text{def}}{=} \{\phi\in\mathcal{H} : \|\phi\|_\vartheta < \infty\} \tag{136}$$

$$\mathcal{H}^{bv} \stackrel{\text{def}}{=} \{\phi\in\mathcal{H} : \|\phi\|_{var} < \infty\} \tag{137}$$

$$\mathcal{H}_\vartheta^{bv} \stackrel{\text{def}}{=} \mathcal{H}^{bv}\cap\mathcal{H}_\vartheta. \tag{138}$$

Then $(\mathcal{H}_\vartheta, \|\cdot\|_\vartheta)$ is a Banach space. For $\phi \in \mathcal{H}^{bv}$ and $\psi \in \mathcal{C}^0(M)$ we define

$$\psi_\Lambda(\mathbf{z}_\Lambda) \stackrel{\text{def}}{=} \int_{(S^1)^{\Lambda^C}} d\mu^{\Lambda^C}(\mathbf{z}_{\Lambda^C})\,\psi(\mathbf{z}_\Lambda\vee\mathbf{z}_{\Lambda^C}) \tag{139}$$

$$\int_M d\mu\,\psi\,\phi \stackrel{\text{def}}{=} \lim_{\Lambda\to\mathbb{Z}^d}\int_{(S^1)^\Lambda} d\mu_\Lambda\,\psi_\Lambda\,\phi_\Lambda. \tag{140}$$

Finally we recall the definition of a transfer operator: Let $\tilde\mu$ be a measure on the (completed) Borel σ-algebra of a metric space $\widetilde{M}$ and $\widetilde{S} : \widetilde{M} \to \widetilde{M}$ be a non-singular measurable map. The **Perron-Frobenius operator** (or **transfer operator**) $\mathcal{L}_{\widetilde{S}}$, acting on $L^1\left(\widetilde{M}\right)$, is defined via the equation

$$\int_{\widetilde{M}} d\tilde\mu\,\psi\circ\widetilde{S}\,\phi = \int_{\widetilde{M}} d\tilde\mu\,\psi\,\mathcal{L}_{\widetilde{S}}\phi \tag{141}$$

that must hold for all $\psi \in L^\infty\left(\widetilde{M}\right)$ and $\phi \in L^1\left(\widetilde{M}\right)$.

The Markov kernels for our stochastic systems are analogous to the composition operator '$\circ S$' (with deterministic S), acting on functions.

Definition 3.1 We define transfer operator for a Markov kernel K analogously to (cf. (229)) by the equation

$$\int_{\widetilde{M}} d\tilde\mu\,(K\psi)\,\phi = \int_{\widetilde{M}} d\tilde\mu\,\psi\,(\mathcal{L}_K\phi). \tag{142}$$

Remark 3.1 1. In the cases we consider, the Markov kernel K_S is given by

$$(K_S\psi)(x) = \int_\Omega dP(\omega)\, \psi \circ S_\omega(x), \tag{143}$$

where Ω is a probability space, S_ω depends measurably on ω and the map $\omega \mapsto \mathcal{L}_{S_\omega}$ is well-defined and integrable. Then

$$\begin{aligned} \int_{\widetilde{M}} d\tilde{\mu}\, (K\psi)\, \phi &= \int_{\widetilde{M}} d\tilde{\mu}(x) \int_\Omega dP(\omega)\, \psi \circ S_\omega(x)\, \phi(x) \\ &= \int_{\widetilde{M}} d\tilde{\mu}(x) \int_\Omega dP(\omega)\, \psi(x)\, (\mathcal{L}_{S_\omega}\phi)(x) \\ &= \int_{\widetilde{M}} d\tilde{\mu}(x)\, \psi(x) \int_\Omega dP(\omega)\, (\mathcal{L}_{S_\omega}\phi)(x). \end{aligned} \tag{144}$$

So $\mathcal{L}_{K_S}$ is given by

$$(\mathcal{L}_{K_S}\psi)(x) = \int_\Omega dP(\omega)\, (\mathcal{L}_{S_\omega}\phi)(x) \tag{145}$$

2. The operator for the infinite dimensional system that we are going to consider act on elements of $\mathcal{H}_\vartheta$ that do not in general correspond to elements of $L^1(M)$. Recall (see [13]) that $\mathcal{H}^{bv}_\vartheta$ can be identified with a subset of $rca(M)$ (or, in other words, a subset of the Borel measures). So for example in Theorem 3.1 we will show that the equation analogous to (142) holds for $\psi \in \mathcal{C}^0(M)$ (rather than $L^\infty(M)$) and $\phi \in \mathcal{H}^{bv}_\vartheta$.

Now we consider a special class of interactions (cf. [13]), namely a family $(S_\Lambda)_{\Lambda\in\mathcal{F}}$ of maps on M that can be written as

$$\begin{aligned} S_\Lambda : M &\to M \\ S_\Lambda(\mathbf{z}) &= F_\Lambda \circ T_\Lambda(\mathbf{z}) \vee \mathbf{z}_{\Lambda^c} \end{aligned} \tag{146}$$

where

$$\begin{aligned} F_\Lambda : (S^1)^\Lambda &\to (S^1)^\Lambda \\ \mathbf{z}_\Lambda = (z_q)_{q\in\Lambda} &\mapsto (f_q(z_q))_{q\in\Lambda} \end{aligned} \tag{147}$$

and

$$T_\Lambda : M \to (S^1)^\Lambda \tag{148}$$

$$(T_\Lambda(\mathbf{z}))_q \stackrel{\text{def}}{=} z_q \exp\left(2\pi i\epsilon \sum_{k=1}^{\infty} g_{q,k}(\mathbf{z})\right) \quad \text{for } q \in \Lambda \tag{149}$$

and f_q and $g_{q,k}$ satisfy the following assumptions:

Assumption I $F(\mathbf{z}) = (f_q(z_q))_{q\in\mathbb{Z}^d}$ where $f_q : S^1 \to S^1$ are real analytic and expanding (i.e. $f_q' \geq \lambda_0 > 1$) maps that extend for some δ_1 holomorphically to the interior of an annulus A_{δ_1}. In Proposition 3.1 and 3.2 of [13] we have shown that the holomorphic extension to a sufficiently thin annulus A_δ is

expanding in the sense that the preimage of A_δ wrt f_q lies in the interior of A_δ. We fix such a δ_1. Then for every $q \in \mathbb{Z}^d$ the Perron-Frobenius operator $\mathcal{L}_{f_q}$, acting on $\mathcal{H}_{\{q\}}$, has a simple largest eigenvalue 1 with eigenvector h_q, such that $\pi_\emptyset(h_q) = 1$ and the restriction of h_q to S^1 is positive and it splits into

$$\mathcal{L}_{f_q} = \mathcal{Q}_q + \mathcal{R}_q, \tag{150}$$

where $\mathcal{Q}_q$ is a projection onto span(h_q). We assume that there are positive constants c_h and c_r such that the following two estimates hold for all $q \in \mathbb{Z}^d$:

$$\|\mathcal{Q}_q\|_{\{q\}} \leq c_h \tag{151}$$

$$\|\mathcal{R}_q^n\|_{\{q\}} \leq c_r \eta^n \tag{152}$$

where $\|\cdot\|_{\{q\}}$ denotes both the uniform norm on $\mathcal{H}_{\{q\}}$ (for this we might have to take δ_1 even smaller) and the induced operator-norm. We note that this holds in particular if f_q does not depend on q.
We further have

$$\mathcal{Q}_q \mathcal{R}_q = \mathcal{R}_q \mathcal{Q}_q = 0 \tag{153}$$

Assumption II For all $q \in \mathbb{Z}^d$ and $k \geq 1$ each map $g_{q,k}$ extends to a holomorphic map $g_{q,k} : A_{\delta_1}^{B_k(q)} \to \mathbb{C}$ (recall definition (23) of $B_k(q)$) and its sup-norm (of modulus) is exponentially bounded by

$$\| g_{q,k} \|_{A_{\delta_1}^{B_k(q)}} \leq c_3 \exp\left(-c_g k^d\right) \tag{154}$$

with $c_3 > 0$ and 'large' $c_g > 0$. (In several statements in Section 3 and 4 a lower bound for c_g will come out of our computations. The idea is always that our estimates work, provided c_g is bigger than a certain constant.)

For $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$ we denote by h_Λ the function

$$h_\Lambda(\mathbf{z}_\Lambda) \stackrel{\text{def}}{=} \prod_{q \in \Lambda} h_q(z_q), \tag{155}$$

where h_q is as in Assumption I. We set $h_\emptyset = 1$ and

$$h_{\mathbb{Z}^d} \stackrel{\text{def}}{=} (h_\Lambda)_{\Lambda \in \mathcal{F}} \in \mathcal{H}. \tag{156}$$

We further define for a fixed $\xi \in M$ and $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$ and $\Lambda_1 \subseteq \Lambda$ the *updating at the Λ_1-sites with fixed boundary conditions ξ_{Λ^C} outside Λ* (or *cut-off* of S_{Λ_1}):

$$\begin{aligned} S_{\Lambda_1,\Lambda} : (S^1)^\Lambda &\to (S^1)^\Lambda \\ \mathbf{z}_\Lambda &\mapsto \pi_\Lambda \circ S_{\Lambda_1}(\mathbf{z}_\Lambda \vee \xi_{\Lambda^C}). \end{aligned} \tag{157}$$

And for $\mathbf{z}_{\Lambda \setminus \Lambda_1} \in (S^1)^{\Lambda \setminus \Lambda_1}$ we define

$$\begin{aligned} \pi_{\Lambda_1} \circ S_{\Lambda_1,\Lambda}(\cdot \vee \mathbf{z}_{\Lambda \setminus \Lambda_1}) : (S^1)^{\Lambda_1} &\to (S^1)^{\Lambda_1} \\ \mathbf{z}_{\Lambda_1} &\mapsto \pi_{\Lambda_1} \circ S_{\Lambda_1,\Lambda}(\mathbf{z}_{\Lambda_1} \vee \mathbf{z}_{\Lambda \setminus \Lambda_1}). \end{aligned} \tag{158}$$

Remark 3.2 1. The map defined in (158) is the cut-off of S wrt Λ_1 and boundary conditions $\mathbf{z}_{\Lambda \setminus \Lambda_1} \vee \xi_\Lambda$. So we can use the special representation in terms of integral kernels for its transfer operator, restricted to $\mathcal{H}_{\Lambda_1}$, for the proposition below.

2. The family $(S_q)_{q\in\mathbb{Z}^d}$, defined by (146), satisfies conditions (73) and (74) as one can see from [13]: The partial derivatives are estimated in the proof of Proposition 3.1 there.

Lemma 3.1 *Let $\Lambda \in \mathcal{F}\setminus\{\emptyset\}$ be the disjoint union of Λ_1 and Λ_2. The transfer operator, restricted to $\mathcal{H}_{\Lambda_1}$, of the map $S_{\Lambda_1,\Lambda} : (S^1)^\Lambda \to (S^1)^\Lambda$, defined in (157) has the following representation in terms of integral kernels:*

$$\begin{aligned}&\left(\mathcal{L}_{S_{\Lambda_1,\Lambda}}\phi\right)(\mathbf{w}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}) \\ &= \int_{\Gamma^{\Lambda_1}} d\mu^{\Lambda_1}(\mathbf{z}_{\Lambda_1})\,\phi(\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2})\prod_{q\in\Lambda_1}\frac{(S_{\Lambda_1,\Lambda}(\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}))_q}{(S_{\Lambda_1,\Lambda}(\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}))_q - w_q}\end{aligned} \tag{159}$$

for $\phi\in\mathcal{H}_\Lambda$.

Proof Let $\psi\in\mathcal{C}^0((S^1)^\Lambda)$. We use the notation $\phi_{\mathbf{w}_{\Lambda_2}}$ for the function $\mathbf{w}_{\Lambda_1}\mapsto\phi(\mathbf{w}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2})$.

$$\begin{aligned}&\int_{(S^1)^\Lambda} d\mu^\Lambda(\mathbf{w}_\Lambda)\,\psi\circ S_{\Lambda_1,\Lambda}(\mathbf{w}_\Lambda)\,\phi(\mathbf{w}_\Lambda) \\ &= \int_{(S^1)^{\Lambda_2}} d\mu^{\Lambda_2}(\mathbf{w}_{\Lambda_2})\int_{(S^1)^{\Lambda_1}} d\mu^{\Lambda_1}(\mathbf{w}_{\Lambda_1})\,\psi_{\mathbf{w}_{\Lambda_2}}\circ\pi_{\Lambda_1}\circ S_{\Lambda_1,\Lambda}(\mathbf{w}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}) \\ &\qquad\qquad \times\phi_{\mathbf{w}_{\Lambda_2}}(\mathbf{w}_{\Lambda_1}) \\ &= \int_{(S^1)^{\Lambda_2}} d\mu^{\Lambda_2}(\mathbf{w}_{\Lambda_2})\int_{(S^1)^{\Lambda_1}} d\mu^{\Lambda_1}(\mathbf{w}_{\Lambda_1})\,\psi_{\mathbf{w}_{\Lambda_2}}(\mathbf{w}_{\Lambda_1}) \\ &\qquad\qquad \times(\mathcal{L}_{\pi_{\Lambda_1}\circ S_{\Lambda_1,\Lambda}(\,\cdot\,\vee\mathbf{w}_{\Lambda_2})}\phi_{\mathbf{w}_{\Lambda_2}})(\mathbf{w}_{\Lambda_1}) \\ &= \int_{(S^1)^\Lambda} d\mu^\Lambda(\mathbf{w}_\Lambda)\,\psi(\mathbf{w}_\Lambda)\,(\mathcal{L}_{\pi_{\Lambda_1}\circ S_{\Lambda_1,\Lambda}(\,\cdot\,\vee\mathbf{w}_{\Lambda_2})}\phi_{\mathbf{w}_{\Lambda_2}})(\mathbf{w}_{\Lambda_1})\end{aligned} \tag{160}$$

Using the representation of the transfer operator for $\pi_{\Lambda_1}\circ S_{\Lambda_1,\Lambda}(\,\cdot\,\vee\mathbf{w}_{\Lambda_2})$ that we established in Proposition 3.3 of [13], we obtain the rhs of (159).

□

Remark 3.3 1. We see in particular that $\mathcal{L}_{S_{\Lambda_1,\Lambda}}$ 'acts on the Λ_1-coordinates' only. There is no integration wrt the Λ_2-coordinates.

For $q\in\Lambda_1$ we can split the factor

$$\frac{(S_{\Lambda_1,\Lambda}(\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}))_q}{(S_{\Lambda_1,\Lambda}(\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}))_q - w_q} = h_q(w_q,z_q) + r_q(w_q,z_q) + \sum_{k=1}^{\infty}\beta_{q,k}(w_q,\mathbf{z}_{\Lambda_1}\vee\mathbf{w}_{\Lambda_2}\vee\xi_{\Lambda^C}) \tag{161}$$

as in [13]. The integral kernels h_q and r_q correspond to the operators $\mathcal{Q}_q$ and $\mathcal{R}_q$, introduced in (150) and $\beta_{q,k}$ to $\mathcal{B}_{q,k}$, say. In addition to (153) we have

$$\mathcal{Q}_q\circ\mathcal{B}_{q,k} = 0 \tag{162}$$

for all k. (Cf. [13] for details.)

2. We have established a representation of the transfer operator also for updatings at more than one point at one time. Such simultaneous updatings happen, for example, in certain discrete time processes with positive probability (cf. [12]).

   As in the systems considered here simultaneous updatings P-almost never happen, we can restrict ourself in the following to the case of updatings at single sites, i.e. $\Lambda_1 = \{q\}$.

Definition 3.2 We define for fixed $\Lambda \in \mathcal{F} \setminus \{\emptyset\}$, $\xi \in M$ and a finite sequence $j = (q_1, \ldots, q_n) \in J$ of points in Λ the map

$$\begin{aligned} S_{j,\Lambda} : (S^1)^\Lambda &\to (S^1)^\Lambda \\ S_{j,\Lambda} &\stackrel{\text{def}}{=} S_{q_n,\Lambda} \circ \ldots \circ S_{q_1,\Lambda}. \end{aligned} \tag{163}$$

Here $S_{q,\Lambda}$ is the map for the updating at site q. Recall that in Definition 1.13 we defined the maps $\mathbf{j}$. For all $\omega \in \Omega$ there is a finite sequence $\mathbf{j}(\omega) = (q_1, \ldots, q_n)$ and so

$$\mathcal{L}_{S_{\mathbf{j}(\omega),\Lambda}} \stackrel{\text{def}}{=} \mathcal{L}_{S_{q_n,\Lambda}} \circ \ldots \circ \mathcal{L}_{S_{q_1,\Lambda}} \tag{164}$$

is well-defined.

Before establishing particular representations of the transfer operators $\mathcal{L}_\Lambda^T$ and $\pi_\Lambda \circ \mathcal{L}^T$ we consider some special examples. For these we need the following definition.

Definition 3.3 We define:

$$\mathcal{R}(t) = e^{-\lambda t} \sum_{k=0}^{\infty} \frac{(\lambda t)^k}{k!} \mathcal{R}^k \tag{165}$$

$$= \exp(\lambda t\,(\mathrm{id} - \mathcal{R})). \tag{166}$$

Then we have, using (152),

$$\mathcal{R}(t_1)\,\mathcal{R}(t_2) = \mathcal{R}(t_1 + t_2) \tag{167}$$

$$\|\mathcal{R}(t)\| \leq c_r e^{-(1-\eta)\lambda t} \tag{168}$$

Example 3.1 Consider a single site system with an updating map $f : S^1 \mapsto S^1$ that satisfies **Assumption I** . We have for fixed time $T > 0$ and jump rate $\lambda > 0$ a Markov kernel K_f^T, acting on functions $\psi \in \mathcal{C}^0(S^1)$ as described in Example 1.3. According to (164) and Example 1.3 we get a transfer operator $\mathcal{L}_f^T$, acting on $\mathcal{H}_p$:

$$\begin{aligned} \mathcal{L}_f^T &= \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \mathcal{L}_f^n \\ &= \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} (\mathcal{Q} + \mathcal{R})^n \qquad (169) \\ &= (1 - e^{-\lambda t})\,\mathcal{Q} + \mathcal{R}(t) \qquad (170) \end{aligned}$$

with $\mathcal{R}(t)$ as in Definition 3.3. Note that we think of the summand $e^{-\lambda t}\mathrm{id}$, corresponding to $n = 0$, as $e^{-\lambda t} R^0$. By (153) we have for $n \geq 1$ that $(\mathcal{Q} + \mathcal{R})^n = \mathcal{Q} + \mathcal{R}^n$, and so we get (170).



Figure 2: single site, unperturbed case: There are only two gum configurations



Figure 3: single site, perturbed case: example for a gum configuration

Example 3.2 Now we consider a small perturbation f_ϵ of the single site system $f_0 = f$ of Example 3.1, that depends on fixed boundary conditions. For simplicity we split the transfer operator for the single updating into $\mathcal{L}_{f_\epsilon} = \mathcal{Q} + \mathcal{R} + \mathcal{B}$ where $\mathcal{B}$ is the difference between the operators for the perturbed and the unperturbed system.
For a given number N of updatings we distribute

$$\mathcal{L}_{f_\epsilon}^N = (\mathcal{Q} + \mathcal{R} + \mathcal{B})^N, \tag{171}$$

using (153) and (162). Let K denote the number of factors $\mathcal{B}$, n_0 the number of factors, either all $\mathcal{Q}$ or all $\mathcal{R}$, before the first factor $\mathcal{B}$ and n_i the number of factors R after the ith $\mathcal{B}$. So the total number of factors is $N = n_0 + \ldots + n_K + K$.
We get

$$\begin{aligned}
\mathcal{L}_{f_\epsilon}^T &= \sum_{N=0}^{\infty} e^{-\lambda T} \frac{(\lambda T)^N}{N!} \mathcal{L}_{f_\epsilon}^N && (172)\\
&= \sum_{N=0}^{\infty} e^{-\lambda T} \sum_{\substack{n_0,\ldots,n_K \geq 0 \\ n_0+\ldots+n_K+K=N}} \frac{(\lambda T)^{N+K}}{(N+K)!} \mathcal{R}^{n_K} \circ \mathcal{B} \circ \ldots \circ \mathcal{B} \circ \mathcal{R}^{n_1} \circ \mathcal{B} \circ (\mathcal{Q}+\mathcal{R})^{n_0} && (173)\\
&= \sum_{K=0}^{\infty} \sum_{\substack{n_0 \geq 1 \\ n_1,\ldots,n_K \geq 0}} e^{-\lambda T} \frac{(\lambda T)^{n_0+\ldots n_K+K}}{(n_0+\ldots n_K+K)!} \mathcal{R}^{n_K} \circ \mathcal{B} \circ \ldots \circ \mathcal{B} \circ \mathcal{R}^{n_1} \circ \mathcal{B} \circ \mathcal{Q} && (174)\\
&\quad + \sum_{K=0}^{\infty} \sum_{n_0,\ldots,n_K \geq 0} e^{-\lambda T} \frac{(\lambda T)^{n_0+\ldots n_K+K}}{(n_0+\ldots n_K+K)!} \mathcal{R}^{n_K} \mathcal{B} \circ \ldots \circ \mathcal{B} \circ \mathcal{R}^{n_1} \circ \mathcal{B} \circ \mathcal{R}^{n_0} \\
&= \sum_{K=0}^{\infty} \int_0^T \lambda dt_1 \int_{t_1}^T \lambda dt_2 \cdots \int_{t_K}^T \lambda dt_K \, (1 - e^{-\lambda t_1}) \, \mathcal{R}(t_n) \circ \mathcal{B} \circ \cdots \circ \mathcal{B} \circ \mathcal{R}(t_1) \circ \mathcal{B} \circ \mathcal{Q} && (175)\\
&\quad + \sum_{K=0}^{\infty} \int_0^T \lambda dt_1 \int_{t_1}^T \lambda dt_2 \ldots \int_{t_K}^T \lambda dt_K \, \mathcal{R}(t_n) \circ \mathcal{B} \circ \ldots \circ \mathcal{R}(t_0)
\end{aligned}$$

For the step from (174) to (175) we have used Lemma 3.2 (s. below). We interpret (175) in the following way. We write the operator $\mathcal{L}_{f_\epsilon}^T$ as a sum of operators $\mathcal{L}_{\mathcal{C}_g,T}$. For the time being we think of $\mathcal{C}_g$ as a diagram like, for example, in Figure 3. The vertical axis from top to bottom corresponds to the positively

oriented time line. Along this axis we draw K thick horizontal bars, denoted by B, that correspond to the operators $\mathcal{B}$. Between consecutive B's or between the first B and the top or the last B and the bottom or, in the case $K = 0$, between the top and the bottom we draw either a thick (r-strip) or a thin (h-strip) line, representing the choice of factors $\mathcal{R}$ or $\mathcal{Q}$ in the product (171), respectively. Note that below a B there must be a thick vertical line. For a fixed T and a K-tuple

$$\mathbf{t} = (t_1, \ldots, t_K) \in \{\mathbf{t} : -T < t_1 < \ldots < t_K < 0\} \tag{176}$$

we think of the top as being fixed at time $-T$, the bottom at 0 and the ith symbol B at t_i. That also fixes the lengths of the particular h- and r-strips. We assign to the triple $(\mathcal{C}_g, T, t)$ an operator $\mathcal{L}_{\mathcal{C}_g,T,\mathbf{t}}$. For example with $\mathcal{C}_g$, T and $\mathbf{t}$ as in Figure 3, we get

$$\mathcal{L}_{\mathcal{C}_g,T,\mathbf{t}} = (1 - e^{-\lambda(t_1+T)})\, \mathcal{R}(0 - t_K) \circ \mathcal{B} \ldots \circ \mathcal{B} \circ \mathcal{R}(t_2 - t_1) \circ \mathcal{B} \circ \mathcal{Q}. \tag{177}$$

To get $\mathcal{L}^T_{\mathcal{C}_g}$ we integrate (177) over the simplex given by (176) wrt the scaled Lebesgue measure $\lambda^K dt_1 \ldots dt_K$. The simplex has dimension K, so for each B in $\mathcal{C}_g$ we get one integration. Heuristically, the measure 'λdt_1' corresponds to the probability that a Poisson process with rate λ jumps in a small time interval. The approach of approximating the continuous time system by discrete time systems is made precise in [12]. We also note the special case $K = 0$ where the simplex degenerates to a single point of measure 1.

Above we have used the following lemma.

Lemma 3.2 Let $(n_{i,j})_{\substack{0\le i\le K\\ 1\le j\le N}}$ be a family of non-negative integers and $\lambda, T > 0$. Then, with the notation $t_0 \stackrel{\text{def}}{=} -T$, $t_{K+1} \stackrel{\text{def}}{=} 0$:

$$\begin{aligned} & e^{-N\lambda T} \frac{(\lambda T)^{n_{0,1}+\ldots+n_{K,N}+K}}{(n_{0,1}+\ldots+n_{K,N}+K)!} \binom{n_{0,1}+\ldots+n_{0,N}}{n_{0,1},\ldots,n_{0,N}} \cdot \ldots \cdot \binom{n_{K,1}+\ldots+n_{K,N}}{n_{K,1},\ldots,n_{K,N}} \\ = \quad & \int_{-T}^{0} \lambda dt_1 \int_{-t_1}^{0} \lambda dt_2 \cdots \int_{-t_{K-1}}^{0} \lambda dt_K \prod_{\substack{0\le i\le K\\ 1\le j\le N}} \left(e^{-\lambda(t_{i+1}-t_i)} \, \frac{(\lambda(t_{i+1}-t_i))^{n_{i,j}}}{n_{i,j}!} \right). \end{aligned} \tag{178}$$

Proof We see that the rhs of (178) is equal to

$$e^{-N\lambda T} \lambda^{n_{0,1}+\ldots+n_{K,N}+K} \prod_{\substack{0\le i\le K\\ 1\le j\le N}} \frac{1}{n_{i,j}!} \int_{-T}^{0} dt_1 \cdots \int_{-t_{K-1}}^{0} dt_K \prod_{0\le i\le K} (t_{i+1}-t_i)^{n_{i,1}+\ldots+n_{i,N}} \tag{179}$$

and so (178) follows by K times applying the identity

$$\int_{-\tau}^{0} dt\, (-t)^n (t+\tau)^m = \frac{n!\, m!}{(n+m)!}\, \tau^{n+m+1} \tag{180}$$

which can be easily shown by iterated integration by parts. □

Example 3.3 Now we consider a system with nearest neighbour coupling where only two adjacent sites (1 and 2) are updated and the states at the other sites are fixed. This time an expansion as in Example 3.2 gives rise to summing operators $\mathcal{L}_{\mathcal{C}_g,T}$ where the $\mathcal{C}_g$ correspond to diagrams like in Figure 4.
We have to consider all finite ordered sequences of symbols B_1 and B_2.
In the example of Figure 4 we have the sequence (B_2, B_1). For each B_i we draw a thick horizontal bar, centered in the column corresponding to the i-th site and connecting to all sites (columns) on which the



Figure 4: two sites, example for a gum configurations

operator $\mathcal{B}_i$ depends. We draw the sequence of B_i's 'downwards'. As we consider only nearest neighbour updatings here, the B_i have width 2.
Then we can choose at sites 1 and 2 between h- and r-strips. Note that in Figure 4 at site 2 an h-strip follows an r-strip. This is possible because they are 'separated' by B_1'. Now we consider the particular $\mathcal{C}_g$ and T shown in Figure 4. Let $n_{0,1}$ denote the number of jumps at site 1 before the jumps that corresponds to $\mathrm{id}_1 \otimes \mathcal{B}_2$ etc. The definition of $n_{i,j}$ in general is analogous. So the total number of jumps is $N = n_{0,1} + \ldots + n_{2,2} + 2$. The first two factors in (182) give the probability that a particular sequence of exactly N jumps occurs. The first binomial coefficient counts the combinations (different sequences) of $n_{0,2}$ jumps at site 1 and $n_{0,2}$ jumps at site 2, the others are explained analogously.
We get the operator:

$$\mathcal{L}_{\mathcal{C}_g,T} \tag{181}$$

$$= \sum_{\substack{n_{0,1},n_{1,1},n_{2,2}\geq 1\\ n_{0,2},n_{1,2},n_{0,2}\geq 0}} e^{-2\lambda T}\frac{(\lambda T)^{n_{0,1}+\ldots+n_{2,2}+2}}{(n_{0,1}+\ldots n_{2,2}+2)!}\binom{n_{0,1}+n_{0,2}}{n_{0,1},n_{0,2}}\binom{n_{1,1}+n_{1,2}}{n_{1,1},n_{1,2}}\binom{n_{2,1}+n_{2,2}}{n_{2,1},n_{2,2}} \tag{182}$$

$$\times\mathcal{R}_1^{n_{2,1}}\otimes\mathcal{Q}_2\circ\mathcal{B}_1\otimes\mathrm{id}_2\circ\mathcal{Q}_1\otimes\mathcal{R}_2^{n_{1,2}}\circ\mathrm{id}_1\otimes\mathcal{B}_2\circ\mathcal{Q}_1\otimes\mathcal{R}_2^{n_{0,2}}$$

$$= \int_{-T}^{0}\lambda dt_1\int_{t_1}^{0}\lambda dt_2\left(\sum_{n_{2,1}\geq 0}e^{-\lambda(|t_2|)}\frac{(\lambda|t_2|)^{n_{2,1}}}{n_{2,1}!}\mathcal{R}_1^{n_{2,1}}\right)\otimes\left(\sum_{n_{2,2}\geq 1}e^{-\lambda|t_2|}\frac{(\lambda|t_2|)^{n_{2,2}}}{n_{2,2}!}\mathcal{Q}_2\right) \tag{183}$$

$$\circ\mathcal{B}_1\otimes\mathrm{id}_2\circ\left(\sum_{n_{1,1}\geq 0}e^{-\lambda(t_2-t_1)}\frac{(\lambda(t_2-t_1))^{n_{1,1}}}{n_{1,1}!}\mathcal{Q}_1\right)\otimes\left(\sum_{n_{1,2}\geq 0}e^{-\lambda(t_2-t_1)}\frac{(\lambda(t_2-t_1)^{n_{1,2}}}{n_{1,2}!}\mathcal{R}_2^{n_{1,2}}\right)$$

$$\circ\mathrm{id}_1\otimes\mathcal{B}_2\circ\left(\sum_{n_{0,1}\geq 0}e^{-\lambda(t_1+T)}\frac{(\lambda(t_1+T))^{n_{0,1}}}{n_{0,1}!}\mathcal{Q}_1\right)\otimes\left(\sum_{n_{0,2}\geq 0}e^{-\lambda(t_1+T)}\frac{(\lambda(t_1+T)^{n_{0,2}}}{n_{0,2}!}\mathcal{R}_2^{n_{0,2}}\right)$$

$$= \int_{0}^{T}\lambda dt_1\int_{t_1}^{T}\lambda dt_2\left(1-e^{-\lambda|t_2|}\right)\left(1-e^{-\lambda(t_2-t_1)}\right)\left(1-e^{-\lambda(t_1-T)}\right)$$

$$\times\mathcal{R}_1(|t_2|)\otimes\mathcal{Q}_1\circ\mathcal{B}_1\otimes\mathrm{id}_2\circ\mathcal{Q}_1\otimes\mathcal{R}_2(t_2-t_1)\circ\mathrm{id}_1\otimes\mathcal{B}_2\circ\mathcal{Q}_1\otimes\mathcal{R}_2(t_1+T)$$

Note that the operator '$\otimes$' here has higher precedence than '$\circ$', so, for example, $A_1 \otimes A_2 \circ A_3 \otimes A_4$ is understood as $(A_1 \otimes A_2) \circ (A_3 \otimes A_4)$.

After these introductory examples we establish a diagrammatic representation of operators $\pi_{\Lambda_1} \circ \mathcal{L}^T_{S,\Lambda}$ and $\pi_{\Lambda_1} \circ \mathcal{L}^T_S$. We begin with some formal definitions for the diagrams.

Definition 3.4 In a labelled tree (as defined in [13]) we call the vertices that have no maximal label (i.e. they are not leaves of the tree and there is a vertex with a greater (wrt to the partial order) label) and that are different from the root vertex, **branchings**. (Each branching corresponds to a particular *star-graph* (like we used to define the term *labelled tree* in [13].) It is called k-**branching** if it corresponds to a star-graph with exactly $v(k)$ vertices (and hence to a k-triangle). The number k is called the **degree of the branching**.
Recall that in [13] we have introduced a linear order $\prec$ on the set of labels of vertices. In the following we will use the same symbol for linear orders on different sets as well. It should be clear from the context to which order we refer.
We say that a linear order $\prec$ on the set of branchings is **compatible with the labelling** if the following condition is satisfied: If $\mathbf{v_1}$ and $\mathbf{v_2}$ are branchings, labelled by label($\mathbf{v_1}$) and label($\mathbf{v_2}$), respectively, then

$$\text{label}(\mathbf{v_1}) \succ \text{label}(\mathbf{v_2}) \quad \Rightarrow \quad \mathbf{v_1} \prec \mathbf{v_2}. \tag{184}$$

We introduce a linear order (also denoted by $\prec$) on $\mathbb{Z}^d$:

$$(k_1, \ldots, k_d) \prec (\tilde{k}_1, \ldots, \tilde{k}_d) \quad \text{if } k_i < \tilde{k}_i \text{ for the lowest index } i \text{ such that } k_i \neq \tilde{k}_i \tag{185}$$

Definition 3.5 A **gum tree** τ_g with parameters $n_\beta = (n_{\beta,1}, n_{\beta,2}, \ldots)$ and $\Lambda_2 \in \mathcal{F} \setminus \{\emptyset\}$ is given by the following data:

1. A labelled tree τ with parameters n_β and $|\Lambda_2|$ (as defined before Lemma 8.2 of [13]).
2. A map **pin** from the set of vertices (except the root) of τ to $\mathbb{Z}^d$ that satisfies the following conditions:
   - The restriction of pin to the set of vertices, that are labelled by $(0,1), \ldots, (0, |\Lambda_2|)$ (We denote the restriction of pin to this set by pin_0.), is an order-preserving bijection onto Λ_2, i.e. for any two such vertices $\mathbf{v}$ and $\tilde{\mathbf{v}}$

     $$\text{label}(\mathbf{v}) \prec \text{label}(\tilde{\mathbf{v}}) \quad \Rightarrow \quad \text{pin}(\mathbf{v}) \prec \text{pin}(\tilde{\mathbf{v}}). \tag{186}$$

   - If $\mathbf{v}$ with $\text{label}(\mathbf{v}) = s = (s_1, \ldots, s_m)$ is a k-branching and $\text{pin}(\mathbf{v}) = q \in \mathbb{Z}^d$ then the restriction of pin to the set of vertices with labels $(s,1), \ldots, (s, v(k))$ (We denote the restriction of pin to this set by $\text{pin}_{\mathbf{v}}$.) is an order-preserving bijection onto $B_k(q) \subset \mathbb{Z}^d$.

A **gum configuration $\mathcal{C}_g$ on Λ ending in Λ_1** is given by the following data:

1. A gum tree τ_g with parameters n_β and Λ_2 such that $\Lambda_2 \subseteq \Lambda_1$. The corresponding tree has branchings $\mathbf{v_1} \prec \ldots \prec \mathbf{v_n}$, say, with branching-degrees $b_1, \ldots, b_n$, respectively.
2. For each $1 \leq i \leq n$ there are maps

$$u_i \quad : \quad B_{b_i}(\text{pin}(\mathbf{v_i})) \cap \Lambda \to \{0,1\} \tag{187}$$
$$d_i \quad : \quad B_{b_i}(\text{pin}(\mathbf{v_i})) \cap \Lambda \to \{0,1\}. \tag{188}$$

such that

(a) If $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i})) \cap \Lambda$ and j is the smallest number greater than i such that $q \in B_{b_j}(\mathrm{pin}(\mathbf{v_j}))$ (if such a j exists at all) then $d_i(q) = u_j(q)$.

(b) For every $1 \leq i \leq n$

$$d_i(\mathrm{pin}(\mathbf{v_i})) = 1. \tag{189}$$

(c) If $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i})) \cap (\Lambda \setminus \Lambda_1)$ and there is no $j > i$ such that $q \in B_{b_j}(q_j)$ then $d_i(q_i) = 0$.

(We will see later that the maps u_i define from a vertex upwards going *h-strips* (if $u_i = 0$) or *r-strips* (if $u_i = 1$). Similarly, the maps d_i determine downwards going strips. For a strip between two vertices it should be well-defined if it is an h-strip or an r-strip. Hence we impose condition (a). Condition (b) says that a strip that goes downwards from a branching must be an r-strip.)

3. A map **long** from $\Lambda \setminus \bigcup_{i=1}^{n} B_{b_i}(\mathrm{pin}(\mathbf{v_i}))$ to $\{0,1\}$ such that

$$\mathrm{long}(q) = 0 \quad \text{if } q \notin \Lambda_1 \tag{190}$$

We define in analogy to Definition 5.2 in [13]

$$\widetilde{\Lambda}(\mathcal{C}_g) \stackrel{\mathrm{def}}{=} \bigcup_{i=1}^{n} B_{b_i}(\mathrm{pin}(\mathbf{v_i})), \tag{191}$$

$$\Lambda_r(\mathcal{C}_g) \stackrel{\mathrm{def}}{=} \{q \in \Lambda \setminus \widetilde{\Lambda}(\mathcal{C}_g) : \mathrm{long}(q) = 1\}, \tag{192}$$

$$\Lambda(\mathcal{C}_g) \stackrel{\mathrm{def}}{=} \widetilde{\Lambda}(\mathcal{C}_g) \cup \Lambda_r(\mathcal{C}_g). \tag{193}$$

We introduce the following notation:

- In the situation of 2.a the point q is the image (wrt pin) of the vertices $\mathrm{pin}^{-1}_{\mathbf{v_i}}(q)$ and $\mathrm{pin}^{-1}_{\mathbf{v_j}}(q)$. We say that $\mathcal{C}_g$ has an h-**strip** (r-**strip**) from $\mathrm{pin}^{-1}_{\mathbf{v_i}}(q)$ to $\mathrm{pin}^{-1}_{\mathbf{v_j}}(q)$ if $d_i(q) = 0$ ($d_i(q) = 1$). (We note that we do not distinguish the order of the vertices in this notation: A strip from $\mathbf{v}$ to $\tilde{\mathbf{v}}$ is the same as a strip from $\tilde{\mathbf{v}}$ to $\mathbf{v}$.)

- If $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i})) \cap \Lambda$ and $\mathbf{v} = \mathrm{pin}^{-1}_{\mathbf{v_i}}(q)$ and there is no $j > i$ such that $q \in B_{b_j}(\mathrm{pin}(\mathbf{v_j}))$ and if $d_i(q) = 0$ ($d_i(q) = 1$) we say that $\mathcal{C}_g$ has an h-**strip (r-strip) from v to the bottom**.

- If $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i})) \cap \Lambda$ and $\mathbf{v} = \mathrm{pin}^{-1}_{\mathbf{v_i}}(q)$ and there is no number $j < i$ such that $q \in B_{b_j}(\mathrm{pin}(\mathbf{v_j}))$ and if $u_i(q) = 0$ ($u_i(q) = 1$) we say that $\mathcal{C}_g$ has an h-**strip (r-strip) from v to the top**.

- In the situation of 2.b we call the corresponding r-strip an **apex-r-strip**.

- If $q \in \Lambda \setminus \widetilde{\Lambda}(\mathcal{C}_g)$ and $\mathrm{long}(q) = 0$ ($\mathrm{long}(q) = 1$) then we say that $\mathcal{C}_g$ has a **long** h-**strip** (**long** r-**strip**) at q. So $\Lambda_r(\mathcal{C}_g) \subset \Lambda_1$ is the set of q where $\mathcal{C}_g$ has long r-strips.

- If $\mathcal{C}_g$ has an r-strip to the top or a long r-strip we say that $\mathcal{C}_g$ **reaches the top**.

We denote by $\mathbf{Conf_g}(\mathbf{\Lambda}, \mathbf{\Lambda_1})$ the set of all gum configurations over Λ ending in Λ_1.

Definition 3.6 Let $\mathcal{C}_g$ be a gum configuration over Λ ending in Λ_1 with branchings $\mathbf{v_1} \prec \ldots \prec \mathbf{v_n}$ of branching-orders $b_1, \ldots, b_n$, respectively, and let $T \in (0, \infty]$. Then we define

$$\mathrm{Simplex}(\mathcal{C}_g, T) \stackrel{\mathrm{def}}{=} \{(t_1, \ldots, t_n) : -T < t_1 < \ldots < t_n < 0\}. \tag{194}$$

$\mathrm{Simplex}(\mathcal{C}_g, T)$ is an open subset of $\mathbb{R}^n$ and so carries the induced Lebesgue-measure.
For the special case $n_\beta(\mathcal{C}_g) = 0$ we define $\mathrm{Simplex}(\mathcal{C}_g, T)$ to be a single point having measure 1.

Definition 3.7 For $\mathcal{C}_g \in \mathrm{Confg}(\Lambda, \Lambda_1)$, $T \in (0, \infty]$ and $\mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, T)$ we call the triple $(\mathcal{C}_g, T, \mathbf{t})$ a **specific gum configuration**.
Specific gum configurations can be viewed graphically: The vertices are placed in $\mathbb{Z}^d \times [0, T]$ and the strips are 'spanned' between vertices, the top ($t = -T$) and the bottom ($t = 0$):

- We assign to each vertex $\mathbf{v}$ in $\tau(\mathcal{C}_g)$ a point in $\mathbb{Z}^d \times [-T, 0]$ in the following way. If $\mathbf{v_i}$ is a branching of degree b_i, $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i}))$ and $\mathbf{v} = \mathrm{pin}_{\mathbf{v_i}}^{-1}(q)$ then $\mathbf{v}$ has time-coordinate t_i. In particular $\mathbf{v_i}$ has time-coordinate t_i. As further $\mathrm{pin}(\mathbf{v}) = q$ we assign $\mathbf{v}$ to (q, t_i).

  Let for the following two vertices $\mathbf{v}$ and $\tilde{\mathbf{v}}$ be assigned to (q, t) and $(q, \tilde{t})$, respectively.

- If $\mathcal{C}_g$ has an h-strip (r-strip) from $\mathbf{v}$ to $\tilde{\mathbf{v}}$ we say that $(\mathcal{C}_g, T, \mathbf{t})$ has a **maximal h-strip** (**maximal r-strip**) from (q, t) to $(q, \tilde{t})$. We define its **length** to be $|t - \tilde{t}|$.

- If $\mathcal{C}_g$ has an h-strip (r-strip) from $\mathbf{v}$ to the bottom (this has time-coordinate 0.) we say that $(\mathcal{C}_g, T, \mathbf{t})$ has a **maximal h-strip** (**maximal r-strip**) from (q, t) to $(q, 0)$. Its length is $|t|$.

- If $\mathcal{C}_g$ has an h-strip (r-strip) from $\mathbf{v}$ to the top (this has time-coordinate $-T$.) we say that $(\mathcal{C}_g, T, \mathbf{t})$ has a **maximal h-strip** (**maximal r-strip**) from (q, t) to $(q, -T)$. Its length is $T - |t|$. (Note that for $T = \infty$ this length is ∞.)

- If $\mathcal{C}_g$ has a long h-strip (long r-strip) at q we say that $(\mathcal{C}_g, T, \mathbf{t})$ has a **long h-strip (long r-strip)** at q. Its length is T. (Long h-strips (long r-strips) are also considered as maximal strips.)

If $(\mathcal{C}_g, T, \mathbf{t})$ has a maximal h-strip (r-strip) from $(q, \tilde{t}_1)$ to $(q, \tilde{t}_4)$ and $\tilde{t}_1 \le \tilde{t}_2 < \tilde{t}_3 \le \tilde{t}_4$ then we say that $(\mathcal{C}_g, T, \mathbf{t})$ has an h-strip (r-strip) from $(q, \tilde{t}_2)$ to $(q, \tilde{t}_3)$ (or from $(q, \tilde{t}_3)$ to $(q, \tilde{t}_2)$).
For a branching $\mathbf{v_i}$ and a $q \in B_{b_i}(\mathrm{pin}(\mathbf{v_i}))$ we call the maximal h-strip (if any) from (q, t_i) to (q, t) with $t_i < t$ ($t_i > t$) a **downwards going (upwards going) h-strip associated to the branching**. (Note that in our pictures the positively oriented time-axis goes downwards.) The notation for r-strips is analogous.
$(\mathcal{C}_g, T, \mathbf{t})$ must have a downwards going r-strip at the points $(\mathrm{pin}(\mathbf{v_i}), t_i)$ because of condition 2.b. We call it an **apex-r-strip**.
An h-strip (r-strip) in $(\mathcal{C}_g, T, \mathbf{t})$ **goes to the bottom (to the top)** if the corresponding h-strip (r-strip) in $\mathcal{C}_g$ goes to the bottom (to the top).
We define

$$\tilde{c}(\mathcal{C}_g, T, \mathbf{t}) \stackrel{\mathrm{def}}{=} \prod_H (1 - \exp(\lambda\, \mathrm{length}(H))) \tag{195}$$

where the product is over all maximal h-strips H that do not end in $(\Lambda \setminus \Lambda_1) \times \{0\}$.

We draw in the specific gum configuration in Figure 5 thick horizontal lines for branchings and thin or thick vertical lines for h-strips or r-strips, respectively. There are two branchings of degree 1, at (q_2, t_1) and at (q_3, t_2). The specific gum configuration has, for example, a long r-strip at site q_5, an r-strip from (q_1, t_1) to the top and an h-strip from (q_1, t_1) to the bottom.
Note that the vertices in the labelled gum tree (except the root) are assigned to points in $\mathbb{Z}^d$ (in this example $d = 1$) by the map pin. For example $\mathrm{pin}(\mathbf{v_1}) = q_2$.

Definition 3.8 We denote by $\mathcal{R}_q(t)$ the operator as defined in Definition 3.3, acting on the q-th coordinate. For $\mathcal{C}_g \in \mathrm{Confg}(\Lambda, \Lambda_1)$ with $n \ge 1$ branchings at $\mathbf{v_1} \prec \ldots \prec \mathbf{v_n}$ of degree $b_1, \ldots, b_n$, respectively, we set $t_0 \stackrel{\mathrm{def}}{=} -T$, $t_{n+1} \stackrel{\mathrm{def}}{=} 0$ and define:



Figure 5: Specific gum configuration and its labelled tree

$$\mathrm{Op}_1(i,\mathcal{C}_g,T,\mathbf{t}) \stackrel{\text{def}}{=} \bigotimes_{q\in\Lambda_{\mathcal{Q}}(i,\mathcal{C}_g)} \mathcal{Q}_q \bigotimes_{q\in\Lambda_{\mathcal{R}}(i,\mathcal{C}_g)} \mathcal{R}_q(t_{i+1}-t_i), \tag{196}$$

$$\mathrm{Op}_2(i,k) \stackrel{\text{def}}{=} \mathcal{B}_{\mathrm{pin}(\mathbf{v_i}),k} \bigotimes_{\bar{q}\in\Lambda\setminus\{q\}} \mathrm{id}_{\bar{q}}, \tag{197}$$

$$\mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}} \stackrel{\text{def}}{=} \tilde{c}(\mathcal{C}_g,T,\mathbf{t})\,\mathrm{Op}_1(n,\mathcal{C}_g,T,\mathbf{t})\circ\mathrm{Op}_2(n,b_n)\circ\ldots \tag{198}$$

$$\circ\mathrm{Op}_1(1,\mathcal{C}_g,T,\mathbf{t})\circ\mathrm{Op}_2(1,b_1)\circ\mathrm{Op}_1(0,\mathcal{C}_g,T,\mathbf{t})$$

$$\text{and}\quad \mathcal{L}^T_{\mathcal{C}_g} \stackrel{\text{def}}{=} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t}\; \mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}} \tag{199}$$

where $\Lambda_{\mathcal{Q}}(i,\mathcal{C}_g)$ is the set of $q\in\Lambda$ such that $(\mathcal{C}_g,T,\mathbf{t})$ has an h-strip from (q,t_i) to (q,t_{i+1}) and $\Lambda_{\mathcal{R}}(i,\mathcal{C}_g)$ is the set of $q\in\mathbb{Z}^d$ such that $(\mathcal{C}_g,T,\mathbf{t})$ has an r-strip from (q,t_i) to (q,t_{i+1}).

If $n(\mathcal{C}_g)=0$ we simply set

$$\mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}} \stackrel{\text{def}}{=} \bigotimes_{q:\,\mathrm{long}(q)=0} Q_q \bigotimes_{q:\,\mathrm{long}(q)=1} \mathcal{R}_q \tag{200}$$

$$\mathcal{L}^T_{\mathcal{C}_g} \stackrel{\text{def}}{=} \mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}}. \tag{201}$$

Finally we set

$$\pi_{\Lambda_1}\circ\mathcal{L}^T_{S,\Lambda} \stackrel{\text{def}}{=} \sum_{\mathcal{C}_g\in\mathrm{Conf_g}(\Lambda,\Lambda_1)} \pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g} \tag{202}$$

Remark 3.4 1. If H is a maximal h-strip from time t_i to time t_j with $1\le i<j\le n+1$ then $\mathrm{length}(H)=|t_i-t_j|$ and so the factor $1-\exp(-\lambda\,|t_i-t_j|)$ does not depend on T. However, in the case $i=0$, i.e. $t_i=-T$, the factor $1-\exp(-\lambda\,(T-|t_j|))$ depends on T. For $T=\infty$ this is equal to 1.

2. From (195), (196), (197) and (198) we see that the map $\mathbf{t} \mapsto \pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}}$, defined on $\mathrm{Simplex}(\mathcal{C}_g, T)$, is uniformly continuous (because all factors are uniformly continuous wrt $\mathbf{t}$), hence integrable if $T < \infty$. We will see in the next proposition that the integral also exists in the case $T = \infty$. So (199) is well-defined.

3. We see that if $(\mathcal{C}_g, \infty, \mathbf{t})$ has an r-strip going to the top then $\mathcal{L}^\infty_{\mathcal{C}_g,\mathbf{t}} = 0$.

Proposition 3.1 There are constants $0 < \tilde{\vartheta} < \vartheta < 1$ and a $c_4 > 0$ such that for sufficiently small $\epsilon > 0$, large c_g, all $T > 0$, $\Lambda_1 \subset \Lambda \in \mathcal{F} \setminus \{\emptyset\}$ and $\phi \in \mathcal{H}_\vartheta$

$$\tilde{\vartheta}^{|\Lambda_1|} \sum_{\mathcal{C}_g \in \mathrm{Confg}(\Lambda,\Lambda_1)} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t} \, \|\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g} \phi_\Lambda\| \le c_4 \, \|\phi\|_\vartheta \tag{203}$$

For sufficiently large T this also holds for suitably chosen $\tilde{\vartheta} = \vartheta$.

Proof For each $\mathcal{C}_g \in \mathrm{Conf_g}(\Lambda, \Lambda_1)$ and $\mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, T)$ we get an estimate analogous to (58) in [13]):

$$\begin{aligned} &\|\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}} \phi_\Lambda\|_{\Lambda_1} \qquad (204)\\ &\le \quad (c_3\epsilon)^{|n_\beta|} \exp\left(-c_g \sum_{k=1}^\infty k^d n_{\beta,k}\right) c_h^{\bar{n}_h} c_r^{\bar{n}_r} \\ &\qquad \times \prod_R \exp\left(-(1-\eta)\,\lambda\,\mathrm{length}(R)\right) \tilde{c}(\mathcal{C}_g, T, \mathbf{t}) \, \|\phi_{\Lambda(\mathcal{C}_g)}\|_{\Lambda(\mathcal{C}_g)}, \end{aligned}$$

where the product is over all maximal r-strips R of $(\mathcal{C}_g, T, \mathbf{t})$ and $\tilde{c}(\mathcal{C}_g, T, \mathbf{t})$ is as defined in (195).
Consider a labelled tree τ with parameters $n_{\beta,k}$ and K, a set $\Lambda_2 \subset \Lambda_1$ with $|\Lambda_2| = K$ and the set $A(\tau, \Lambda_2)$ of all $\mathcal{C}_g \in \mathrm{Conf}_g(\Lambda, \Lambda_1)$ whose labelled tree is τ and whose gum tree has parameter Λ_2. Note that there can be different linear orders on the branchings of τ. We want to estimate

$$\tilde{\vartheta}^{|\Lambda_1|} \sum_{\mathcal{C}_g \in A(\tau,\Lambda_2)} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t} \, \|\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}} \phi_\Lambda\| \tag{205}$$

and consider this expression as integral over the union of all sets $\mathrm{Simplex}(\mathcal{C}_g, T)$. We change the variables of integration: Let $\tilde{t}_1, \ldots, \tilde{t}_{|n_\beta|}$ denote the lengths of the apex-r-strips (where $\tilde{t}_i$ corresponds to the branching $\mathbf{v_i}$). They are bounded by T. Let t_i correspond to $\mathbf{v_i}$. (So the t_i's are not necessarily ordered according to their indices now!) For each $\mathbf{t} = (t_1, \ldots, t_{|n_\beta|}) \in \bigcup_{\mathcal{C}_g \in A(\tau)} \mathrm{Simplex}(\mathcal{C}_g, T)$ there is a unique $\tilde{\mathbf{t}} = (\tilde{t}_1, \ldots, \tilde{t}_{|n_\beta|})$. Note that the union is of disjoint sets and its image is a subset of $[0, T]^n$. Further the change of variables from $\mathbf{t}$ to $\tilde{\mathbf{t}}$ is linear and has a determinant of modulus 1. We see that by doing the transformation successively: $\tilde{t}_1$ is given by a linear equation

$$\tilde{t}_1 = \mathrm{Lin}_1(t_2, \ldots, t_n) - t_1 \tag{206}$$

and $\tilde{t}_2$ by

$$\tilde{t}_2 = \mathrm{Lin}_2(\tilde{t}_1, t_3, \ldots, t_n) - t_2 \tag{207}$$

etc. and the statement about the determinant follows. So we can estimate in (205) the term
'$\sum_{\mathcal{C}_g \in A(\tau,A_2)} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} d\mathbf{t} \ldots$' by '$\int_{[0,T]^{|n_\beta(\tau)|}} d\tilde{\mathbf{t}} \ldots$' and so in the estimate of (203) we replace
'$\sum_{\mathcal{C}_g} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} d\mathbf{t} \ldots$' by '$\sum_{\Lambda_2,\tau} \int_{[0,T]^{|n_\beta(\tau)|}} d\tilde{\mathbf{t}} \ldots$' where the sum is over all $\Lambda_2 \subseteq \Lambda_1$ and labelled trees τ with parameters $|\Lambda_2|$ and n_β.
We use the 'usual estimates' similar to the ones in the proof of Proposition 8.1 in [13]): For fixed $\Lambda_2 \subseteq \Lambda_1$ and n_β with $|n_\beta| \ge |\Lambda_2|$ the number of labelled trees with parameter $|\Lambda_2|$ and n_β is bounded

by $4^{|\Lambda_2|}\prod_{k=1}^{\infty}(\exp(\tilde{c}_d k^d))^{n_{\beta,k}}$. For each k-branching we get a factor $\lambda\,\epsilon\exp(-c_g k^d)$ from the uniform estimates for the corresponding operator. Summing over all possible choices of upwards or downwards going h-strips or r-strips associated with the branching, we get a factor $\exp(c_{12}k^d)$. There are not more than $|\Lambda_1|-|\Lambda_2|$ sites for which we can choose between long h-strips and long r-strips. A long r-strip gives rise to a factor $c_r\exp(-(1-\eta)\lambda T)$, and a long h-strip to a factor at most c_h. The norm $\|\phi_{\Lambda(\mathcal{C}_g)}\|_{\Lambda(\mathcal{C}_g)}$ is estimated by

$$\begin{aligned}\|\phi_{\Lambda(\mathcal{C})}\|_{\Lambda(\mathcal{C})} &\leq \vartheta^{-|\Lambda_r|-\sum_{k=1}^{\infty}(3k)^d n_{\beta,k}}\|\phi\|_\vartheta \\ &\leq \vartheta^{-|\Lambda_r|}\prod_{k=1}^{\infty}\vartheta^{-(3k)^d n_{\beta,k}}\|\phi\|_\vartheta.\end{aligned} \tag{208}$$

Gum configurations $\mathcal{C}_g$ without branchings (i.e. $n_\beta(\mathcal{C}_g)=0$) can only have long r-chains (that must end in Λ_1) or long h-chains. This case corresponds to the summand for $K=0$ in (210).
We remark that the sum '$\sum_{n=0}^{\infty}$' in (210) also includes the estimate for the special case $K=0$. Then the gum configurations have no branchings and so $n_\beta=0$. The sum '$\sum_{n=0}^{\infty}$' should then be replaced by a factor 1 (to avoid confusion). However, this sum is at least 1 and so the estimate is correct.
We estimate the left-hand side (lhs) of (203):

$$\tilde{\vartheta}^{|\Lambda_1|}\sum_{\mathcal{C}_g\in\mathrm{Confg}(\Lambda,\Lambda_1)}\int_{\mathrm{Simplex}(\mathcal{C}_g,T)}\lambda^{|n_\beta(\mathcal{C}_g)|}d\mathbf{t}\,\|\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}}\phi_\Lambda\| \tag{209}$$

$$\begin{aligned}&\leq \tilde{\vartheta}^{|\Lambda_1|}\sum_{K=0}^{|\Lambda_1|}\binom{|\Lambda_1|}{K}(c_h+\vartheta^{-1}c_r\exp(-(1-\eta)\lambda T))^{|\Lambda_1|-K}4^K \\ &\quad\times\sum_{n=K}^{\infty}\left(\epsilon\lambda\sum_{k=1}^{\infty}\exp(-c_g k^d)\,\exp((\tilde{c}_d+c_{12})k^d)\,c^{k^d}\vartheta^{-ck^d}\right. \\ &\quad\left.\times\int_0^T dt\,c_r\exp(-(1-\eta)\lambda T)\right)^n\|\phi\|_\vartheta \\ &\leq c_4\left(\tilde{\vartheta}c_h+\frac{\tilde{\vartheta}}{\vartheta}\,c_r\,\exp(-(1-\eta)\lambda T)+\tilde{\vartheta}\epsilon_1\right)^{|\Lambda_1|}\|\phi\|_\vartheta\end{aligned} \tag{210}$$

with $\lim_{\epsilon\to 0}\epsilon_1=0$. So there are $0<\tilde{\vartheta}<\vartheta$ such that for sufficiently small ϵ

$$\tilde{\vartheta}c_h+\frac{\tilde{\vartheta}}{\vartheta}\,c_r\,\exp(-(1-\eta)T)+\tilde{\vartheta}\epsilon_1<1 \tag{211}$$

and so (203) holds uniformly in Λ_1 and Λ. For sufficiently large T we can choose $\tilde{\vartheta}=\vartheta$ such that (211) holds. So (203) is proved. □

Proposition 3.2 *$\mathcal{L}^T_{S,\Lambda}\stackrel{\mathrm{def}}{=}\pi_\Lambda\circ\mathcal{L}^T_{S,\Lambda}$ is the transfer operator, restricted to $\mathcal{H}_\Lambda$, for $K^T_{S,\Lambda}$, i.e.*

$$\int_{(S^1)^\Lambda}d\mu\,(K^T_{S,\Lambda}\psi_\Lambda)\,\phi_\Lambda=\int_{(S^1)^\Lambda}d\mu\,\psi_\Lambda\,(\mathcal{L}^T_{S,\Lambda}\phi_\Lambda) \tag{212}$$

For all $\psi\in\mathcal{C}^0((S^1)^\Lambda)$ and $\phi_\Lambda\in\mathcal{H}_\Lambda$ the operator $\pi_{\Lambda_1}\circ\mathcal{L}^T_{S,\Lambda}$ is the composite of π_{Λ_1} and $\mathcal{L}^T_{S,\Lambda}$ for any $\Lambda_1\subset\Lambda$.

Proof The first claim is a special case of the last statement. Note that $\pi_{\Lambda_1} \circ \mathcal{L}^T_{S,\Lambda}$ is a priori the operator defined in (202). Now we prove that it is actually the composite of π_{Λ_1} and the transfer operator, restricted to $\mathcal{H}_\Lambda$, for $K^T_{S,\Lambda}$. The convergence of the following expressions follows from Proposition 3.1.
We consider all ordered finite sequences of jump-sites in Λ. A particular sequence $(q_1, \ldots, q_k)$ occurs in a fixed time interval $(-T, 0)$ with probability $e^{-|\Lambda|\lambda T} \frac{(\lambda T)^N}{N!}$ because this is the probability of having exactly N jumps in total, divided by the number $|\Lambda|^N$ of different sequences of length k (which have all the same probability).
The sequence corresponds to a map $S_{q_N} \circ \ldots \circ S_{q_1}$ (cf. Definition 3.2) and so by (164), to a transfer operator

$$\pi_{\Lambda_1} \circ \mathcal{L}_{S_{q_N}} \circ \ldots \circ \mathcal{L}_{S_{q_1}} \tag{213}$$

So the composite of π_{Λ_1} and the transfer operator for $K^T_{S,\Lambda}$ is equal to the following sum over all (possibly empty) sequences:

$$\sum_{(q_1,\ldots,q_N)} e^{-|\Lambda|\lambda T} \frac{(\lambda T)^N}{N!} \, \pi_{\Lambda_1} \circ \mathcal{L}_{S_{q_N}} \circ \ldots \circ \mathcal{L}_{S_{q_1}} \tag{214}$$

because this is equal to the rhs of (145). The probability space Ω is partitioned into countably many sets, each corresponding to a particular sequence of jump sites. So we can write the integral here as a weighted sum.
The factors in (213) can be split

$$\mathcal{L}_{S_q} = \left(\mathcal{Q}_q + \mathcal{R}_q + \sum_{l=1}^{\infty} \mathcal{B}_{q,l} \right) \otimes \mathrm{id}_{\Lambda\setminus\{q\}} \tag{215}$$

Expanding the product in (213), we get a sum of operators. Recall the rules (153) and (162), and also that we have

$$\begin{aligned} \pi_q \circ \mathcal{R}_q &= 0 \\ \pi_q \circ \mathcal{B}_{q,l} &= 0, \end{aligned} \tag{216}$$

for all $q \in \Lambda$ and $l \geq 1$ (as $\mathcal{Q}_q = h_q \cdot \pi_q$). So some of the summands in the expansion are zero, namely if $\mathcal{Q}_q$ is followed by $\mathcal{R}_q$, or $\mathcal{R}_q$ or $\mathcal{B}_q$ are followed by $\mathcal{Q}_q$ or a projection at site $\tilde{q} \notin \Lambda_1$ due to the oprator π_{Λ_1}. 'Following' here means that there is in between no $\mathcal{B}_{q,l}$ with $\|\tilde{q} - q\| \leq l$. In the following we rule out these combinations. (Compare this to the notion of non-zero configurations in [13].)
Now we represent each summand

$$\pi_{\Lambda_1} \circ A_N \circ \ldots \circ A_1 \tag{217}$$

in the expansion of (213) which is the composite of operators π_{Λ_1}, $\mathcal{B}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$, $\mathcal{R}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ and $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ with variable $q \in \Lambda$, by a gum configuration as follows:
We define the obvious order of the factors in (217) such that A_1 comes before A_2 etc. Each $\mathcal{B}_{q,l}$ corresponds to a l-branching (which is assigned by the map pin to q) and the order of the operators $\mathcal{B}_{q,l}$ defines the linear order of the branchings. The other factors $\mathcal{Q}_q$ and $\mathcal{R}_q$ determine the h- and r-strips in the following way:

1. *strips between two vertices:* Let $\mathbf{v}_i \prec \mathbf{v}_j$ be two branchings of degree b_i and b_j, respectively, and $q \in B_{b_i}(\mathrm{pin}(\mathbf{v}_i)) \cap B_{b_j}(\mathrm{pin}(\mathbf{v}_j)) \cap \Lambda$ such that there is no other branching $\mathbf{v}_k$, of degree b_k, say, with $\mathbf{v}_i \prec \mathbf{v}_k \prec \mathbf{v}_j$ and $q \in B_{b_k}(\mathrm{pin}(\mathbf{v}_k))$. Then $\mathcal{C}_g$ has an h-strip between $\mathrm{pin}^{-1}_{\mathbf{v}_i}(q)$ and $\mathrm{pin}^{-1}_{\mathbf{v}_j}(q)$ if there is a factor $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ in (217) between the two factors corresponding to $\mathbf{v}_i$ and $\mathbf{v}_j$. Otherwise $\mathcal{C}_g$ has an r-strip there (even if there is no factor $\mathcal{R}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$.)

2. *strips from a vertex and the top:* Let $\mathbf{v}_j$ be a b_j-branching and assume there is no b_i-branching $\mathbf{v}_i$ with $\mathbf{v}_i \prec \mathbf{v}_j$ and $q \in B_{b_i}(\mathrm{pin}(\mathbf{v}_i))$. Then $\mathcal{C}_g$ has an h-strip from $\mathrm{pin}_{\mathbf{v}_j}^{-1}(q)$ to the top if there is a factor $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ in (217) before the factor corresponding to $\mathbf{v}_j$. Otherwise $\mathcal{C}_g$ has an r-strip there.

3. *strips from a vertex and the bottom:* Let $\mathbf{v}_i$ be a b_i-branching and assume there is no b_j-branching $\mathbf{v}_j$ with $\mathbf{v}_i \prec \mathbf{v}_j$ and $q \in B_{b_j}(\mathrm{pin}(\mathbf{v}_j))$. Then $\mathcal{C}_g$ has an h-strip from $\mathrm{pin}_{\mathbf{v}_i}^{-1}(q)$ to the bottom if there is a factor $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ in (217) after the factor corresponding to $\mathbf{v}_j$ or if $q \in \Lambda \setminus \Lambda_1$. Otherwise $\mathcal{C}_g$ has an r-strip there.

4. *long strips:* Let $q \in \Lambda$ and assume that there is no b_i-branching $\mathbf{v}_i$ with $q \in B_{b_i}(\mathrm{pin}(\mathbf{v}_i))$. Then $\mathcal{C}_g$ has a long h-strip at site q if there is a factor $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ in (217) or if $q \in \Lambda \setminus \Lambda_1$. Otherwise $\mathcal{C}_g$ has a long r-strip there.

The assignment of a summand in the sequence (217) of operators to a gum configuration is not injective, as we have already seen in the simple Example 3.1. Now we consider a fixed gum configuration $\mathcal{C}_g$ on Λ ending in Λ_1 and all sequences corresponding to it. We assume $\mathcal{C}_g$ to have at least one branching. The case of no branching is treated in a similar but easier way. To keep the notation simpler, that $\Lambda = \{1, \ldots, |\Lambda|\}$. Any such sequence has the factors $\mathcal{B}_{q_i,l_i}$ $(i = 1, \ldots, K)$ that correspond 1-1 and order preserving to the K branchings of $\mathcal{C}_g$ as described in above. Between two consecutive factors $\mathcal{B}_{q_i,l_i}$ and $\mathcal{B}_{q_{i+1},l_{i+1}}$ there can be factors $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ (factors $\mathcal{R}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$) if for any $\mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, T)$ the specific gum configuration $(\mathcal{C}_g, T, \mathbf{t})$ has an h-strip (an r-strip) from (q, t_i) to (q, t_{i+1}). Similarly, the option of having such factors before $\mathcal{B}_{q_1,l_1}$ or after $\mathcal{B}_{q_K,l_K}$ depends on the h-strips and r-strips of $\mathcal{C}_g$ in the obvious way. Further such factors belong to particular maximal r-strips or h-strips in the obvous way.
Let us denote the number of factors $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ or $\mathcal{R}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ before $\mathcal{B}_{q_1,l_1}$ by $n_{0,q}$ and the number of such factors after $\mathcal{B}_{q_i,l_i}$ by $n_{i,q}$.
Note that for every maximal h-strip to the bottom at a site $q \in \Lambda_1$ there must be at least one factor $Q_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$ because of rule 3 on page 40. We denote this condition on the family $(n_{i,q})_{\substack{0\le i\le K\\ q\in\Lambda}}$ of numbers by 'condition A'. In total there are

$$N = \sum_{q\in\Lambda} \sum_{i=0}^{K} n_{i,q} + K \tag{218}$$

factors that correspond to a particular sequence of N jump sites and such a sequence occurs with probability $e^{-|\Lambda|\lambda T} \frac{(\lambda T)^N}{N!}$. The $\sum_{q\in\Lambda} n_{0,q}$ factors before $\mathcal{B}_{q_1,l_1}$ can occur in any of the $\binom{n_{0,1}+\ldots+n_{0,|\Lambda|}}{n_{0,1},\ldots,n_{0,|\Lambda|}}$ different orders, that all have the same probability.
So if we sum over all (products of) operators that correspond to $\mathcal{C}_g$, weighted with the probability that the corresponding sequence of jumps (in the underlying Poisson process) occurs, we get In (219) each operator $A_{k,q}$ stands for either $\mathcal{Q}_q$ or $\mathcal{R}_q$.

$$\begin{aligned}
&\sum_{\substack{(n_{i,q})\\ \text{condition A}}} e^{-|\Lambda|\lambda T} \frac{(\lambda T)^N}{N!} \binom{n_{0,1}+\ldots+n_{0,|\Lambda|}}{n_{0,1},\ldots,n_{0,|\Lambda|}} \cdots \binom{n_{K,1}+\ldots+n_{K,|\Lambda|}}{n_{K,1},\ldots,n_{K,|\Lambda|}} \qquad (219)\\
&\quad \times A_{K,1}^{n_{K,1}} \otimes \ldots \otimes A_{K,|\Lambda|}^{K,n_{|\Lambda|}} \circ \mathcal{B}_{q_K,l_K} \otimes \mathrm{id}_{\Lambda\setminus\{q_K\}} \circ \ldots \circ \mathcal{B}_{q_1,l_1} \otimes \mathrm{id}_{\Lambda\setminus\{q_1\}} \circ A_{0,1}^{n_{0,1}} \otimes \ldots \otimes A_{0,|\Lambda|}^{n_{0,|\Lambda|}}\\
&= \sum_{\substack{(n_{i,q})\\ \text{condition A}}} \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} \lambda^K d\mathbf{t} \left(e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{K,1}}}{n_{K,1}!} A_{K,1}^{n_{K,1}} \right) \otimes \ldots\\
&\quad \otimes \left(e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{K,|\Lambda|}}}{n_{K,|\Lambda|}!} A_{K,|\Lambda|}^{n_{K,|\Lambda|}} \right) \circ \mathcal{B}_{q_K,l_K} \otimes \mathrm{id}_{\Lambda\setminus\{q_K\}} \circ \ldots \circ \mathcal{B}_{q_1,l_1} \otimes \mathrm{id}_{\Lambda\setminus\{q_1\}}
\end{aligned}$$

$$\circ \left(e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{0,1}}}{n_{0,1}!} A_{0,1}^{n_{0,1}} \right) \otimes \ldots \otimes \left(e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{0,|\Lambda|}}}{n_{0,|\Lambda|}!} A_{0,|\Lambda|}^{n_{0,|\Lambda|}} \right)$$

$$= \int_{\mathrm{Simplex}(\mathcal{C}_g,T)} \lambda^k d\mathbf{t} \prod_H \left(1 - e^{-\lambda \mathrm{length}(H)}\right) \left(\sum_{n_{K,1} \geq 0} e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{K,1}}}{n_{K,1}!} A_{K,1}^{n_{K,1}} \right) \otimes \ldots$$

$$\otimes \left(\sum_{n_{K,|\Lambda|} \geq 0} e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{K,|\Lambda|}}}{n_{K,|\Lambda|}!} A_{K,|\Lambda|}^{n_{K,|\Lambda|}} \right) \circ \mathcal{B}_{q_K,l_K} \otimes \mathrm{id}_{\Lambda \setminus \{q_K\}} \circ \ldots \circ \mathcal{B}_{q_1,l_1} \otimes \mathrm{id}_{\Lambda \setminus \{q_1\}}$$

$$\circ \left(\sum_{n_{0,1} \geq 0} e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{0,1}}}{n_{0,1}!} A_{0,1}^{n_{0,1}} \right) \otimes \ldots \otimes \left(\sum_{n_{0,|\Lambda|} \geq 0} e^{-\lambda(t_1+T)} \frac{(\lambda(t_1+T))^{n_{0,|\Lambda|}}}{n_{0,|\Lambda|}!} A_{0,|\Lambda|}^{n_{0,|\Lambda|}} \right)$$

$$= \mathcal{L}_{\mathcal{C}_g,T}^T \tag{220}$$

We have interchanged summation with and integration and multiplication and applied the following computations to each set of factors belonging to the same maximal h-strip that does not end in $(\Lambda \setminus \Lambda_1) \times \{0\}$. For notational simplicity, let the maximal h-strip under consideration cover the time intervals $(t_0,t_1),\ldots,(t_{l-1},t_l)$. So its length is $t_l - t_0$. And let the operators $\mathcal{Q}_q$ have exponents $n_1,\ldots,n_l$:

$$\sum_{\substack{n_0,\ldots,n_l \geq 0 \\ n_0+\ldots+n_l \geq 1}} \prod_{i=1}^{l} \left(e^{-\lambda(t_i-t_{i-1})} \frac{(\lambda(t_i-t_{i-1}))^{n_i}}{n_i!} \mathcal{Q}_q \right) \tag{221}$$

$$= \prod_{i=1}^{l} \left(\sum_{n_i=0}^{\infty} e^{-\lambda(t_i-t_{i-1})} \frac{(\lambda(t_i-t_{i-1}))^{n_i}}{n_i!} \mathcal{Q}_q \right) - \prod_{i=1}^{l} e^{-\lambda(t_i-t_{i-1})} \mathcal{Q}_q$$

$$= \left(1 - e^{-\lambda(t_l-t_0)}\right) Q_q \tag{222}$$

This explains the appearance of the factor (195) in (220). Recall that each operator $A_{i,q}$ in (220) stands for either $\mathcal{Q}_q$ or $\mathcal{R}_q$. So we can replace

$$\sum_{n_{i,q} \geq 0} e^{-\lambda(t_i-t_{i-1})} \frac{(\lambda(t_i-t_{i-1}))^{n_{i,q}}}{n_{i,q}!} \mathcal{Q}_q^{n_{i,q}} = \mathcal{Q}_q \tag{223}$$

$$\sum_{n_{i,q} \geq 0} e^{-\lambda(t_i-t_{i-1})} \frac{(\lambda(t_i-t_{i-1}))^{n_{i,q}}}{n_{i,q}!} \mathcal{R}_q^{n_{i,q}} = \mathcal{R}_q(t_i-t_{i-1}) \tag{224}$$

and get (220).
So we have seen that (214) is equal to $\pi_{\Lambda_1} \circ \mathcal{L}_{S,\Lambda}^T$, as defined in (202).

□

For the representation of the transfer operator for the infinite dimensional system we need the following definition.

Definition 3.9 Let $\Lambda_1, \Lambda_2 \subseteq \Lambda \in \mathcal{F} \setminus \{\emptyset\}$ and $\mathcal{C}_g \in \mathrm{Confg}(\Lambda, \Lambda_1)$. We say that $\mathcal{C}_\mathbf{g}$ **lies in** $\mathbf{\Lambda_2}$ if $\Lambda(\mathcal{C}_g) \cup \Lambda_1 \subseteq \Lambda_2$. (Recall that $\Lambda(\mathcal{C}_g)$ was defined in (193).) Let both $\mathcal{C}_g \in \mathrm{Confg}(\Lambda, \Lambda_1)$ and $\widetilde{\mathcal{C}_g} \in$

$\mathrm{Confg}(\widetilde{\Lambda}, \Lambda_1)$ lie in $\Lambda \cap \widetilde{\Lambda}$. If further $\mathcal{C}_g$ and $\widetilde{\mathcal{C}}_g$ have the same gum tree with the same linear order and if they have the same r-strips then we say that $\mathcal{C}_g$ is **equivalent** to $\widetilde{\mathcal{C}}_g$. Then we have defined an equivalence relation and further, for $\mathcal{C}_g$ equivalent to $\widetilde{\mathcal{C}}_g$, we have:

$$\begin{aligned} \mathrm{Simplex}(\mathcal{C}_g, T) &= \mathrm{Simplex}(\widetilde{\mathcal{C}}_g, T) \quad \text{for all } T \in (0, \infty], && (225)\\ \pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g, \mathbf{t}} \circ \pi_{\Lambda} &= \pi_{\Lambda_1} \circ \mathcal{L}^T_{\widetilde{\mathcal{C}}_g, \mathbf{t}} \circ \pi_{\widetilde{\Lambda}} \quad \text{for all } \mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, T) && (226)\\ \text{and} \quad \pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g} \circ \pi_{\Lambda} &= \pi_{\Lambda_1} \circ \mathcal{L}^T_{\widetilde{\mathcal{C}}_g} \circ \pi_{\widetilde{\Lambda}}. && (227) \end{aligned}$$

(226) and (227) say that the operators in $L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})$ are the same. We define by $\mathbf{Confg}(\mathbb{Z}^\mathbf{d}, \mathbf{\Lambda_1})$ the set of equivalence classes. Because of (225) and (226) the simplices and operators for each equivalent class can be defined as being equal to the corresponding object for any representative.
We will write $\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g, \mathbf{t}}$ instead of $\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g, \mathbf{t}} \circ \pi_\Lambda$ and $\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g}$ instead of $\pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g} \circ \pi_\Lambda$ for the operators from $\mathcal{H}_\vartheta$ to $\mathcal{H}_{\Lambda_1}$.

Theorem 3.1 *1. There are $0 < \tilde{\vartheta} < \vartheta < 1$ such that for sufficiently small ϵ, large c_g and every $T \in [0, \infty]$ we can define an operator $\mathcal{L}^T_S$ from $\mathcal{H}_\vartheta$ to $\mathcal{H}_{\tilde{\vartheta}}$ by*

$$\pi_{\Lambda_1} \circ \mathcal{L}^T_S \phi = \sum_{\mathcal{C}_g \in \mathrm{Confg}(\mathbb{Z}^d, \Lambda_1)} \pi_{\Lambda_1} \circ \mathcal{L}^T_{\mathcal{C}_g} \phi. \tag{228}$$

There is a $T_0 > 0$ such that for $T \geq T_0$ the operator $\mathcal{L}^T_S$ maps $\mathcal{H}_\vartheta$ into $\mathcal{H}_\vartheta$.
$\mathcal{L}^T_S$ is the transfer operator, restricted to $\mathcal{H}^{bv}_\vartheta$, for the kernel K^T_S, i.e.

$$\int_M d\mu \, (K^T_S \psi) \, \phi = \int_M d\mu \, \psi \, (\mathcal{L}^T_S \phi) \tag{229}$$

for all $\psi \in \mathcal{C}^0(M)$ and $\phi \in \mathcal{H}^{bv}_\vartheta$.

2. The family $(\mathcal{L}^T_S)_{T \geq 0}$ in $L(\mathcal{H}_\vartheta)$ converges exponentially fast to $\mathcal{L}^\infty_S$:

$$\| \mathcal{L}^\infty_S - \mathcal{L}^T_S \|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\tilde{\vartheta}})} \leq c_5 e^{-c_6 T} \tag{230}$$

for some positive constants c_5, c_6. For sufficiently large T estimate (230) holds also in the norm of $L(\mathcal{H}_\vartheta)$. So among the probability measures corresponding to elements in $\mathcal{H}_\vartheta$ there is a unique K^T_S-invariant probability measure ν^* on M, say corresponding to $\nu \in \mathcal{H}_\vartheta$. The operator $\mathcal{L}^\infty_S$ is a projection onto $\mathrm{span}\, \nu$*:*

$$\mathcal{L}^\infty_S \phi = \mu(\phi) \, \nu. \tag{231}$$

Proof The infinite sum on the rhs of (228) converges as the prove of estimate (203) applies literally to the case $\Lambda = \mathbb{Z}^d$. Next we want to show that $\pi_{\Lambda_1} \circ \mathcal{L}^T_S$ is the limit of $\pi_{\Lambda_1} \circ \mathcal{L}^T_{S,\Lambda}$ (as $\Lambda \to \mathbb{Z}^d$). The difference between these two operators is due to configurations $\mathcal{C}_g$ in $\mathrm{Confg}(\Lambda, \Lambda_1)$ or in $\mathrm{Confg}(\mathbb{Z}^d, \Lambda_1)$ with $\Lambda(\mathcal{C}_g) \not\subset \Lambda$. For these we can split in estimate (204) the factor that arises from the decay of interaction in the following way (which is the same as the splitting (110) in [13]).

$$\exp\left(-c_g \sum_{k=1}^\infty k^d n_{\beta,k} \right) \leq \exp\left(-\tilde{c}_g \sum_{k=1}^\infty k^d n_{\beta,k} \right) \exp(-\xi \, \mathrm{dist}(\Lambda_1, \Lambda^C)) \tag{232}$$

with a suitably chosen $\xi > 0$ such that $\tilde{c}_g = c_g - \xi > 0$. (Note that we can choose ξ so small that the estimates, formerly done with c_g work with $\tilde{c}_g$ instead as well.) So we can estimate

$$
\begin{aligned}
&\tilde{\vartheta}^{|\Lambda_1|}\|\pi_{\Lambda_1}\circ\mathcal{L}^T_{S,\Lambda}-\pi_{\Lambda_1}\circ\mathcal{L}^T_S\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (233)\\
&\quad\le\quad 2\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf}(\mathbb{Z}^d,\Lambda_1),\\ \Lambda(\mathcal{C}_g)\not\subset\Lambda}}\tilde{\vartheta}^{|\Lambda_1|}\|\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})}\\
&\quad\le\quad c_7\,\exp(-\xi\,\mathrm{dist}(\Lambda_1,\Lambda^C)).
\end{aligned}
$$

Next we show (229) for the special case that ψ depends only on the Λ_1-coordinates, using (212):

$$
\begin{aligned}
&\int_M d\mu(\mathbf{z})\;(K^T_S\psi)(\mathbf{z})\,\phi(\mathbf{z}) \qquad (234)\\
&\quad=\quad\lim_{\Lambda\to\mathbb{Z}^d}\int_M d\mu^\Lambda(\mathbf{z}_\Lambda)\;(K^T_{S,\Lambda}\psi)(\mathbf{z}_\Lambda)\,\phi_\Lambda(\mathbf{z}_\Lambda)\\
&\quad=\quad\lim_{\Lambda\to\mathbb{Z}^d}\int_{(S^1)^\Lambda} d\mu^\Lambda(\mathbf{z}_{\mathbf{\Lambda}})\;\psi(\mathbf{z}_\Lambda)\;(\mathcal{L}^T_{S,\Lambda}\phi_\Lambda)(\mathbf{z}_\Lambda)\\
&\quad=\quad\lim_{\Lambda\to\mathbb{Z}^d}\int_{(S^1)^{\Lambda_1}} d\mu^{\Lambda_1}(\mathbf{z}_{\mathbf{\Lambda}})\;\psi(\mathbf{z}_{\Lambda_1})\;(\pi_{\Lambda_1}\circ\mathcal{L}^T_s\phi_\Lambda)(\mathbf{z}_{\Lambda_1})\\
&\quad=\quad\int_M d\mu(\mathbf{z})\;\psi(\mathbf{z})\;(\mathcal{L}^T_S\phi)(\mathbf{z}).
\end{aligned}
$$

We conclude (229) for general $\psi\in\mathcal{C}^0(M)$ by approximating it by ψ_{Λ_1} (cf. (139)), depending only on the Λ_1-coordinates and using continuity wrt ψ of both sides of (229). So 1. is proved.
Next we show (230). We note that for $\Lambda_1=\emptyset$ the lhs (235) in the following estimate is equal to zero as both transfer operators preserve the Lebesgue integral (μ is a 'left eigenvector' with eigenvalue 1.) So we only have to consider the case $|\Lambda_1|\ge 1$.

$$
\begin{aligned}
&\tilde{\vartheta}^{|\Lambda_1|}\|\pi_{\Lambda_1}\circ\mathcal{L}^\infty_S-\pi_{\Lambda_1}\circ\mathcal{L}^T_S\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (235)\\
&\quad\le\quad\tilde{\vartheta}^{|\Lambda_1|}\sum_{\mathcal{C}_g\in\mathrm{Conf}(\mathbb{Z}^d,\Lambda_1)}\|\pi_{\Lambda_1}\circ\mathcal{L}^\infty_{\mathcal{C}_g}-\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})}\\
&\quad\le\quad\tilde{\vartheta}^{|\Lambda_1|}\,\|\mathcal{Q}_{\Lambda_1}\circ\pi_{\Lambda_1}-(1-e^{-\lambda T})^{|\Lambda_1|}\mathcal{Q}_{\Lambda_1}\circ\pi_{\Lambda_1}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (236)\\
&\qquad+\tilde{\vartheta}^{|\Lambda_1|}\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1),\\ \mathcal{C}_g\text{reaches the top}}}\|\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (237)\\
&\qquad+\tilde{\vartheta}^{|\Lambda_1|}\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1),\\ \mathcal{C}_g\text{does not}\\ \text{reach the top},\\ |n_\beta(\mathcal{C}_g)|\ge 1}}\int_{\mathrm{Simplex}(\mathcal{C}_g,\frac{T}{2})}\lambda^{|n_\beta(\mathcal{C}_g)|}d\mathbf{t}\;\|\pi_{\Lambda_1}\circ\mathcal{L}^\infty_{\mathcal{C}_g,\mathbf{t}}-\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (238)\\
&\qquad+\tilde{\vartheta}^{|\Lambda_1|}\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1),\\ \mathcal{C}_g\text{does not reach the top},\\ |n_\beta(\mathcal{C}_g)|\ge 1}}\int_{\substack{\mathrm{Simplex}(\mathcal{C}_g,\infty)\\ \setminus\mathrm{Simplex}(\mathcal{C}_g,\frac{T}{2})}}\lambda^{|n_\beta(\mathcal{C}_g)|}d\mathbf{t}\;\|\pi_{\Lambda_1}\circ\mathcal{L}^\infty_{\mathcal{C}_g,\mathbf{t}}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (239)\\
&\qquad+\tilde{\vartheta}^{|\Lambda_1|}\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1),\\ \mathcal{C}_g\text{does not reach the top},\\ |n_\beta(\mathcal{C}_g)|\ge 1}}\int_{\substack{\mathrm{Simplex}(\mathcal{C}_g,T)\\ \setminus\mathrm{Simplex}(\mathcal{C}_g,\frac{T}{2})}}\lambda^{|n_\beta(\mathcal{C}_g)|}d\mathbf{t}\;\|\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g,\mathbf{t}}\|_{L(\mathcal{H}_\vartheta,\mathcal{H}_{\Lambda_1})} \qquad (240)
\end{aligned}
$$

We have distinguished between the following classes of gum configurations. The first summand (236) corresponds to the operator $\pi_{\Lambda_1}\circ\mathcal{L}^\infty_{\mathcal{C}_g}-\pi_{\Lambda_1}\circ\mathcal{L}^T_{\mathcal{C}_g}$ where $\mathcal{C}_g$ is the gum configuration that has only long

h-strips (no branchings or r-strips). The second summand (237) takes all $\mathcal{C}_g$ into account that reach the top. So all specified configurations $(\mathcal{C}_g, T, \mathbf{t})$ have an r-strip ending at time $-T$. All $(\mathcal{C}_g, \infty, \mathbf{t})$ have an infinitely long r-strip and so the corresponding operator is zero (cf. Remark 3.4.3) and does not appear in (237). The last three summands, (238), (239) and (240), correspond to $\mathcal{C}_g$ that do not reach the top and do not consist only of h-strips. That implies that it has at least one branching and the corresponding domains of integration, $\mathrm{Simplex}(\mathcal{C}_g, \infty)$ and $\mathrm{Simplex}(\mathcal{C}_g, T)$, are not degenerated to a point. We divide them into $\mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})$ and the particular complements. The reason for this will become clear when we do the estimates. In (238) we integrate the norm of the operator difference $\pi_{\Lambda_1} \circ \mathcal{L}^{\infty}_{\mathcal{C}_g} - \pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g}$ over $\mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})$ and in (239) and (240) we integrate the norms of the two operators separately over the particular complement sets.
Now we estimate each summand: The first summand (236) is estimated by

$$\begin{aligned} \tilde{\vartheta}^{|\Lambda_1|} c_h^{|\Lambda_1|} (1 - (1 - e^{-\lambda T})^{|\Lambda_1|}) &\leq (\tilde{\vartheta}\, c_h)^{|\Lambda_1|} \sum_{k=1}^{|\Lambda_1|} \binom{|\Lambda_1|}{k} (e^{-\lambda T})^k \\ &\leq (\tilde{\vartheta}\, c_h\, (1 + e^{-\frac{1}{2}\lambda T}))^{|\Lambda_1|} e^{-\frac{1}{2}\lambda T} \\ &\leq e^{-\frac{1}{2}\lambda T} \end{aligned} \tag{241}$$

where the last inequality holds if $\tilde{\vartheta}$ is chosen sufficiently small.
For estimating the last summand (240) we note that for $\mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, T) \setminus \mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})$ the sum of the lengths of all r-strips of $(\mathcal{C}_g, T, \mathbf{t})$ is at least $\frac{T}{2}$. (This is because $(\mathcal{C}_g, T, \mathbf{t})$ has a branching, say at time t_i with $|t_i| \geq \frac{T}{2}$ and there must be a sequence of apex-r-strips whose lengths add up to at least $\frac{T}{2}$.) So if we split in the estimate (168) for each maximal r-chain, of length t say, the rhs

$$\begin{aligned} \|\mathcal{R}(t)\| &\leq c_r e^{-(1-\eta)\lambda t} \\ &= c_r e^{-\frac{1-\eta}{2}\lambda t}\, e^{-\frac{1-\eta}{2}\lambda t} \end{aligned} \tag{242}$$

we can extract the second factor $\exp(-\frac{1-\eta}{2}\lambda t)$. Their product is bounded from above by $\exp(-\frac{1-\eta}{2}\,\lambda\,\frac{T}{2})$. We assume that the coupling parameters ϵ (small) and c_g (large) are such that our analysis still holds with the (in the sum) remaining factors $\exp(-\frac{1-\eta}{2}\,\lambda\,\mathrm{length}(R))$ for each maximal r-chain R.
We get

$$\tilde{\vartheta}^{|\Lambda_1|} \sum_{\substack{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda_1), \\ \mathcal{C}_g \text{does not reach the top}, \\ |n_\beta(\mathcal{C}_g)| \geq 1}} \int_{\substack{\mathrm{Simplex}(\mathcal{C}_g, T) \\ \setminus \mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})}} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t}\, \|\pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g, \mathbf{t}}\|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})} \leq c_8\, \exp\left(-\lambda \frac{1-\eta}{2}\,\frac{T}{2}\right). \tag{243}$$

Similarly we can estimate the second (237) and the fourth (239) summand:

$$\begin{aligned} \tilde{\vartheta}^{|\Lambda_1|} \sum_{\substack{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda_1), \\ \mathcal{C}_g \text{reaches the top}}} \|\pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g}\|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})} \\ \leq c_9\, \exp\left(-\frac{1-\eta}{2}\,\lambda\, T\right). \end{aligned} \tag{244}$$

$$\begin{aligned} \tilde{\vartheta}^{|\Lambda_1|} \sum_{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda_1)} \int_{\substack{\mathrm{Simplex}(\mathcal{C}_g, \infty) \\ \setminus \mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})}} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t}\, \|\pi_{\Lambda_1} \circ \mathcal{L}^{\infty}_{\mathcal{C}_g, \mathbf{t}}\|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})} \\ \leq c_{10}\, \exp\left(-\frac{1-\eta}{2}\,\lambda\,\frac{T}{2}\right). \end{aligned} \tag{245}$$

We estimate the third summand (238) we use Lemma 3.3 (see below). For that we note that for $\mathcal{C}_g \in \mathrm{Confg}(\mathbb{Z}^d, \Lambda_1)$ and $\mathbf{t} \in \mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})$ the operators $\pi_{\Lambda_1} \circ \mathcal{L}^{\infty}_{\mathcal{C}_g,\mathbf{t}}$ and $\pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g,\mathbf{t}}$ can both be written as a product of numbers $(1 - e^{-\mathrm{length}(H)})$ and operators $\mathcal{Q}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$, corresponding to maximal h-strips, $\mathcal{R}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$, corresponding to maximal r-strips, and $\mathcal{B}_q \otimes \mathrm{id}_{\Lambda\setminus\{q\}}$, corresponding to branchings. They have the same structure in the sense that these factors are in 1-1-correspondence and the quantitative difference is only due to h-strips going to the top or long h-strips in Λ_1 as we can see from representation (198) for $\mathcal{L}^{\infty}_{\mathcal{C}_g,\mathbf{t}}$ and $\mathcal{L}^{T}_{\mathcal{C}_g,\mathbf{t}}$ and also from Remark 3.4.1. So they differ only in the constants $\tilde{c}(\mathcal{C}_g, \infty, \mathbf{t})$ and $\tilde{c}(\mathcal{C}_g, T, \mathbf{t})$. More precisely, an h-strip in $\mathcal{C}_g$ that goes to the top and therefore corresponds to an h-strip in $(\mathcal{C}_g, T, \mathbf{t})$, say from (q, t_i) to $(q, -T)$, and so gives rise to a factor $1 - \exp(-\lambda(T - |t_i|))$ (note that $|t_i| < \frac{T}{2}$) whilst the corresponding h-strip in $(\mathcal{C}_g, \infty, \mathbf{t})$ ends at time $-\infty$ and gives rise to a factor 1. Similarly a long h-strip of $\mathcal{C}_g$ in Λ_1 gives rise to factors $1 - \exp(-\lambda T)$ and 1, respectively. In both cases the difference between the scalar factors (for each h-strip to the top) is bounded by

$$\delta^2 = \exp\left(-\frac{\lambda}{2}\,T\right). \tag{246}$$

The number of h-strips to the top is bounded by $\sum_{k=1}^{\infty} 3^d n_{\beta,k} k^d$ and the number of long h-strips at sites in Λ_1 by $|\Lambda_1| - K$ (where $n_{\beta,k}$ and K are the parameters of the labelled tree of $\mathcal{C}_g$.)
So we estimate

$$\begin{aligned} &\|\pi_{\Lambda_1} \circ \mathcal{L}^{\infty}_{\mathcal{C}_g,\mathbf{t}} - \pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g,\mathbf{t}}\|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})} \\ &\leq \quad \delta \prod_{k=1}^{\infty} (1+\delta)^{3^d n_{\beta,k} k^d} (1+\delta)^{|\Lambda_1| - K} (c_3\epsilon)^{|n_\beta|} \exp\left(-c_g \sum_{k=1}^{\infty} k^d n_{\beta,k}\right) c_h^{\bar{n}_h} c_r^{\bar{n}_r} \\ &\qquad \times \prod_R \exp(-(1-\eta)\lambda\,\mathrm{length}(R))\; \tilde{c}(\mathcal{C}_g, T, \mathbf{t})\, \|\phi_{\Lambda(\mathcal{C}_g)}\|_{\Lambda(\mathcal{C}_g)}. \end{aligned} \tag{247}$$

The factors $(1+\delta)^{3^d n_{\beta,k} k^d}$ are compensated for by $\exp(-c_g k^d n_{\beta,k})$ 'in the usual way'. If ϵ is sufficiently small and c_g large we can estimate

$$\begin{aligned} \tilde{\vartheta}^{|\Lambda_1|} \sum_{\substack{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda_1), \\ \mathcal{C}_g \text{does not reach the top}, \\ |n_\beta(\mathcal{C}_g)| \geq 1}} & \int_{\mathrm{Simplex}(\mathcal{C}_g, \frac{T}{2})} \lambda^{|n_\beta(\mathcal{C}_g)|} d\mathbf{t}\; \|\pi_{\Lambda_1} \circ \mathcal{L}^{\infty}_{\mathcal{C}_g,\mathbf{t}} - \pi_{\Lambda_1} \circ \mathcal{L}^{T}_{\mathcal{C}_g,\mathbf{t}}\|_{L(\mathcal{H}_\vartheta, \mathcal{H}_{\Lambda_1})} \\ &\leq c_{11}\, e^{-\frac{\lambda}{4}T}. \end{aligned} \tag{248}$$

From (241), (244), (248), (245), and (243) we conclude (230) with $c_6 = \frac{1-\eta}{4}\lambda$ and c_5 sufficiently large. For any $\phi \in \mathcal{H}_\vartheta$ and any $\Lambda \in \mathcal{F}$ we have

$$\begin{aligned} \pi_\Lambda \circ \mathcal{L}^{\infty}_{S}\,\phi \quad &= \sum_{\substack{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda), \\ \mathcal{C}_g \text{does not reach the top}}} \pi_\Lambda \circ \mathcal{L}^{\infty}_{\mathcal{C}_g}\,\phi \\ &= \sum_{\substack{\mathcal{C}_g \in \mathrm{Conf_g}(\mathbb{Z}^d, \Lambda), \\ \mathcal{C}_g \text{does not reach the top}}} \left(\pi_\Lambda \circ \mathcal{L}^{\infty}_{\mathcal{C}_g} h_{\mathbb{Z}^d}\right)\, \mu(\phi) \end{aligned} \tag{249}$$

The sum in (249) is a priori over all $\mathcal{C}_g \in \mathrm{Conf}_g(\mathbb{Z}^d, \Lambda)$ but, as we have seen before, if $\mathcal{C}_g$ reaches the top the corresponding operator $\pi_\Lambda \circ \mathcal{L}^{\infty}_{\mathcal{C}_g}$ is zero. If $\mathcal{C}_g$ does not reach the top there are only h-strips going to the top $(-\infty)$ and $\pi_\Lambda \circ \mathcal{L}^{\infty}_{\mathcal{C}_g}$ is a projection onto $\mathrm{span}(h_{\mathbb{Z}^d})$.

We set $\nu_\Lambda \stackrel{\text{def}}{=} \pi_\Lambda \circ \mathcal{L}_{\mathcal{C}_g}^\infty h_{\mathbb{Z}^d}$ and this defines $\nu = (\nu_\Lambda)_{\Lambda \in \mathcal{F}}$. Note that the transfer operator $\mathcal{L}_S^\infty$ preserves the Lebesgue-integral and so $\nu_\emptyset = 1$, i.e. ν corresponds to a probability measure.

□

In the proof of Theorem 3.1 we have used the following lemma.

Lemma 3.3 *Let $A_1, \ldots, A_n, \tilde{A}_1, \ldots, \tilde{A}_n$ be operators on the same Banach space, $0 < \delta < 1$ and $a_1, \ldots, a_n$ positive numbers such that:*

$$\|A_i\| \leq a_i \qquad \text{for all } 1 \leq i \leq n \tag{250}$$

$$\text{and} \quad \|A_i - \tilde{A}_i\| \leq \delta^2 a_i. \tag{251}$$

Then

$$\|A_1 \circ \ldots \circ A_n - \tilde{A}_1 \circ \ldots \circ \tilde{A}_n\| \leq \delta \, (1+\delta)^n \, a_1 \cdot \ldots \cdot a_n. \tag{252}$$

Proof From (251) we get

$$\|\tilde{A}_i\| \leq (1 + \delta^2) \, a_i. \tag{253}$$

So we get via 'telescope expansion':

$$\begin{aligned}
&\|A_1 \circ \ldots \circ A_n - \tilde{A}_1 \circ \ldots \circ \tilde{A}_n\| \qquad (254)\\
&\leq \|A_1 \circ \ldots \circ A_n - \tilde{A}_1 \circ A_2 \circ \ldots \circ A_n\| + \ldots \\
&\quad + \|\tilde{A}_1 \circ \ldots \circ \tilde{A}_{n-1} \circ A_n - \tilde{A}_1 \circ \ldots \circ \tilde{A}_n\| \\
&\leq \delta^2 \, (1 + (1+\delta^2) + \ldots + (1+\delta^2)^{n-1}) \, a_1 \cdot \ldots \cdot a_n \\
&= ((1+\delta^2)^n - 1) \, a_1 \cdot \ldots \cdot a_n \\
&= \sum_{k=1}^{n} \binom{n}{k} \delta^{2k} \, a_1 \cdot \ldots \cdot a_n \\
&\leq \delta \sum_{k=1}^{n} \binom{n}{k} \delta^{k} \, a_1 \cdot \ldots \cdot a_n \\
&\leq \delta (1+\delta)^n \, a_1 \cdot \ldots \cdot a_n
\end{aligned}$$

and the lemma is proved.

□

Remark 3.5 Analogously to Proposition 6.1.3 in [13], one can also prove a semigroup-like property of the family $(\mathcal{L}_S^t)_{t \geq 0}$, using Remark 2.5 and the diagrammatic representation of the operators.

4 Decay of Correlations

In the following theorem which is completely analogous to Theorem 7.1 in [13], we state the mixing properties for the invariant probability measure ν^* in terms of the weighted norms.

Theorem 4.1 *For sufficiently small ϑ, $\tilde{\vartheta}$, ϵ and big c_g there is a $\kappa \in (0,1)$ and positive constants c_{12}, c_{13}, c_{14} and c_{15} such that for all finite disjoint $\Lambda_1, \Lambda_2 \subset \mathbb{Z}^d$ and $\psi \in \mathcal{H}_{\Lambda_2}$ the following holds:*

$$\|\nu_{\Lambda_1\cup\Lambda_2} - \nu_{\Lambda_1}\nu_{\Lambda_2}\|_{\Lambda_1\cup\Lambda_2} \leq c_{12}\,\vartheta^{|\Lambda_1\cup\Lambda_2|}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}, \tag{255}$$

$$\|\pi_{\Lambda_1}(\psi\nu) - \nu(\psi)\,\nu_{\Lambda_1}\|_{\Lambda_1} \leq c_{13}\,\vartheta^{-|\Lambda_1\cup\Lambda_2|}\|\psi\|_{\Lambda_2}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}, \tag{256}$$

$$\begin{aligned}\|\pi_{\Lambda_1}\circ\mathcal{L}_S^T(\psi\nu) - \nu(\psi)\,\nu_{\Lambda_1}\|_{\Lambda_1} &\leq c_{14}\,\vartheta^{-|\Lambda_2|}\,\tilde{\vartheta}^{-|\Lambda_1|}\|\psi\|_{\Lambda_2}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)} \\ &\quad\times\exp\left(-c_{15}T\right)\end{aligned} \tag{257}$$

for every $T>0$.

Proof For a gum configuration $\mathcal{C}_g$ we define in analogy to (109) in [13]

$$b(\mathcal{C}_g) \stackrel{\mathrm{def}}{=} \sum_{k=1}^{\infty} k\,n_{\beta,k}(\mathcal{C}_g). \tag{258}$$

In the following we split gum configurations $\mathcal{C}_g \in \mathrm{Conf}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2)$ with $b(\mathcal{C}_g)\leq\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)$ into $\mathcal{C}_g=\mathcal{C}_g^1\cup\mathcal{C}_g^2$ with $\mathcal{C}_g^1\in\mathrm{Conf}(\mathbb{Z}^d,\Lambda_1)$, $\mathcal{C}_g^2\in\mathrm{Conf}(\mathbb{Z}^d,\Lambda_2)$ and $\Lambda(\mathcal{C}_g^1)\cap\Lambda(\mathcal{C}_g^2)=\emptyset$.
We write, using (231) and the notation of (156):

$$\begin{aligned}\nu_{\Lambda_1\cup\Lambda_2} &= \sum_{\substack{\mathcal{C}_g=\mathcal{C}_g^1\cup\mathcal{C}_g^2\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)\leq\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} (\pi_{\Lambda_1}\circ\mathcal{L}_{\mathcal{C}_g^1}^T h_{\mathbb{Z}^d})\,(\pi_{\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g^2}^T h_{\mathbb{Z}^d}) \\ &\quad+\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)>\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} \pi_{\Lambda_1\cup\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g}^T h_{\mathbb{Z}^d}\end{aligned}$$

In estimating the norm of the second summand in (259) we can take out from the estimate for $\|\pi_{\Lambda_1}\circ\mathcal{L}_{\mathcal{C}_g}^{\infty}h_{\mathbb{Z}^d}\|$ a factor

$$\exp\left(-\xi\,\frac{1}{2}\,\mathrm{dist}(\Lambda_1,\Lambda_2)\right) = \kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)} \tag{259}$$

like in (232) such that we get

$$\|\sum_{\substack{\mathcal{C}_g\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)>\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} \pi_{\Lambda_1\cup\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g}^T h_{\mathbb{Z}^d}\| \leq c_{16}\,\vartheta^{|\Lambda_1\cup\Lambda_2|}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}. \tag{260}$$

We write the first summand in (259) as

$$\begin{aligned}&\sum_{\substack{\mathcal{C}_g=\mathcal{C}_g^1\cup\mathcal{C}_g^2\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)\leq\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} (\pi_{\Lambda_1}\circ\mathcal{L}_{\mathcal{C}_g^1}^T h_{\mathbb{Z}^d})\,(\pi_{\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g^2}^T h_{\mathbb{Z}^d}) \\ &= \nu_{\Lambda_1}\nu_{\Lambda_2} - \sum_{\substack{\mathcal{C}_g=\mathcal{C}_g^1\cup\mathcal{C}_g^2\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)>\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} (\pi_{\Lambda_1}\circ\mathcal{L}_{\mathcal{C}_g^1}^T h_{\mathbb{Z}^d})\,(\pi_{\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g^2}^T h_{\mathbb{Z}^d})\end{aligned} \tag{261}$$

and estimate

$$\|\sum_{\substack{\mathcal{C}_g=\mathcal{C}_g^1\cup\mathcal{C}_g^2\in\mathrm{Conf_g}(\mathbb{Z}^d,\Lambda_1\cup\Lambda_2),\\ b(\mathcal{C}_g)>\frac{1}{2}\mathrm{dist}(\Lambda_1,\Lambda_2)}} (\pi_{\Lambda_1}\circ\mathcal{L}_{\mathcal{C}_g^1}^T h_{\mathbb{Z}^d})\,(\pi_{\Lambda_2}\circ\mathcal{L}_{\mathcal{C}_g^2}^T h_{\mathbb{Z}^d})\| \leq c_{17}\,\vartheta^{|\Lambda_1\cup\Lambda_2|}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}. \tag{262}$$

From (260), (261) and (262) we conclude (255). The proof of (256), using (255), is the same as in [13]. To prove (257) we set $\phi = \psi\nu - \nu(\psi)\,\nu$. So

$$\pi_{\Lambda_1} \circ \mathcal{L}_S^T \phi = \pi_{\Lambda_1} \circ \mathcal{L}_S^T (\psi\,\nu) - \nu(\psi)\,\nu_{\Lambda_1} \tag{263}$$

and in particular

$$\mathcal{L}_S^\infty \phi = 0. \tag{264}$$

We estimate (263), analogously to (129) in [13], using the finer estimate

$$\begin{aligned} &\|\phi_{\Lambda(\mathcal{C})}\|_{\Lambda(\mathcal{C})} \\ &\quad \leq \quad c_{13}\,\vartheta^{-|\Lambda_2|}\,\|\psi\|_{\Lambda_2}\,\vartheta^{-|\Lambda_r(\mathcal{C})|-\sum_{k=1}^\infty (3k)^d n_{\beta,k}}\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)-\sum_{k=1}^\infty k n_{\beta,k}} \end{aligned} \tag{265}$$

that we get from (256). For each $\mathcal{C}_g$ we get a 'good' factor $\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}$ that we can take out of the sum (over gum configurations), and a 'bad' factor $\kappa^{-\sum_{k=1}^\infty k n_{\beta,k}}$. The latter is compensated for in the usual way by the factor $\exp(-c_g \sum_{k=1}^\infty k^d n_{\beta,k})$, provided that c_g is sufficiently large.
Using (264) and (265), we get with the same argument as for the proof of (230):

$$\tilde{\vartheta}^{|\Lambda_1|}\,\|\pi_{\Lambda_1} \circ \mathcal{L}_S^T \phi\| \leq c_{18}\,\vartheta^{-|\Lambda_2|}\,\|\psi\|\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}\,\exp(-c_{15}T) \tag{266}$$

and (257) follows.

□

We can state the mixing properties of ν^* wrt spatio-temporal shifts in terms of correlation functions for observables $\psi_1, \psi_2 \in \mathcal{C}^0(M)$ like in Theorem 2.2 of [13] and, using Theorem 4.1, prove them in exactly the same way. We denote by τ a spatial shift and by $m(\tau)$ its size. (Cf. [13].)

Theorem 4.2 *For sufficiently small ϑ, ϵ and large c_g there is a $\kappa \in (0,1)$ such that for all nonempty $\Lambda_1, \Lambda_2 \in \mathcal{F}$ the following holds with the constant $c(\Lambda_1,\Lambda_2,\kappa) \stackrel{\mathrm{def}}{=} \kappa^{-\max\{\|p-q\|:p\in\Lambda_1,q\in\Lambda_2\})}$ and some positive constants c_{19}, c_{20}:*

1. *If $\psi_1 \in \mathcal{C}((S^1)^{\Lambda_1})$ and $\psi_2 \in \mathcal{C}((S^1)^{\Lambda_2})$ then*

$$\begin{aligned} &\left|\int_M d\nu^*\,\psi_1\,\psi_2 - \left(\int_M d\nu^*\,\psi_1\right)\left(\int_M d\nu^*\,\psi_2\right)\right| \\ &\quad \leq \quad c_{19}\,\vartheta^{-|\Lambda_1|-|\Lambda_2|}\,\|\psi_1\|_\infty\,\|\psi_2\|_\infty\,\kappa^{\mathrm{dist}(\Lambda_1,\Lambda_2)}. \end{aligned} \tag{267}$$

2. *If $\psi_1 \in \mathcal{C}((S^1)^{\Lambda_1})$ and $\psi_2 \in \mathcal{H} \cap \mathcal{C}((S^1)^{\Lambda_2})$ then*

$$\begin{aligned} &\left|\int_M d\nu^*\,K_S^T(\psi_1 \circ \tau)\,\psi_2 - \left(\int_M d\nu^*\,\psi_1 \circ \tau\right)\left(\int_M d\nu^* \psi_2\right)\right| \\ &\quad \leq \quad c(\Lambda_1,\Lambda_2,\kappa)\,c_{20}^{|\Lambda_1|+|\Lambda_2|}\,\|\psi_1\|_\infty\,\|\psi_2\|_{\Lambda_2}\,\kappa^{m(\tau)}\,\exp(-c_{21}T). \end{aligned} \tag{268}$$

3. *If the system is translation-invariant and $\psi_1, \psi_2 \in \mathcal{C}(M)$ then*

$$\lim_{\max\{m(\tau),T\}\to\infty} \int_M d\nu^*\,K_S^T(\psi_1 \circ \tau)\,\psi_2 \quad = \quad \left(\int_M d\nu^*\,\psi_1\right)\left(\int_M d\nu^*\,\psi_2\right). \tag{269}$$

□

Acknowledgements

This paper is based on a part of my thesis [12]. I would like to thank my supervisor Hans Henrik Rugh for his great support, and also Roger Tribe for many helpful discussions and suggestions. Further I thank Viviane Baladi and Véronique Maume-Deschamps for helpful discussions. Finally I thank the EC for financial support through the TMR-Fellowship ERBFMBICT-961157.

References

[1] L. Arnold: Random Dynamical Systems, *Springer*, 1998

[2] V. Baladi, M. Degli Esposti, S. Isola, E. Järvenpää, A. Kupiainen: The spectrum of weakly coupled map lattices, *J. Math. Pures et Appl.*, 77, pp. 539-584, 1998

[3] H. Bauer: Wahrscheinlichkeitstheorie, *de Gruyter*, 4th ed., 1991

[4] J. Bricmont and A. Kupiainen: Coupled Analytic Maps, *Nonlinearity* 8, pp.379-396, 1995

[5] J. Bricmont and A. Kupiainen: High Temperature Expansions and Dynamical Systems, *Comm. Math. Phys.* 178, 1996, pp.73-732

[6] J. Bricmont and A. Kupiainen: Infinite dimensional SRB-measures. Lattice dynamics (Paris, 1995), *Physica D*, 103 (1997), no. 1-4, 18–33.

[7] L.A. Bunimovich: Coupled map lattices: One step forward and two steps back, *Physica D 86*, 1995, pp. 248-255

[8] L.A. Bunimovich, Y.G. Sinai: Space-time chaos in coupled map lattices, *Nonlinearity 1*, pp. 491-516, 1988

[9] R.L. Dobrushin: Markov processes with a large number of locally interacting components: existence of a limit process and its ergodicity, *Problems Inform. Transmission*, 7, pp. 149-164, 1971

[10] R.L. Dobrushin: Markov processes with many locally interacting components- the reversible case and some generalizations, *Problems Inform. Transmission*, 7, pp. 235-241, 1971

[11] R.L. Dobrushin and Ya.G. Sinai (editors): Multicomponent Random Systems, *Advances in probability and related topics*, vol. 6 (originally published in Russian), 1980

[12] T. Fischer: Transfer Operators for Deterministic and Stochastic Coupled Map Lattices, *thesis*, University of Warwick, 1998

[13] T. Fischer, H.H. Rugh: Transfer Operators for Coupled Analytic Maps, to appear in *Ergodic Theory & Dynamical Systems*

[14] R.J. Glauber: Time-dependent statistics of the Ising model. *J. Math. Physics*, 4. pp. 294-307, 1963

[15] T.E. Harris: Nearest-Neighbor Markov Interaction Processes on Multidimensional Lattices, *Advances in Mathematics* **9**, pp. 66-89, 1972

[16] R. Holley: A class of interactions in an infinite particle system, *Adv. Math.*, 5, pp. 291-309, 1970

[17] M. Jiang:Equilibrium states for lattice models of hyperbolic type *Nonlinearity 8*, no.5, 1994, pp.631-659

[18] M. Jiang: Ergodic Properties of Coupled Map Lattices of Hyperbolic Type, *Penns. State University Dissertation*, 1995

[19] M. Jiang and A. Mazel: uniqueness of Gibbs States and exponential Decay of Correlation for some Lattice models, *Journal of Statistical Physics* 82, no. 3-4, 1995

[20] M. Jiang and Ya.B. Pesin: Equilibrium Measures for Coupled Map Lattices: Existence, Uniqueness and Finite-Dimensional Approximations, to appear in CMP

[21] G. Keller and M. Künzle: Transfer Operators for Coupled Map Lattices, *Ergodic Theory & Dynamical Systems* Vol. 12, pp.297-318, 1992

[22] S. Lang: Real and Functional Analysis, 3rd edition, *Springer*, 1993

[23] T.M. Liggett: Existence theorems for infinite particle systems, *Trans. Amer. Math. Soc.*, 165, pp. 471-481, 1972

[24] T.M. Liggett: Interacting Particle Systems, *Grundlehren der mathematischen Wissenschaften*, 276, *Springer*, 1985

[25] C. Maes and A. Van Moffaert: Stochastic Stability of Weakly Coupled Lattice Maps, *Nonlinearity*, 10, 1997, pp.715-730

[26] Ya.B. Pesin and Ya.G. Sinai: Space-time chaos in chains of weakly interacting hyperbolic mappings, *Advances in Soviet Mathematics* Vol.3, 1991, pp.165-198

[27] F. Spitzer: Random Processes Defined through the Interaction of an Infinite Particle System, *Springer Lecture Notes in Mathematics*, Vol. 89, pp. 201-223, 1969

[28] F. Spitzer: Interaction of Markov processes, *Adv. Math.*, 5, pp. 246-290, 1970

[29] D.L. Volevich: The Sinai-Bowen-Ruelle Measure for a Multidimensional Lattice of interacting Hyperbolic Mappings, *Russ. Akad. Dokl. Math.* 47, 1993, pp.117-121

[30] D.L. Volevich: Construction of an analogue of Bowen-Ruell-Sinai measure for a multidimensional lattice of interacting hyperbolic mappings *Russ. Akad. Math. Sbornik* Vol.79, 1994, pp.347-363

Torsten Fischer
Niels Bohr Institute
Blegdamsvej 17
DK-2100 Copenhagen Ø
email: fischer@nbi.dk